

# ABATACEPT IV

## Products Affected

- ORENCIA (WITH MALTOSE)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST.
Coverage Duration	RA, PJIA, PSA: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. AGVHD: 12 MONTHS.
Other Criteria	INITIAL: RA: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PJIA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ IR. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, TREMFYA, XELJANZ, RINVOQ, SKYRIZI. RENEWAL: RA, PJIA, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ABATACEPT SQ

## Products Affected

- ORENCIA
- ORENCIA CLICKJECT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: RA: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PJIA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ IR. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, COSENTYX, ENBREL, TREMFYA, XELJANZ, RINVOQ, SKYRIZI. RENEWAL: RA, PJIA, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ABEMACICLIB

---

## Products Affected

- VERZENIO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ABIRATERONE

---

## Products Affected

- *abiraterone*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC HIGH-RISK CASTRATION-SENSITIVE PROSTATE CANCER (MCSPC), METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (MCRPC): 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ABIRATERONE SUBMICRONIZED

---

## Products Affected

- YONSA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ACALABRUTINIB

---

## Products Affected

- CALQUENCE (ACALABRUTINIB MAL)

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT BRUKINSA, WHERE INDICATIONS ALIGN.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# ADAGRASIB

---

## Products Affected

- KRAZATI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ADALIMUMAB

## Products Affected

- HUMIRA PEN
- HUMIRA PEN CROHNS-UC-HS START
- HUMIRA PEN PSOR-UVEITS-ADOL HS
- HUMIRA SUBCUTANEOUS SYRINGE KIT 40 MG/0.8 ML
- HUMIRA(CF)
- HUMIRA(CF) PEDI CROHNS STARTER
- HUMIRA(CF) PEN
- HUMIRA(CF) PEN CROHNS-UC-HS
- HUMIRA(CF) PEN PEDIATRIC UC
- HUMIRA(CF) PEN PSOR-UV-ADOL HS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, OR GENITAL AREA.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RA, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA), ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST. UVEITIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OPHTHALMOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PJIA, PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD. PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. CD, UC: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPURINE, METHOTREXATE, OR MESALAMINE. UVEITIS: NO ISOLATED ANTERIOR UVEITIS. RENEWAL: RA, PJIA, PSA, AS, PSO, HIDRADENITIS SUPPURATIVA, UVEITIS: CONTINUES TO BENEFIT FROM THE MEDICATION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# AFATINIB

---

## Products Affected

- GILOTRIF

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS.
<b>Other Criteria</b>	METASTATIC NSCLC WITH EGFR MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ALECTINIB

---

## Products Affected

- ALECENSA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ALPELISIB-PIQRAY

---

## Products Affected

- PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1), 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2)

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# AMBRISENTAN

---

## Products Affected

- *ambrisentan*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	PAH: INITIAL: DOES NOT HAVE IDIOPATHIC PULMONARY FIBROSIS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# AMIVANTAMAB-VMJW

---

## Products Affected

- RYBREVANT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# ANAKINRA

---

## Products Affected

- KINERET

PA Criteria	Criteria Details
Exclusion Criteria	CORONAVIRUS DISEASE 2019 (COVID-19) IN HOSPITALIZED ADULTS.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	RA: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. ALL OTHERS: 12 MONTHS.
Other Criteria	RA: INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, RINVOQ, ENBREL, XELJANZ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# APALUTAMIDE

---

## Products Affected

- ERLEADA ORAL TABLET 240 MG, 60 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: NON-METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (NMCRPC): HIGH RISK PROSTATE CANCER (I.E., RAPIDLY INCREASING PROSTATE SPECIFIC ANTIGEN [PSA] LEVELS). NMCRPC, METASTATIC CASTRATION-SENSITIVE PROSTATE CANCER (MCSPC): 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. RENEWAL: NMCRPC, MCSPC: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GNRH ANALOG.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# APOMORPHINE

---

## Products Affected

- *apomorphine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PARKINSONS DISEASE (PD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	PD: INITIAL: PHYSICIAN HAS OPTIMIZED DRUG THERAPY FOR PD. RENEWAL: IMPROVEMENT WITH MOTOR FLUCTUATIONS DURING OFF EPISODES WHILE ON THERAPY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# APOMORPHINE - SL

---

## Products Affected

- KYNMOBI SUBLINGUAL FILM 10 MG, 10-15-20-25-30 MG, 15 MG, 20 MG, 25 MG, 30 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	PARKINSONS DISEASE (PD): INITIAL: 18 YEARS OF AGE OR OLDER.
Prescriber Restrictions	PD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	PD: INITIAL: PHYSICIAN HAS OPTIMIZED DRUG THERAPY FOR PARKINSONS DISEASE. RENEWAL: IMPROVEMENT WITH MOTOR FLUCTUATIONS DURING OFF EPISODES WITH THE USE OF THERAPY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# APREMILAST

---

## Products Affected

- OTEZLA
- OTEZLA STARTER

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: MILD PLAQUE PSORIASIS (PSO): ONE OF THE FOLLOWING: 1) PSORIASIS COVERING 2 PERCENT OF BODY SURFACE AREA (BSA), 2) STATIC PHYSICIAN GLOBAL ASSESSMENT (SPGA) SCORE OF 2, OR 3) PSORIASIS AREA AND SEVERITY INDEX (PASI) SCORE OF 2 TO 9. MODERATE TO SEVERE PSO: 1) PSORIASIS COVERING 3 PERCENT OR MORE OF BSA, OR 2) PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, OR GENITAL AREA.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. BEHCETS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, TREMFYA, XELJANZ, RINVOQ, SKYRIZI. MILD PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL SYSTEMIC AGENT (E.G., METHOTREXATE, ACITRETIN, CYCLOSPORINE) AND ONE CONVENTIONAL TOPICAL AGENT (E.G., PUVA, UVB, TOPICAL CORTICOSTEROIDS). MODERATE TO SEVERE PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, SKYRIZI, TREMFYA. BEHCETS DISEASE: 1) HAS ORAL ULCERS OR A HISTORY OF RECURRENT ORAL ULCERS BASED ON CLINICAL SYMPTOMS, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE OR MORE CONSERVATIVE TREATMENTS (E.G., COLCHICINE, TOPICAL CORTICOSTEROID, ORAL CORTICOSTEROID). RENEWAL: PSA, PSO, BEHCETS DISEASE: CONTINUES TO BENEFIT FROM THE MEDICATION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ASCIMINIB

---

## Products Affected

- SCEMBLIX ORAL TABLET 20 MG, 40 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	PHILADELPHIA CHROMOSOME-POSITIVE CHRONIC MYELOID LEUKEMIA (Ph+ CML); MUTATIONAL ANALYSIS PRIOR TO INITIATION AND SCEMBLIX IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ASFOTASE ALFA

---

## Products Affected

- STRENSIQ

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	HYPOPHOSPHATASIA (HPP): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST, GENETICIST, OR METABOLIC SPECIALIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.



PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: PERINATAL/INFANTILE-ONSET HPP: 1) 6 MONTHS OF AGE OR YOUNGER AT ONSET OF HPP, AND 2) POSITIVE FOR A TISSUE NON-SPECIFIC ALKALINE PHOSPHATASE (TNSALP) (ALPL) GENE MUTATION AS CONFIRMED BY GENETIC TESTING OR TWO OF THE FOLLOWING: (A) SERUM ALKALINE PHOSPHATASE (ALP) LEVEL BELOW THAT OF NORMAL RANGE FOR PATIENT AGE, (B) ELEVATED SERUM PYRIDOXAL-5'-PHOSPHATE (PLP) LEVELS AND NO VITAMIN B6 SUPPLEMENTATION IN THE PREVIOUS WEEK, (C) URINE PHOSPHOETHANOLAMINE (PEA) LEVEL ABOVE THAT OF NORMAL RANGE FOR PATIENT AGE, (D) RADIOGRAPHIC EVIDENCE OF HPP, (E) AT LEAST TWO OF THE FOLLOWING: (I) RACHITIC CHEST DEFORMITY, (II) CRANIOSYNOSTOSIS, (III) DELAY IN SKELETAL GROWTH RESULTING IN DELAY OF MOTOR DEVELOPMENT, (IV) HISTORY OF VITAMIN B6 DEPENDENT SEIZURES, (V) NEPHROCALCINOSIS OR HISTORY OF ELEVATED SERUM CALCIUM, (VI) HISTORY OR PRESENCE OF NON-TRAUMATIC POSTNATAL FRACTURE AND DELAYED FRACTURE HEALING. JUVENILE-ONSET HPP: 1) 18 YEARS OF AGE OR YOUNGER AT ONSET OF HPP, AND 2) POSITIVE FOR A TNSALP ALPL GENE MUTATION AS CONFIRMED BY GENETIC TESTING OR TWO OF THE FOLLOWING: (A) SERUM ALP LEVEL BELOW THAT OF NORMAL RANGE FOR PATIENT AGE, (B) ELEVATED SERUM PLP LEVELS AND NO VITAMIN B6 SUPPLEMENTATION IN THE PREVIOUS WEEK, (C) URINE PEA LEVEL ABOVE THAT OF NORMAL RANGE FOR PATIENT AGE, (D) RADIOGRAPHIC EVIDENCE OF HPP, (E) AT LEAST TWO OF THE FOLLOWING: (I) RACHITIC DEFORMITIES, (II) PREMATURE LOSS OF PRIMARY TEETH PRIOR TO 5 YEARS OF AGE, (III) DELAY IN SKELETAL GROWTH RESULTING IN DELAY OF MOTOR DEVELOPMENT, (IV)</p>

PA Criteria	Criteria Details
	<p>HISTORY OR PRESENCE OF NON-TRAUMATIC FRACTURES OR DELAYED FRACTURE HEALING. ALL INDICATIONS: 1) NOT CURRENTLY RECEIVING TREATMENT WITH A BISPHTHOSPHONATE, 2) CALCIUM OR PHOSPHATE LEVELS ARE NOT BELOW THE NORMAL RANGE, 3) NOT HAVE A TREATABLE FORM OF RICKETS. RENEWAL: ALL INDICATIONS: 1) IMPROVEMENT IN THE SKELETAL CHARACTERISTICS OF HPP, AND 2) NOT CURRENTLY RECEIVING TREATMENT WITH A BISPHTHOSPHONATE.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ATOGEPANT

## Products Affected

- QULIPTA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	MIGRAINE PREVENTION: INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREVENTIVE MIGRAINE TREATMENTS: DIVALPROEX SODIUM, TOPIRAMATE, PROPRANOLOL, TIMOLOL, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. RENEWAL: 1) REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# AVAPRITINIB

---

## Products Affected

- AYVAKIT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# AVATROMBOPAG

## Products Affected

- DOPTELET (10 TAB PACK)
- DOPTELET (15 TAB PACK)
- DOPTELET (30 TAB PACK)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: CHRONIC LIVER DISEASE (CLD): PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST, GASTROENTEROLOGIST, HEPATOLOGIST, IMMUNOLOGIST, ENDOCRINOLOGIST, OR A SURGEON. CHRONIC IMMUNE THROMBOCYTOPENIA (ITP): PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR IMMUNOLOGIST.
Coverage Duration	CLD: 1 MONTH. CHRONIC ITP: INITIAL: 2 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: CLD: 1) PLANNED PROCEDURE 10 TO 13 DAYS AFTER INITIATION OF DOPTELET, AND 2) NOT RECEIVING OTHER THROMBOPOIETIN RECEPTOR AGONISTS (E.G., ROMIPLOSTIM, ELTROMBOPAG, ETC.). CHRONIC ITP: TRIAL OF OR CONTRAINDICATION TO CORTICOSTEROIDS OR IMMUNOGLOBULINS OR INSUFFICIENT RESPONSE TO SPLENECTOMY. RENEWAL: CHRONIC ITP: PATIENT HAD A CLINICAL RESPONSE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# AXITINIB

---

## Products Affected

- INLYTA ORAL TABLET 1 MG, 5 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# AZACITIDINE

---

## Products Affected

- ONUREG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# AZTREONAM INHALED

---

## Products Affected

- CAYSTON

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	7 YEARS OF AGE OR OLDER
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# BARICITINIB

## Products Affected

- OLUMIANT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	CORONAVIRUS DISEASE 2019 (COVID-19) IN HOSPITALIZED ADULTS.
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. SEVERE ALOPECIA AREATA: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST.
<b>Coverage Duration</b>	RA, SEVERE ALOPECIA AREATA: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO ANY TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ. SEVERE ALOPECIA AREATA: 1) AT LEAST 50 PERCENT SCALP HAIR LOSS AS MEASURED BY THE SEVERITY OF ALOPECIA TOOL (SALT) FOR MORE THAN 6 MONTHS, AND 2) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGICS FOR SEVERE ALOPECIA AREATA OR OTHER JAK INHIBITORS FOR ANY INDICATION. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION. SEVERE ALOPECIA AREATA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGICS FOR SEVERE ALOPECIA AREATA OR OTHER JAK INHIBITORS FOR ANY INDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No

# BEDAQUILINE

---

## Products Affected

- SIRTURO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	24 WEEKS
<b>Other Criteria</b>	PULMONARY MULTI-DRUG RESISTANT TUBERCULOSIS (MDR-TB): SIRTURO USED IN COMBINATION WITH AT LEAST 3 OTHER ANTIBIOTICS FOR THE TREATMENT OF PULMONARY MDR-TB.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BELIMUMAB

---

## Products Affected

- BENLYSTA SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: SYSTEMIC LUPUS ERYTHEMATOSUS (SLE): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. LUPUS NEPHRITIS (LN): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR NEPHROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: SLE: CURRENTLY TAKING CORTICOSTEROIDS, ANTIMALARIALS, NSAIDS, OR IMMUNOSUPPRESSIVE AGENTS. RENEWAL: SLE: PATIENT HAD CLINICAL IMPROVEMENT. LN: IMPROVEMENT IN RENAL RESPONSE FROM BASELINE LABORATORY VALUES (I.E., EGFR OR PROTEINURIA) AND/OR CLINICAL PARAMETERS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# BELUMOSUDIL

---

## Products Affected

- REZUROCK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BELZUTIFAN

---

## Products Affected

- WELIREG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BENDAMUSTINE

---

## Products Affected

- *bendamustine intravenous recon soln*
- BENDAMUSTINE INTRAVENOUS SOLUTION
- BENDEKA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# BENRALIZUMAB

---

## Products Affected

- FASENRA
- FASENRA PEN

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	ASTHMA: INITIAL: BLOOD EOSINOPHIL LEVEL GREATER THAN OR EQUAL TO 150 CELLS/MCL WITHIN THE PAST 12 MONTHS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	ASTHMA: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE.
<b>Coverage Duration</b>	INITIAL: 4 MONTHS, RENEWAL: 12 MONTHS.



PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>ASTHMA: INITIAL: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE, OR MAXIMALLY TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND ONE OTHER MAINTENANCE MEDICATION, AND 2) ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS, OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA, AND 3) NO CONCURRENT USE WITH XOLAIR, DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA. RENEWAL: 1) NO CONCURRENT USE WITH XOLAIR, DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA, 2) CONTINUED USE OF ICS AND ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) INCREASE IN PERCENT PREDICTED FEV1 FROM PRETREATMENT BASELINE, OR (D) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BETAINE

---

## Products Affected

- *betaine*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BEVACIZUMAB-ADCD

---

## Products Affected

- VEGZELMA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BEVACIZUMAB-AWWB

---

## Products Affected

- MVASI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BEVACIZUMAB-BVZR

---

## Products Affected

- ZIRABEV

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BEXAROTENE

---

## Products Affected

- *bexarotene*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BINIMETINIB

---

## Products Affected

- MEKTOVI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BORTEZOMIB

---

## Products Affected

- *bortezomib injection*
- VELCADE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# BOSENTAN

## Products Affected

- *bosentan*
- TRACLEER ORAL TABLET FOR SUSPENSION

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	PAH: INITIAL: 1) DOES NOT HAVE ELEVATED LIVER ENZYMES (ALT, AST) MORE THAN 3 TIMES UPPER LIMIT OF NORMAL (ULN) OR INCREASE IN BILIRUBIN BY 2 OR MORE TIMES ULN, AND 2) NOT CONCURRENTLY TAKING CYCLOSPORINE A OR GLYBURIDE. RENEWAL: NOT CONCURRENTLY TAKING CYCLOSPORINE A OR GLYBURIDE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BOSUTINIB

---

## Products Affected

- BOSULIF ORAL CAPSULE 100 MG, 50 MG
- BOSULIF ORAL TABLET 100 MG, 400 MG, 500 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	PREVIOUSLY TREATED (Ph+ CML): MUTATIONAL ANALYSIS PRIOR TO INITIATION AND BOSULIF IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BRIGATINIB

---

## Products Affected

- ALUNBRIG ORAL TABLET 180 MG, 30 MG, 90 MG
- ALUNBRIG ORAL TABLETS,DOSE PACK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# C1 ESTERASE INHIBITOR-CINRYZE

## Products Affected

- CINRYZE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	HEREDITARY ANGIOEDEMA (HAE): INITIAL: DIAGNOSIS CONFIRMED BY COMPLEMENT TESTING.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	HAE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST, IMMUNOLOGIST, OR ALLERGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	HAE: INITIAL: NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS. RENEWAL: 1) IMPROVEMENT IN HAE ATTACKS (I.E., REDUCTIONS IN ATTACK FREQUENCY OR ATTACK SEVERITY), AND 2) NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# C1 ESTERASE INHIBITOR-HAEGARDA

## Products Affected

- HAEGARDA SUBCUTANEOUS  
RECON SOLN 2,000 UNIT, 3,000 UNIT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	HEREDITARY ANGIOEDEMA (HAE): INITIAL: DIAGNOSIS CONFIRMED BY COMPLEMENT TESTING.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	HAE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST, IMMUNOLOGIST, OR ALLERGIST.
<b>Coverage Duration</b>	INITIAL AND RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	HAE: INITIAL: NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS. RENEWAL: 1) IMPROVEMENT COMPARED TO BASELINE IN HAE ATTACKS (I.E., REDUCTIONS IN ATTACK FREQUENCY OR ATTACK SEVERITY), AND 2) NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# CABOZANTINIB CAPSULE

---

## Products Affected

- COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X1), 140 MG/DAY(80 MG X1-20 MG X3), 60 MG/DAY (20 MG X 3/DAY)

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# CABOZANTINIB TABLET

---

## Products Affected

- CABOMETYX ORAL TABLET 20 MG, 40 MG, 60 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# CANNABIDIOL

## Products Affected

- EPIDIOLEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	DRAVET SYNDROME (DS), LENNOX-GASTAUT SYNDROME (LGS), TUBEROUS SCLEROSIS COMPLEX (TSC): PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL: 12 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: LENNOX-GASTAUT SYNDROME (LGS): TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING ANTIEPILEPTIC MEDICATIONS: RUFINAMIDE, FELBAMATE, CLOBAZAM, TOPIRAMATE, LAMOTRIGINE, CLONAZEPAM. RENEWAL: DS, LGS, TSC: CONFIRMATION OF DIAGNOSIS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# CAPIVASERTIB

---

## Products Affected

- TRUQAP

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# CAPLACIZUMAB YHDP

## Products Affected

- CABLIVI INJECTION KIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	ACQUIRED THROMBOTIC THROMBOCYTOPENIA PURPURA (ATTP): PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST
Coverage Duration	12 MONTHS
Other Criteria	ATTP: CABLIVI WAS PREVIOUSLY INITIATED AS PART OF THE FDA APPROVED TREATMENT REGIMEN IN COMBINATION WITH PLASMA EXCHANGE AND IMMUNOSUPPRESSIVE THERAPY WITHIN AN INPATIENT SETTING. THE PATIENT HAS NOT EXPERIENCED MORE THAN TWO RECURRENCES OF ATTP WHILE ON CABLIVI THERAPY (I.E., NEW DROP IN PLATELET COUNT REQUIRING REPEAT PLASMA EXCHANGE DURING 30 DAYS POST-PLASMA EXCHANGE THERAPY [PEX] AND UP TO 28 DAYS OF EXTENDED THERAPY).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# CAPMATINIB

---

## Products Affected

- TABRECTA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# CARGLUMIC ACID

## Products Affected

- *carglumic acid*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: ACUTE OR CHRONIC HYPERAMMONEMIA (HA) DUE TO N ACETYLGLUTAMATE SYNTHASE (NAGS) DEFICIENCY: NAGS GENE MUTATION IS CONFIRMED BY BIOCHEMICAL OR GENETIC TESTING. ACUTE HA DUE TO PROPIONIC ACIDEMIA (PA): 1) CONFIRMED BY ELEVATED METHYLCITRIC ACID AND NORMAL METHYLMALONIC ACID, OR 2) GENETIC TESTING CONFIRMS MUTATION IN THE PCCA OR PCCB GENE. ACUTE HA DUE TO METHYLMALONIC ACIDEMIA (MMA): 1) CONFIRMED BY ELEVATED METHYLMALONIC ACID, METHYLCITRIC ACID, OR 2) GENETIC TESTING CONFIRMS MUTATION IN THE MMUT, MMA, MMAB OR MMADHC GENES.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	ACUTE HA DUE TO NAGS/PA/MMA: 7 DAYS. CHRONIC HA DUE TO NAGS: INITIAL: 6 MOS, RENEWAL: 12 MOS.
<b>Other Criteria</b>	RENEWAL: CHRONIC HA DUE TO NAGS: PATIENT HAS SHOWN CLINICAL IMPROVEMENT.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# CERITINIB

---

## Products Affected

- ZYKADIA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# CETUXIMAB

---

## Products Affected

- ERBITUX

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# CLADRIBINE

## Products Affected

- MAVENCLAD (10 TABLET PACK)
- MAVENCLAD (4 TABLET PACK)
- MAVENCLAD (5 TABLET PACK)
- MAVENCLAD (6 TABLET PACK)
- MAVENCLAD (7 TABLET PACK)
- MAVENCLAD (8 TABLET PACK)
- MAVENCLAD (9 TABLET PACK)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL/RENEWAL: 48 WEEKS.
Other Criteria	RELAPSING FORM OF MULTIPLE SCLEROSIS (MS): INITIAL: HAS NOT RECEIVED A TOTAL OF TWO YEARS OF MAVENCLAD TREATMENT (I.E., TWO YEARLY TREATMENT COURSES OF TWO CYCLES IN EACH). RENEWAL: 1) HAS DEMONSTRATED CLINICAL BENEFIT COMPARED TO PRE-TREATMENT BASELINE, 2) DOES NOT HAVE LYMPHOPENIA, AND 3) HAS NOT RECEIVED A TOTAL OF TWO YEARS OF MAVENCLAD TREATMENT (I.E., TWO YEARLY TREATMENT COURSES OF TWO CYCLES IN EACH).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# CLOBAZAM-SYMPAZAN

---

## Products Affected

- SYMPAZAN

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	LENNOX-GASTAUT SYNDROME (LGS): THERAPY IS PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	LGS: 1) UNABLE TO TAKE TABLETS OR SUSPENSIONS, AND 2) TRIAL OF OR CONTRAINDICATION TO A FORMULARY VERSION OF CLOBAZAM.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# COBIMETINIB

---

## Products Affected

- COTELLIC

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# COLCHICINE

---

## Products Affected

- *colchicine oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	PROPHYLAXIS OF GOUT FLARES: 16 YEARS AND OLDER
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PROPHYLAXIS OF GOUT FLARES: TRIAL OF OR CONTRAINDICATION TO COLCHICINE CAPSULES (MITIGARE) IF AGE 18 YEARS OR OLDER.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# CORTICOTROPIN

## Products Affected

- ACTHAR
- CORTROPHIN GEL

PA Criteria	Criteria Details
Exclusion Criteria	INITIAL: NOT APPROVED FOR DIAGNOSTIC PURPOSES.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS AND MULTIPLE SCLEROSIS (MS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, ALLERGIST/IMMUNOLOGIST, OPHTHALMOLOGIST, PULMONOLOGIST OR NEPHROLOGIST.
Coverage Duration	INFANTILE SPASMS AND MS: 28 DAYS. OTHER FDA APPROVED INDICATIONS: INITIAL AND RENEWAL: 28 DAYS
Other Criteria	INITIAL: ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS: TRIAL OF OR CONTRAINDICATION TO INTRAVENOUS (IV) CORTICOSTEROIDS. ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS AND MS: TRIAL OF OR CONTRAINDICATION TO A STANDARD OF CARE THERAPY. RENEWAL: ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS AND MS: 1) DEMONSTRATED CLINICAL BENEFIT WHILE ON THERAPY AS INDICATED BY SYMPTOM RESOLUTION AND/OR NORMALIZATION OF LABORATORY TESTS, AND 2) CONTINUES TO POSSESS CONTRAINDICATION TO IV CORTICOSTEROIDS. PART B BEFORE PART D STEP THERAPY, APPLIES ONLY TO BENEFICIARIES IN AN MA-PD PLAN.
Indications	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# CRIZOTINIB

---

## Products Affected

- XALKORI ORAL CAPSULE

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# CRIZOTINIB PELLETS

---

## Products Affected

- XALKORI ORAL PELLETT 150 MG, 20 MG, 50 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	NON-SMALL CELL LUNG CANCER (NSCLC), ANAPLASTIC LARGE CELL LYMPHOMA (ALCL), INFLAMMATORY MYOFIBROBLASTIC TUMOR (IMT): UNABLE TO SWALLOW CAPSULES.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# CYSTEAMINE HYDROCHLORIDE

---

## Products Affected

- CYSTARAN

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DABRAFENIB

---

## Products Affected

- TAFINLAR ORAL CAPSULE

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# DABRAFENIB SUSPENSION

---

## Products Affected

- TAFINLAR ORAL TABLET FOR SUSPENSION

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	UNABLE TO SWALLOW TAFINILAR CAPSULES.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DACOMITINIB

---

## Products Affected

- VIZIMPRO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC): NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DALFAMPRIDINE

---

## Products Affected

- *dalfampridine*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	MULTIPLE SCLEROSIS (MS): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
<b>Coverage Duration</b>	INITIAL: 3 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	MS: INITIAL: HAS SYMPTOMS OF A WALKING DISABILITY SUCH AS MILD TO MODERATE BILATERAL LOWER EXTREMITY WEAKNESS OR UNILATERAL WEAKNESS PLUS LOWER EXTREMITY OR TRUNCAL ATAXIA. RENEWAL: IMPROVEMENT IN WALKING ABILITY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DAROLUTAMIDE

## Products Affected

- NUBEQA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	INITIAL: NON-METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (NMCRPC): HIGH RISK PROSTATE CANCER (I.E., RAPIDLY INCREASING PROSTATE SPECIFIC ANTIGEN [PSA] LEVELS). NMCRPC, METASTATIC HORMONE-SENSITIVE PROSTATE CANCER (MHSPC): 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. RENEWAL: NMCRPC, MHSPC: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GNRH ANALOG.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DASATINIB

---

## Products Affected

- SPRYCEL ORAL TABLET 100 MG, 140 MG, 20 MG, 50 MG, 70 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PREVIOUSLY-TREATED PHILADELPHIA CHROMOSOME-POSITIVE CHRONIC MYELOID LEUKEMIA (Ph+ CML): MUTATIONAL ANALYSIS PRIOR TO INITIATION AND SPRYCEL IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# DECITABINE/CEDAZURIDINE

---

## Products Affected

- INQOVI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DEFERASIROX

## Products Affected

- *deferasirox*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: CHRONIC IRON OVERLOAD DUE TO BLOOD TRANSFUSIONS: SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 1000 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS). NON-TRANSFUSION DEPENDENT THALASSEMIA (NTDT): 1) SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 300 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS) AND 2) LIVER IRON CONCENTRATION (LIC) OF 5 MG FE/G DRY WEIGHT OR GREATER. RENEWAL: CHRONIC IRON OVERLOAD DUE TO BLOOD TRANSFUSIONS: SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 500 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS). NTDT: 1) SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 300 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS) OR 2) LIC OF 3 MG FE/G DRY WEIGHT OR GREATER.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: CHRONIC IRON OVERLOAD: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR HEMATOLOGIST/ONCOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
<b>Other Criteria</b>	INITIAL FOR ALL INDICATIONS: FORMULARY VERSION OF DEFERASIROX SPRINKLE: TRIAL OF OR CONTRAINDICATION TO GENERIC DEFERASIROX TABLET OR TABLET FOR ORAL SUSPENSION.
<b>Indications</b>	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# DEFERIPRONE

## Products Affected

- *deferiprone*
- FERRIPROX (2 TIMES A DAY)
- FERRIPROX ORAL SOLUTION

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	TRANSFUSIONAL IRON OVERLOAD: RENEWAL: SERUM FERRITIN LEVELS CONSISTENTLY ABOVE 500 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	TRANSFUSIONAL IRON OVERLOAD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR HEMATOLOGIST/ONCOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
<b>Other Criteria</b>	INITIAL: TRANSFUSIONAL IRON OVERLOAD DUE TO THALASSEMIA SYNDROMES: 1) TRIAL OF, CONTRAINDICATION, INTOLERABLE TOXICITIES, OR CLINICALLY SIGNIFICANT ADVERSE EFFECTS TO A FORMULARY VERSION OF DEFERASIROX OR DEFEROXAMINE, OR 2) CURRENT CHELATION THERAPY (I.E., FORMULARY VERSION OF DEFERASIROX OR DEFEROXAMINE) IS INADEQUATE. TRANSFUSIONAL IRON OVERLOAD DUE TO SICKLE CELL DISEASE OR OTHER ANEMIAS: TRIAL OF OR CONTRAINDICATION TO A FORMULARY VERSION OF DEFERASIROX OR DEFEROXAMINE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No

# DENOSUMAB-XGEVA

---

## Products Affected

- XGEVA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DEUTETRABENAZINE

## Products Affected

- AUSTEDO ORAL TABLET 12 MG, 6 MG, 9 MG
- AUSTEDO XR ORAL TABLET EXTENDED RELEASE 24 HR 12 MG, 24 MG, 6 MG
- AUSTEDO XR TITRATION KIT(WK1-4)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	HUNTINGTON DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR MOVEMENT DISORDER SPECIALIST. TARDIVE DYSKINESIA: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR MOVEMENT DISORDER SPECIALIST.
Coverage Duration	12 MONTHS
Other Criteria	TARDIVE DYSKINESIA: HISTORY OF USING AGENTS THAT CAUSE TARDIVE DYSKINESIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# DICLOFENAC TOPICAL GEL

---

## Products Affected

- *diclofenac sodium topical gel 3%*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DICLOFENAC TOPICAL SOLUTION

---

## Products Affected

- *diclofenac sodium topical solution in metered-dose pump*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	OSTEOARTHRITIS OF THE KNEE: TRIAL OF OR CONTRAINDICATION TO A FORMULARY VERSION OF DICLOFENAC SODIUM 1% TOPICAL GEL AND A FORMULARY VERSION OF DICLOFENAC SODIUM 1.5% TOPICAL DROPS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# DIMETHYL FUMARATE

---

## Products Affected

- *dimethyl fumarate oral capsule, delayed release(dr/lec) 120 mg, 120 mg (14)- 240 mg (46), 240 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# DIROXIMEL FUMARATE

---

## Products Affected

- VUMERITY

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# DOSTARLIMAB-GXLY

---

## Products Affected

- JEMPERLI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DRONABINOL CAPSULE

---

## Products Affected

- *dronabinol*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	NAUSEA AND VOMITING ASSOCIATED WITH CANCER CHEMOTHERAPY: TRIAL OF OR CONTRAINDICATION TO ONE ANTIEMETIC THERAPY. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D FOR THE INDICATION OF NAUSEA AND VOMITING ASSOCIATED WITH CANCER CHEMOTHERAPY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# DROXIDOPA

## Products Affected

- *droxidopa*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	NEUROGENIC ORTHOSTATIC HYPOTENSION (NOH): INITIAL: 1) BASELINE BLOOD PRESSURE READINGS WHILE THE PATIENT IS SITTING AND ALSO WITHIN 3 MINUTES OF STANDING FROM A SUPINE POSITION. 2) A DECREASE OF AT LEAST 20 MMHG IN SYSTOLIC BLOOD PRESSURE OR 10 MMHG DIASTOLIC BLOOD PRESSURE WITHIN THREE MINUTES AFTER STANDING FROM A SITTING POSITION.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	NOH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR CARDIOLOGIST.
<b>Coverage Duration</b>	INITIAL: 3 MONTHS RENEWAL: 12 MONTHS
<b>Other Criteria</b>	NOH: RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DUPILUMAB

## Products Affected

- DUPIXENT PEN
- DUPIXENT SYRINGE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: EOSINOPHILIC ASTHMA: BLOOD EOSINOPHIL LEVEL OF 150 TO 1500 CELLS/MCL WITHIN THE PAST 12 MONTHS. EOSINOPHILIC ESOPHAGITIS (EOE): DIAGNOSIS CONFIRMED BY ESOPHAGOGASTRODUODENOSCOPY (EGD) WITH BIOPSY.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: AD, PN: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST, ALLERGIST OR IMMUNOLOGIST. ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE. CRSWNP: PRESCRIBED BY OR IN CONSULTATION WITH AN OTOLARYNGOLOGIST, ALLERGIST OR IMMUNOLOGIST. EOE: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST, ALLERGIST, OR IMMUNOLOGIST.
<b>Coverage Duration</b>	INITIAL: AD, CRSWNP, EOE, PN: 6 MOS, ASTHMA: 4 MOS. RENEWAL: ALL INDICATIONS: 12 MOS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: AD: 1) AD COVERING AT LEAST 10 PERCENT OF BODY SURFACE AREA OR AD AFFECTING THE FACE, HEAD, NECK, HANDS, FEET, GROIN, OR INTERTRIGINOUS AREAS, 2) INTRACTABLE PRURITUS OR CRACKING/OOZING/BLEEDING OF AFFECTED SKIN, 3) TRIAL OF OR CONTRAINDICATION TO ONE TOPICAL (CORTICOSTEROID, CALCINEURIN INHIBITOR, PDE4 INHIBITOR, OR JAK INHIBITOR), AND 4) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGICS OR JAK INHIBITORS FOR AD. ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY-TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND ONE OTHER MAINTENANCE MEDICATION, 2) ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS, OR ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA, AND 3) NO CONCURRENT USE WITH XOLAIR OR OTHER ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA. CRSWNP: 1) EVIDENCE OF NASAL POLYPS BY DIRECT EXAMINATION, ENDOSCOPY OR SINUS CT SCAN, 2) INADEQUATELY CONTROLLED DISEASE AS DETERMINED BY USE OF SYSTEMIC STEROIDS IN THE PAST 2 YEARS OR ENDOSCOPIC SINUS SURGERY, AND 3) A 56 DAY TRIAL OF ONE TOPICAL NASAL CORTICOSTEROID. PN: 1) CHRONIC PRURITIS (ITCH MORE THAN 6 WEEKS), MULTIPLE PRURIGINOUS LESIONS, AND</p>

PA Criteria	Criteria Details
	<p>HISTORY OR SIGN OF A PROLONGED SCRATCHING BEHAVIOR, 2) TRIAL OF OR CONTRAINDICATION TO ONE TOPICAL (CORTICOSTEROID OR CALCIPOTRIOL). RENEWAL: AD: 1) IMPROVEMENT WHILE ON THERAPY, AND 2) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGICS OR JAK INHIBITORS FOR AD. CRSWNP, EOE: IMPROVEMENT WHILE ON THERAPY. ASTHMA: 1) NO CONCURRENT USE WITH XOLAIR, OR OTHER ANTI-IL5 BIOLOGICS FOR ASTHMA, 2) CONTINUED USE OF ICS AND ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) INCREASE IN PERCENT PREDICTED FEV1 FROM PRETREATMENT BASELINE, OR (D) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS. PN: IMPROVEMENT OR REDUCTION OF PRURITIS OR PRURIGINOUS LESIONS.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DUVELISIB

---

## Products Affected

- COPIKTRA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# EFLORNITHINE

---

## Products Affected

- IWILFIN

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	24 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# ELACESTRANT

---

## Products Affected

- ORSERDU ORAL TABLET 345 MG, 86 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ELAGOLIX

## Products Affected

- ORLISSA ORAL TABLET 150 MG, 200 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS.
<b>Age Restrictions</b>	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: 18 YEARS OF AGE OR OLDER.
<b>Prescriber Restrictions</b>	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS
<b>Other Criteria</b>	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, AND 2) TRIAL OF OR CONTRAINDICATION TO AN NSAID AND A PROGESTIN-CONTAINING PREPARATION. RENEWAL: 1) IMPROVEMENT IN PAIN ASSOCIATED WITH ENDOMETRIOSIS WHILE ON THERAPY, AND 2) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No

# ELEXACAFITOR-TEZACAFITOR-IVACAFITOR

## Products Affected

- TRIKAFTA ORAL TABLETS, SEQUENTIAL

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	CYSTIC FIBROSIS (CF): INITIAL: CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CYSTIC FIBROSIS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CYSTIC FIBROSIS EXPERT.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: LIFETIME.
<b>Other Criteria</b>	CF: RENEWAL: 1) MAINTAINED, IMPROVED, OR DEMONSTRATED LESS THAN EXPECTED DECLINE IN FEV1 OR BODY MASS INDEX (BMI), OR 2) REDUCTION IN NUMBER OF PULMONARY EXACERBATIONS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ELIGLUSTAT

---

## Products Affected

- CERDELGA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ELRANATAMAB-BCMM

---

## Products Affected

- ELREXFIO 44 MG/1.1 ML VIAL OUTER, SUV, P/F
- ELREXFIO SUBCUTANEOUS SOLUTION 40 MG/ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	RELAPSED OR REFRACTORY MULTIPLE MYELOMA: RENEWAL: 1) HAS RECEIVED AT LEAST 24 WEEKS OF TREATMENT WITH ELREXFIO, AND 2) HAS RESPONDED TO TREATMENT (PARTIAL RESPONSE OR BETTER), AND HAS MAINTAINED THIS RESPONSE FOR AT LEAST 2 MONTHS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ELTROMBOPAG

## Products Affected

- PROMACTA ORAL POWDER IN PACKET 12.5 MG, 25 MG
- PROMACTA ORAL TABLET 12.5 MG, 25 MG, 50 MG, 75 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: ITP: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR IMMUNOLOGIST.
Coverage Duration	ITP: INITIAL: 2 MO, RENEWAL: 12 MO. HEPATITIS C, SEVERE APLASTIC ANEMIA: 12 MO.
Other Criteria	INITIAL: PERSISTENT OR CHRONIC IMMUNE THROMBOCYTOPENIA PURPURA (ITP): TRIAL OF OR CONTRAINDICATION TO CORTICOSTEROIDS, IMMUNOGLOBULINS, OR AN INSUFFICIENT RESPONSE TO SPLENECTOMY. ALL INDICATIONS: APPROVAL FOR PROMACTA ORAL SUSPENSION PACKETS REQUIRES A TRIAL OF PROMACTA TABLETS OR PATIENT IS UNABLE TO TAKE TABLET FORMULATION. RENEWAL: ITP: PATIENT HAS SHOWN A CLINICAL RESPONSE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# EMAPALUMAB-LZSG

## Products Affected

- GAMIFANT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS (HLH): INITIAL: 1) A GENETIC TEST IDENTIFYING HLH-ASSOCIATED GENE MUTATION (E.G., PRF1, UNC13D), OR 2) HAS AT LEAST FIVE OF THE FOLLOWING EIGHT DIAGNOSTIC CRITERIA FOR HLH: (A) FEVER, (B) SPLENOMEGALY, (C) CYTOPENIAS (AFFECTING AT LEAST 2 OF 3 CELL LINEAGES), (D) HYPERTRIGLYCERIDEMIA OR HYPOFIBRINOGENEMIA, (E) HEMOPHAGOCYTOSIS IN BONE MARROW OR SPLEEN OR LYMPH NODES AND NO EVIDENCE OF MALIGNANCY, (F) LOW OR ABSENT NATURAL KILLER-CELL ACTIVITY, (G) FERRITIN LEVEL OF 500 MCG/L OR GREATER, (H) SOLUBLE CD25 LEVEL OF 2,400 U/ML OR GREATER.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	HLH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN IMMUNOLOGIST, HEMATOLOGIST, OR ONCOLOGIST.
<b>Coverage Duration</b>	INITIAL AND RENEWAL: 8 WEEKS.



PA Criteria	Criteria Details
<b>Other Criteria</b>	HLH: INITIAL: 1) CONCURRENT THERAPY WITH DEXAMETHASONE, AND 2) ONE OF THE FOLLOWING: (A) HAS REFRACTORY, RECURRENT, OR PROGRESSIVE DISEASE, OR (B) HAD A TRIAL OF OR INTOLERANCE TO CONVENTIONAL HLH THERAPY (I.E., CHEMOTHERAPY, STEROIDS, IMMUNOTHERAPY). RENEWAL: 1) HAS NOT RECEIVED SUCCESSFUL HEMATOPOIETIC STEM CELL TRANSPLANTATION, AND 2) DEMONSTRATED IMPROVED IMMUNE SYSTEM RESPONSE FROM BASELINE (E.G., RESOLUTION OF FEVER, DECREASED SPLENOMEGALY, IMPROVEMENT IN CNS SYMPTOMS, IMPROVED CBC, INCREASED FIBRINOGEN LEVELS, REDUCED D-DIMER, REDUCED FERRITIN, REDUCED SOLUBLE CD25 LEVELS.)
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ENASIDENIB

---

## Products Affected

- IDHIFA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ENCORAFENIB

---

## Products Affected

- BRAFTOVI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ENTRECTINIB

---

## Products Affected

- ROZLYTREK ORAL CAPSULE 100 MG, 200 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ENTRECTINIB PELLETS

---

## Products Affected

- ROZLYTREK ORAL PELLETS IN PACKET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC), SOLID TUMORS: 1) TRIAL OF OR CONTRAINDICATION TO ROZLYTREK CAPSULES MADE INTO AN ORAL SUSPENSION, AND 2) DIFFICULTY OR UNABLE TO SWALLOW CAPSULES.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ENZALUTAMIDE

## Products Affected

- XTANDI ORAL CAPSULE
- XTANDI ORAL TABLET 40 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	INITIAL: NON-METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (NMCRPC): HIGH RISK PROSTATE CANCER (I.E. RAPIDLY INCREASING PROSTATE SPECIFIC ANTIGEN [PSA] LEVELS). NON-METASTATIC CASTRATION-SENSITIVE PROSTATE CANCER (NMCSPC): HIGH RISK FOR METASTASIS (I.E. PSA DOUBLED OVER 9 MONTHS OR LESS). METASTATIC CRPC (MCRPC), NMCRPC, METASTATIC CSPC (MCSPC), NMCSPC : 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. RENEWAL: MCRPC, NMCRPC, MCSPC: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GNRH ANALOG.
Indications	All FDA-approved Indications.
Off Label Uses	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No

# EPCORITAMAB-BYSP

---

## Products Affected

- EPKINLY

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# EPOETIN ALFA-EPBX

## Products Affected

- RETACRIT INJECTION SOLUTION  
10,000 UNIT/ML, 2,000 UNIT/ML,  
20,000 UNIT/2 ML, 20,000 UNIT/ML,  
3,000 UNIT/ML, 4,000 UNIT/ML, 40,000  
UNIT/ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	<p>INITIAL: CHRONIC KIDNEY DISEASE (CKD), ANEMIA RELATED TO ZIDOVUDINE, OR CANCER CHEMOTHERAPY: HEMOGLOBIN LEVEL OF LESS THAN 10G/DL. ELECTIVE, NON-CARDIAC, NON-VASCULAR SURGERY: HEMOGLOBIN LEVEL LESS THAN 13G/DL. RENEWAL: 1) CKD IN ADULTS NOT ON DIALYSIS: (A) HEMOGLOBIN LEVEL IS LESS THAN 10G/DL, OR (B) HEMOGLOBIN LEVEL HAS REACHED 10G/DL AND THE DOSE IS BEING REDUCED/INTERRUPTED TO DECREASE THE NEED FOR BLOOD TRANSFUSIONS. CKD IN PEDIATRIC PATIENTS: (A) HEMOGLOBIN LEVEL IS LESS THAN 10G/DL, OR (B) HEMOGLOBIN LEVEL HAS APPROACHED OR EXCEEDS 12G/DL AND THE DOSE IS BEING REDUCED/INTERRUPTED TO DECREASE THE NEED FOR BLOOD TRANSFUSIONS. ANEMIA RELATED TO ZIDOVUDINE: HEMOGLOBIN LEVEL BETWEEN 10G/DL AND 12G/DL. CANCER CHEMOTHERAPY: 1) HEMOGLOBIN LEVEL OF LESS THAN 10 G/DL, OR 2) THE HEMOGLOBIN LEVEL DOES NOT EXCEED A LEVEL NEEDED TO AVOID RBC TRANSFUSION.</p>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<p>ANEMIA FROM CHEMO/CKD WITHOUT DIALYSIS/ZIDOVUDINE: INITIAL/RENEWAL: 12 MONTHS. SURGERY: 1 MONTH.</p>

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Other Criteria</b>	RENEWAL: CKD: NOT RECEIVING DIALYSIS TREATMENT. THIS DRUG MAY BE EITHER BUNDLED WITH AND COVERED UNDER END STAGE RENAL DISEASE DIALYSIS RELATED SERVICES OR COVERED UNDER MEDICARE D DEPENDING UPON THE CIRCUMSTANCES.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ERDAFITINIB

---

## Products Affected

- BALVERSA ORAL TABLET 3 MG, 4 MG, 5 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ERLOTINIB

---

## Products Affected

- *erlotinib oral tablet 100 mg, 150 mg, 25 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC NSCLC WITH EGFR MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ESKETAMINE

## Products Affected

- SPRAVATO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: TREATMENT-RESISTANT DEPRESSION (TRD), MAJOR DEPRESSIVE DISORDER (MDD): PRESCRIBED BY OR IN CONSULTATION WITH A PSYCHIATRIST.
<b>Coverage Duration</b>	INITIAL: TRD: 3 MONTHS. MDD: 4 WEEKS. RENEWAL: TRD, MDD: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: TRD: 1) NON-PSYCHOTIC, UNIPOLAR DEPRESSION, 2) NO ACTIVE SUBSTANCE ABUSE, AND 3) ADEQUATE TRIAL (AT LEAST 4 WEEKS) OF AT LEAST TWO ANTIDEPRESSANT AGENTS FROM DIFFERENT CLASSES THAT ARE INDICATED FOR DEPRESSION. MDD: 1) NON-PSYCHOTIC, UNIPOLAR DEPRESSION, AND 2) NO ACTIVE SUBSTANCE ABUSE. RENEWAL: TRD, MDD: DEMONSTRATED CLINICAL BENEFIT (IMPROVEMENT IN DEPRESSION) COMPARED TO BASELINE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ETANERCEPT

---

## Products Affected

- ENBREL
- ENBREL MINI
- ENBREL SURECLICK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE OR GENITAL AREA.
Age Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS): 18 YEARS OR OLDER. PSORIATIC ARTHRITIS (PSA): 2 YEARS OR OLDER.
Prescriber Restrictions	INITIAL: RA, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA), AS: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSA: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PJIA, PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD. PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. RENEWAL: RA, PJIA, PSA, AS, PSO: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# EVEROLIMUS-AFINITOR

---

## Products Affected

- *everolimus (antineoplastic) oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# EVEROLIMUS-AFINITOR DISPERZ

---

## Products Affected

- *everolimus (antineoplastic) oral tablet for suspension*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# FECAL MICROBIOTA CAPSULE

---

## Products Affected

- VOWST

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	30 DAYS
Other Criteria	CLOSTRIDIODES DIFFICILE INFECTION (CDI): 1) HAS NOT PREVIOUSLY RECEIVED VOWST: COMPLETION OF ANTIBIOTIC TREATMENT FOR RECURRENT CDI (AT LEAST 3 CDI EPISODES), OR 2) PREVIOUSLY RECEIVED VOWST: (A) TREATMENT FAILURE (DEFINED AS THE PRESENCE OF CDI DIARRHEA WITHIN 8 WEEKS OF FIRST DOSE OF VOWST AND A POSITIVE STOOL TEST FOR C. DIFFICILE), AND (B) HAS NOT RECEIVED MORE THAN ONE TREATMENT COURSE OF VOWST WHICH WAS AT LEAST 12 DAYS AND NOT MORE THAN 8 WEEKS PRIOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# FEDRATINIB

---

## Products Affected

- INREBIC

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
<b>Other Criteria</b>	MYELOFIBROSIS: INITIAL: TRIAL OF OR CONTRAINDICATION TO JAKAFI (RUXOLITINIB). RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# FENFLURAMINE

---

## Products Affected

- FINTEPLA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: DRAVET SYNDROME, LENNOX-GASTAUT SYNDROME (LGS): PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	DRAVET SYNDROME: INITIAL/RENEWAL: 12 MONTHS. LGS: 12 MONTHS.
Other Criteria	INITIAL: LGS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING ANTIEPILEPTIC MEDICATIONS: RUFINAMIDE, FELBAMATE, CLOBAZAM, TOPIRAMATE, LAMOTRIGINE, CLONAZEPAM. RENEWAL: DRAVET SYNDROME: PATIENT HAS SHOWN CONTINUED CLINICAL BENEFIT (E.G. REDUCTION OF SEIZURES, REDUCED LENGTH OF SEIZURES, SEIZURE CONTROL MAINTAINED).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# FENTANYL CITRATE

---

## Products Affected

- *fentanyl citrate buccal lozenge on a handle*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	CANCER RELATED PAIN: 1) CURRENTLY ON A MAINTENANCE DOSE OF CONTROLLED-RELEASE OPIOID PAIN MEDICATION, AND 2) TRIAL OF OR CONTRAINDICATION TO AT LEAST ONE IMMEDIATE-RELEASE ORAL OPIOID PAIN AGENT OR PATIENT HAS DIFFICULTY SWALLOWING TABLETS/CAPSULES. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# FILGRASTIM-AAFI

---

## Products Affected

- NIVESTYM

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# FILGRASTIM-AYOW

---

## Products Affected

- RELEUKO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT NIVESTYM, WHERE INDICATIONS ALIGN.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# FILGRASTIM-SNDZ

---

## Products Affected

- ZARXIO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: NIVESTYM.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# FINERENONE

---

## Products Affected

- KERENDIA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# FINGOLIMOD

---

## Products Affected

- *fingolimod*
- GILENYA ORAL CAPSULE 0.25 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# FINGOLIMOD LAURYL SULFATE

---

## Products Affected

- TASCENSO ODT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	MULTIPLE SCLEROSIS (MS): (1) UNABLE TO SWALLOW FINGOLIMOD CAPSULES, AND (2) TRIAL OF OR CONTRAINDICATION TO FINGOLIMOD CAPSULES.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# FREMANEZUMAB-VFRM

## Products Affected

- AJOVY AUTOINJECTOR
- AJOVY SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	MIGRAINE PREVENTION: INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREVENTIVE MIGRAINE TREATMENTS: DIVALPROEX SODIUM, TOPIRAMATE, PROPRANOLOL, TIMOLOL, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. RENEWAL: 1) REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# FRUQUINTINIB

---

## Products Affected

- FRUZAQLA ORAL CAPSULE 1 MG, 5 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# FUTIBATINIB

---

## Products Affected

- LYTGOBI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INTRAHEPATIC CHOLANGIOCARCINOMA (ICCA): COMPLETE A COMPREHENSIVE OPHTHALMOLOGICAL EXAMINATION, INCLUDING OPTICAL COHERENCE TOMOGRAPHY (OCT), PRIOR TO THE INITIATION OF THERAPY AND AT THE RECOMMENDED SCHEDULED INTERVALS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# GALCANEZUMAB-GNLM

## Products Affected

- EMGALITY PEN 3)
- EMGALITY SYRINGE  
SUBCUTANEOUS SYRINGE 120  
MG/ML, 300 MG/3 ML (100 MG/ML X

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: MIGRAINE PREVENTION: 6 MOS. EPISODIC CLUSTER HEADACHE: 3 MOS. RENEWAL (ALL): 12 MOS.
Other Criteria	INITIAL: MIGRAINE PREVENTION: 1) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREVENTIVE MIGRAINE TREATMENTS: DIVALPROEX SODIUM, TOPIRAMATE, PROPRANOLOL, TIMOLOL. RENEWAL: MIGRAINE PREVENTION: 1) REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. EPISODIC CLUSTER HEADACHE: IMPROVEMENT IN EPISODIC CLUSTER HEADACHE FREQUENCY AS COMPARED TO BASELINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# GANAXOLONE

---

## Products Affected

- ZTALMY

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# GEFITINIB

---

## Products Affected

- *gefitinib*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC NSCLC WITH EGFR MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# GILTERITINIB

---

## Products Affected

- XOSPATA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# GLASDEGIB

---

## Products Affected

- DAURISMO ORAL TABLET 100 MG,  
25 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# GLATIRAMER

---

## Products Affected

- COPAXONE SUBCUTANEOUS SYRINGE 20 MG/ML, 40 MG/ML
- *glatiramer subcutaneous syringe 20 mg/ml, 40 mg/ml*
- *glatopa subcutaneous syringe 20 mg/ml, 40 mg/ml*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# GLP1-DULAGLUTIDE

---

## Products Affected

- TRULICITY

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# GLP1-SEMAGLUTIDE

---

## Products Affected

- OZEMPIC SUBCUTANEOUS PEN INJECTOR 0.25 MG OR 0.5 MG (2 MG/3 ML), 0.25 MG OR 0.5 MG(2 MG/1.5 ML), 1 MG/DOSE (4 MG/3 ML), 2 MG/DOSE (8 MG/3 ML)
- RYBELSUS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# GLP1-TIRZEPATIDE

---

## Products Affected

- MOUNJARO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# GLYCEROL PHENYL BUTYRATE

---

## Products Affected

- RAVICTI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	UREA CYCLE DISORDER (UCD): INITIAL: DIAGNOSIS IS CONFIRMED BY ENZYMATIC, BIOCHEMICAL OR GENETIC TESTING
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	UCD: INITIAL: TRIAL OF OR CONTRAINDICATION TO SODIUM PHENYL BUTYRATE. RENEWAL: PATIENT HAS CLINICAL BENEFIT FROM BASELINE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# GOSERELIN

## Products Affected

- ZOLADEX

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	ENDOMETRIOSIS: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	ENDOMETRIOSIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST.
<b>Coverage Duration</b>	STAGE B2-C PROSTATIC CARCINOMA: 4 MOS. ENDOMETRIOSIS: 6 MOS PER LIFETIME. ALL OTHERS: 12 MONTHS.
<b>Other Criteria</b>	ENDOMETRIOSIS: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, 2) TRIAL OF OR CONTRAINDICATION TO NSAID AND PROGESTIN-CONTAINING PREPARATION, AND 3) HAS NOT RECEIVED A TOTAL OF 6 MONTHS OF TREATMENT PER LIFETIME. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# GUSELKUMAB

## Products Affected

- TREMFYA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE OR GENITAL AREA.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
<b>Other Criteria</b>	INITIAL: PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG). PSO: TRIAL OF OR CONTRAINDICATION ONE CONVENTIONAL THERAPY SUCH AS A PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. RENEWAL: PSO, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# HIGH CONCENTRATION ORAL OPIOID SOLUTIONS

## Products Affected

- *morphine concentrate oral solution*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	OPIOID TOLERANT: 12 MONTHS. HOSPICE, PALLIATIVE CARE OR END OF LIFE CARE: LIFETIME.
Other Criteria	1) OPIOID TOLERANT (I.E. PREVIOUS USE OF 60 MG ORAL MORPHINE PER DAY, 25 MCG TRANSDERMAL FENTANYL PER HOUR, 30 MG ORAL OXYCODONE PER DAY, 8 MG ORAL HYDROMORPHONE PER DAY, 25 MG ORAL OXYMORPHONE PER DAY, 60 MG ORAL HYDROCODONE PER DAY, OR AN EQUIANALGESIC DOSE OF ANOTHER OPIOID) AND HAS TROUBLE SWALLOWING OPIOID TABLETS, CAPSULES, OR LARGE VOLUMES OF LIQUID, OR 2) ENROLLED IN HOSPICE OR PALLIATIVE CARE OR END OF LIFE CARE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# HIGH RISK DRUGS IN THE ELDERLY - CYPROHEPTADINE

---

## Products Affected

- *cyproheptadine oral syrup*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# HIGH RISK DRUGS IN THE ELDERLY - BUTALBITAL-CONTAINING AGENTS

## Products Affected

- *ascomp with codeine*
- *butalbital-acetaminophen-caff oral tablet*
- *butalbital-aspirin-caffeine*
- *codeine-butalbital-asa-caff*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS ARE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# HIGH RISK DRUGS IN THE ELDERLY - CONJUGATED ESTROGEN

---

## Products Affected

- PREMARIN ORAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	VULVAR/VAGINAL ATROPHY, OSTEOPOROSIS AND VASOMOTOR SYMPTOMS OF MENOPAUSE: PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HYPOESTROGENISM TREATMENT, PALLIATIVE TREATMENT, AND HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# HIGH RISK DRUGS IN THE ELDERLY - DIPYRIDAMOLE

---

## Products Affected

- *dipyridamole oral*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# HIGH RISK DRUGS IN THE ELDERLY - DISOPYRAMIDE

---

## Products Affected

- *disopyramide phosphate oral capsule*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# HIGH RISK DRUGS IN THE ELDERLY - ESTRADIOL

## Products Affected

- *dotti*
- *estradiol oral*
- *estradiol transdermal patch semiweekly*
- *estradiol transdermal patch weekly*
- *lyllana*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	VULVAR/VAGINAL ATROPHY, OSTEOPOROSIS AND VASOMOTOR SYMPTOMS OF MENOPAUSE: PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HYPOESTROGENISM TREATMENT, PALLIATIVE TREATMENT, AND HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# HIGH RISK DRUGS IN THE ELDERLY - ESTRADIOL-NORETHINDRONE

---

## Products Affected

- *amabelz*
- *estradiol-norethindrone acet oral tablet 0.5-0.1 mg*
- *mimvey*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	VULVAR/VAGINAL ATROPHY, OSTEOPOROSIS, AND VASOMOTOR SYMPTOMS OF MENOPAUSE: PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HYPOESTROGENISM TREATMENT AND HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# HIGH RISK DRUGS IN THE ELDERLY - ESTROGEN-BAZEDOXIFENE

---

## Products Affected

- DUAVEE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# HIGH RISK DRUGS IN THE ELDERLY - ESTROGEN-MEDROXYPROGESTERONE

---

## Products Affected

- PREMPHASE
- PREMPRO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# HIGH RISK DRUGS IN THE ELDERLY - GLYBURIDE FORMULATIONS

---

## Products Affected

- *glyburide*
- *glyburide micronized*
- *glyburide-metformin*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	TRIAL OF GLIMEPIRIDE, GLIPIZIDE, OR PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS ARE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# HIGH RISK DRUGS IN THE ELDERLY - KETOROLAC

---

## Products Affected

- *ketorolac oral*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	30 DAYS
<b>Other Criteria</b>	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# HIGH RISK DRUGS IN THE ELDERLY - NORETHINDRONE-ESTRADIOL

## Products Affected

- *fyavolv*
- *jinteli*
- *norethindrone ac-eth estradiol oral tablet*  
0.5-2.5 mg-mcg, 1-5 mg-mcg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# HIGH RISK DRUGS IN THE ELDERLY - PHENOBARBITAL

---

## Products Affected

- *phenobarbital*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	FOR TREATMENT OF EPILEPSY/SEIZURES IN PATIENTS WHO ARE NEWLY PRESCRIBED PHENOBARBITAL: PATIENT HAS NOT RESPONDED TO AT LEAST ONE ANTICONVULSANT OR PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# HIGH RISK DRUGS IN THE ELDERLY - PROMETHAZINE

## Products Affected

- *promethazine injection solution* *mg*
- *promethazine oral*
- *promethazine rectal*
- *promethegan rectal suppository 12.5 mg, 25*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	PRURITUS/URTICARIA/SEASONAL/PERENNIAL ALLERGY: 1) TRIAL OF OR CONTRAINDICATION TO A NON-SEDATING ANTIHISTAMINE SUCH AS LEVOCETIRIZINE, OR 2) PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. NAUSEA AND VOMITING: PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH-RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS REQUIRE PHYSICIAN ATTESTATION THAT REQUESTED MEDICATION IS USED TO TREAT A DIAGNOSIS UNRELATED TO THE TERMINAL ILLNESS OR RELATED CONDITION, AND ARE APPROVED WITHOUT TRIAL OF FORMULARY ALTERNATIVES NOR REQUIRING PRESCRIBER ACKNOWLEDGEMENT.
<b>Indications</b>	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# HIGH RISK DRUGS IN THE ELDERLY - SCOPOLAMINE

## Products Affected

- *scopolamine base*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS REQUIRE PHYSICIAN ATTESTATION THAT REQUESTED MEDICATION IS USED TO TREAT A DIAGNOSIS UNRELATED TO THE TERMINAL ILLNESS OR RELATED CONDITION, AND ARE APPROVED WITHOUT REQUIRING PRESCRIBER ACKNOWLEDGEMENT.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# HIGH RISK DRUGS IN THE ELDERLY - SKELETAL MUSCLE RELAXANTS

---

## Products Affected

- *chlorzoxazone oral tablet 500 mg*
- *cyclobenzaprine oral tablet 10 mg, 5 mg*
- *methocarbamol oral tablet 500 mg, 750 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED A HIGH RISK MEDICATION FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# HIGH RISK DRUGS IN THE ELDERLY- DIPHENHYDRAMINE ELIXIR

## Products Affected

- *diphenhydramine hcl oral elixir*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	PRURITUS OR URTICARIA: TRIAL OF OR CONTRAINDICATION TO A NON-SEDATING ANTIHISTAMINE SUCH AS LEVOCETIRIZINE. INSOMNIA: TRIAL OF SILENOR AND BELSOMRA. MOTION SICKNESS AND ANTIPARKINSONISM: PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS AND ANAPHYLACTIC REACTIONS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# HIGH RISK DRUGS IN THE ELDERLY- DIPHENOXYLATE-ATROPINE

---

## Products Affected

- *diphenoxylate-atropine oral tablet*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# HIGH RISK DRUGS IN THE ELDERLY- INDOMETHACIN

---

## Products Affected

- *indomethacin oral capsule 25 mg, 50 mg*
- *indomethacin oral capsule, extended release*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# HIGH RISK DRUGS IN THE ELDERLY- MEGESTROL

---

## Products Affected

- *megestrol oral suspension 400 mg/10 ml (40 mg/ml)*
- *megestrol oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# HIGH RISK DRUGS IN THE ELDERLY- PAROXETINE

---

## Products Affected

- *paroxetine hcl oral suspension*
- *paroxetine hcl oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# IBRUTINIB

---

## Products Affected

- IMBRUVICA ORAL CAPSULE 140 MG, 280 MG, 420 MG
- IMBRUVICA ORAL SUSPENSION
- IMBRUVICA ORAL TABLET 140 MG,

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: BRUKINSA, WHERE INDICATIONS ALIGN.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ICATIBANT

---

## Products Affected

- *icatibant*
- *sajazir*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HEREDITARY ANGIOEDEMA (HAE): DIAGNOSIS CONFIRMED BY COMPLEMENT TESTING.
Age Restrictions	
Prescriber Restrictions	HAE: PRESCRIBED BY OR IN CONSULTATION WITH AN ALLERGIST, IMMUNOLOGIST, OR HEMATOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	HAE: NO CONCURRENT USE WITH OTHER MEDICATIONS FOR TREATMENT OF ACUTE HAE ATTACKS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# IDELALISIB

---

## Products Affected

- ZYDELIG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# IMATINIB

---

## Products Affected

- *imatinib oral tablet 100 mg, 400 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	ADJUVANT GASTROINTESTINAL STROMAL TUMOR TREATMENT: 36 MONTHS. ALL OTHER DIAGNOSES: 12 MONTHS.
Other Criteria	PHILADELPHIA CHROMOSOME POSITIVE CHRONIC MYELOID LEUKEMIA: PATIENT HAS NOT RECEIVED A PREVIOUS TREATMENT WITH ANOTHER TYROSINE KINASE INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# INFIGRATINIB

---

## Products Affected

- TRUSELTIQ

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	CHOLANGIOCARCINOMA: COMPREHENSIVE OPHTHALMOLOGICAL EXAMINATION, INCLUDING OPTICAL COHERENCE TOMOGRAPHY (OCT), WILL BE COMPLETED PRIOR TO INITIATION OF THERAPY AND AT THE RECOMMENDED SCHEDULED INTERVALS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# INFLIXIMAB

---

## Products Affected

- *infliximab*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, OR FACE.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: RA: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, XELJANZ, TREMFYA, RINVOQ, SKYRIZI. PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, SKYRIZI, TREMFYA. AS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, COSENTYX, ENBREL, XELJANZ, RINVOQ. CD: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, RINVOQ, SKYRIZI. UC: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, XELJANZ, RINVOQ. RENEWAL: RA, AS, PSO, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# INTERFERON FOR MS-AVONEX

---

## Products Affected

- AVONEX INTRAMUSCULAR PEN INJECTOR KIT
- AVONEX INTRAMUSCULAR SYRINGE KIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# INTERFERON FOR MS-BETASERON

---

## Products Affected

- BETASERON SUBCUTANEOUS KIT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# INTERFERON FOR MS-PLEGRIDY

---

## Products Affected

- PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML, 63 MCG/0.5 ML- 94 MCG/0.5 ML
- PLEGRIDY SUBCUTANEOUS SYRINGE 125 MCG/0.5 ML, 63 MCG/0.5 ML- 94 MCG/0.5 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# INTERFERON GAMMA-1B

---

## Products Affected

- ACTIMMUNE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: CHRONIC GRANULOMATOUS DISEASE (CGD): PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST, INFECTIOUS DISEASE SPECIALIST, OR IMMUNOLOGIST. SEVERE MALIGNANT OSTEOPECTOSIS (SMO): PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	RENEWAL: CGD, SMO: 1) DEMONSTRATED CLINICAL BENEFIT COMPARED TO BASELINE, AND 2) HAS NOT RECEIVED HEMATOPOIETIC CELL TRANSPLANTATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# IPILIMUMAB

## Products Affected

- YERVOY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: UNRESECT/MET MEL: 4MO, RCC/CRC/HCC: 3MO, ALL OTHERS: 12MO. INITIAL/RENEWAL: CUTAN MEL: 6MO
Other Criteria	RENEWAL: ADJUVANT CUTANEOUS MELANOMA: NO EVIDENCE OF DISEASE RECURRENCE (DEFINED AS THE APPEARANCE OF ONE OR MORE NEW MELANOMA LESIONS: LOCAL, REGIONAL OR DISTANT METASTASIS). THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# IVACAF TOR

---

## Products Affected

- KALYDECO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	CYSTIC FIBROSIS (CF); INITIAL: CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CYSTIC FIBROSIS
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CYSTIC FIBROSIS EXPERT
<b>Coverage Duration</b>	INITIAL: 12 MONTHS. RENEWAL: LIFETIME
<b>Other Criteria</b>	CF: INITIAL: NOT HOMOZYGOUS FOR F508DEL MUTATION IN CFTR GENE. RENEWAL: 1) MAINTAINED, IMPROVED, OR DEMONSTRATED LESS THAN EXPECTED DECLINE IN FEV1 OR BODY MASS INDEX (BMI), OR 2) REDUCTION IN NUMBER OF PULMONARY EXACERBATIONS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# IVOSIDENIB

---

## Products Affected

- TIBSOVO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# IXAZOMIB

---

## Products Affected

- NINLARO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# IXEKIZUMAB

## Products Affected

- TALTZ AUTOINJECTOR
- TALTZ SYRINGE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, OR GENITAL AREA. NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN (CRP) LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. ANKYLOSING SPONDYLITIS (AS), NR-AXSPA: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, COSENTYX, STELARA, ENBREL, SKYRIZI, TREMFYA. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, COSENTYX, STELARA, ENBREL, XELJANZ, TREMFYA, RINVOQ, SKYRIZI. AS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA, COSENTYX, XELJANZ, RINVOQ. NR-AXSPA: 1) TRIAL OF OR CONTRAINDICATION TO BOTH OF THE PREFERRED AGENTS: COSENTYX, RINVOQ, OR 2) TRIAL OF COSENTYX AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. RENEWAL: PSO, PSA, AS, NR-AXSPA: CONTINUES TO BENEFIT FROM THE MEDICATION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# LANADELUMAB-FLYO

## Products Affected

- TAKHZYRO SUBCUTANEOUS SOLUTION (150 MG/ML)
- TAKHZYRO SUBCUTANEOUS SYRINGE 150 MG/ML, 300 MG/2 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HEREDITARY ANGIOEDEMA (HAE): INITIAL: DIAGNOSIS CONFIRMED BY COMPLEMENT TESTING.
Age Restrictions	
Prescriber Restrictions	HAE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ALLERGIIST, IMMUNOLOGIST, OR HEMATOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	HAE: INITIAL: NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS. RENEWAL: 1) IMPROVEMENT COMPARED TO BASELINE IN HAE ATTACKS (I.E., REDUCTIONS IN ATTACK FREQUENCY OR ATTACK SEVERITY), AND 2) NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# LANREOTIDE

## Products Affected

- *lanreotide* ML
- SOMATULINE DEPOT  
SUBCUTANEOUS SYRINGE 120  
MG/0.5 ML, 60 MG/0.2 ML, 90 MG/0.3

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	ACROMEGALY: INITIAL: THERAPY IS PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	ACROMEGALY: INITIAL: 3 MOS, RENEWAL: 12 MOS.GEP-NETS, CARCINOID SYNDROME: 12 MOS.
Other Criteria	ACROMEGALY: INITIAL: TRIAL OF OR CONTRAINDICATION TO ONE GENERIC OCTREOTIDE INJECTION. RENEWAL: 1) REDUCTION, NORMALIZATION, OR MAINTENANCE OF IGF-1 LEVELS BASED ON AGE AND GENDER, AND 2) IMPROVEMENT OR SUSTAINED REMISSION OF CLINICAL SYMPTOMS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# LAPATINIB

---

## Products Affected

- *lapatinib*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# LAROTRECTINIB

---

## Products Affected

- VITRAKVI ORAL CAPSULE 100 MG, 25 MG
- VITRAKVI ORAL SOLUTION

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	APPROVAL FOR VITRAKVI ORAL SOLUTION: TRIAL OF VITRAKVI CAPSULES OR PATIENT IS UNABLE TO TAKE CAPSULE FORMULATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# LEDIPASVIR-SOFOSBUVIR

## Products Affected

- HARVONI ORAL PELLETS IN PACKET 33.75-150 MG, 45-200 MG
- HARVONI ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HCV RNA LEVEL WITHIN PAST 6 MONTHS.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
Other Criteria	1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, AND 2) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING: CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, ROSUVASTATIN, TIPRANA VIR/RITONAVIR, SOFOSBUVIR (AS A SINGLE AGENT), EPCLUSA, ZEPATIER, MAVYRET, OR VOSEVI. REQUESTS FOR HARVONI 45MG-200MG PELLETS: PATIENT IS UNABLE TO SWALLOW TABLETS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# LENALIDOMIDE

---

## Products Affected

- *lenalidomide*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# LENVATINIB

---

## Products Affected

- LENVIMA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# LETERMIVIR

## Products Affected

- PREVYMIS INTRAVENOUS SOLUTION 240 MG/12 ML, 480 MG/24 ML
- PREVYMIS ORAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	HSCT: NOT AT RISK FOR LATE CMV: 4 MOS, AT RISK FOR LATE CMV: 7 MOS. KIDNEY TRANSPLANT: 7 MOS.
Other Criteria	HEMATOPOIETIC STEM CELL TRANSPLANT (HSCT): 1) THERAPY WILL BE INITIATED BETWEEN DAY 0 AND DAY 28 POST TRANSPLANT, AND 2) WILL NOT RECEIVE THE MEDICATION BEYOND 100 DAYS POST TRANSPLANT IF NOT AT RISK FOR LATE CYTOMEGALOVIRUS (CMV) INFECTION AND DISEASE, OR BEYOND 200 DAYS POST TRANSPLANT IF AT RISK FOR LATE CMV INFECTION AND DISEASE. KIDNEY TRANSPLANT: 1) THERAPY WILL BE INITIATED BETWEEN DAY 0 AND DAY 7 POST TRANSPLANT, AND 2) WILL NOT RECEIVE THE MEDICATION BEYOND 200 DAYS POST TRANSPLANT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# LEUPROLIDE

---

## Products Affected

- *leuprolide subcutaneous kit*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	PROSTATE CANCER: 12 MONTHS.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# LEUPROLIDE DEPOT

---

## Products Affected

- *leuprolide (3 month)*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# LEUPROLIDE-ELIGARD

---

## Products Affected

- ELIGARD
- ELIGARD (3 MONTH)
- ELIGARD (4 MONTH)
- ELIGARD (6 MONTH)

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS.
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# LEUPROLIDE-LUPRON DEPOT

## Products Affected

- LUPRON DEPOT
- LUPRON DEPOT (3 MONTH)
- LUPRON DEPOT (4 MONTH)
- LUPRON DEPOT (6 MONTH)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: ENDOMETRIOSIS: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: ENDOMETRIOSIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST.
<b>Coverage Duration</b>	PROSTATE CA: 12 MOS. UTERINE FIBROIDS: 3 MOS. ENDOMETRIOSIS: INITIAL/RENEWAL: 6 MOS.
<b>Other Criteria</b>	INITIAL: ENDOMETRIOSIS: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, 2) TRIAL OF OR CONTRAINDICATION TO NSAID AND PROGESTIN-CONTAINING PREPARATION, AND 3) HAS NOT RECEIVED A TOTAL OF 12 MONTHS OF TREATMENT PER LIFETIME. RENEWAL: ENDOMETRIOSIS: 1) IMPROVEMENT OF PAIN RELATED TO ENDOMETRIOSIS WHILE ON THERAPY, 2) RECEIVING CONCOMITANT ADD-BACK THERAPY (I.E., COMBINATION ESTROGEN-PROGESTIN OR PROGESTIN-ONLY CONTRACEPTIVE PREPARATION), 3) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, AND 4) HAS NOT RECEIVED A TOTAL OF 12 MONTHS OF TREATMENT PER LIFETIME. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# LEVODOPA

---

## Products Affected

- INBRIJA INHALATION CAPSULE,  
W/INHALATION DEVICE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PARKINSONS DISEASE (PD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	PD: INITIAL: 1) NOT CURRENTLY TAKING MORE THAN 1600MG OF LEVODOPA PER DAY, AND 2) PHYSICIAN HAS OPTIMIZED DRUG THERAPY FOR PARKINSONS DISEASE. RENEWAL: IMPROVEMENT WITH MOTOR FLUCTUATIONS DURING OFF EPISODES WITH THE USE OF INBRIJA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# L-GLUTAMINE

## Products Affected

- ENDARI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	SICKLE CELL DISEASE(SCD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST
Coverage Duration	INITIAL: 12 MONTHS. RENEWAL: LIFETIME.
Other Criteria	SCD: INITIAL: AGES 18 YEARS OR OLDER: 1) AT LEAST 2 SICKLE CELL CRISES IN THE PAST YEAR, 2) SICKLE-CELL ASSOCIATED SYMPTOMS WHICH ARE INTERFERING WITH ACTIVITIES OF DAILY LIVING, OR 3) HISTORY OF OR HAS RECURRENT ACUTE CHEST SYNDROME. AGES 5 TO 17 YEARS: APPROVED WITHOUT ADDITIONAL CRITERIA. RENEWAL: MAINTAINED OR EXPERIENCED A REDUCTION IN ACUTE COMPLICATIONS OF SCD.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# LIDOCAINE OINTMENT

---

## Products Affected

- *lidocaine topical ointment*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION. THIS DRUG MAY BE EITHER BUNDLED WITH AND COVERED UNDER END STAGE RENAL DISEASE DIALYSIS RELATED SERVICES OR COVERED UNDER MEDICARE D DEPENDING UPON THE CIRCUMSTANCES.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# LIDOCAINE PATCH

---

## Products Affected

- *lidocaine topical adhesive patch, medicated 5%*
- ZTLIDO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# LIDOCAINE PRILOCAINE

---

## Products Affected

- *lidocaine-prilocaine topical cream*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG MAY BE EITHER BUNDLED WITH AND COVERED UNDER END STAGE RENAL DISEASE DIALYSIS RELATED SERVICES OR COVERED UNDER MEDICARE D DEPENDING UPON THE CIRCUMSTANCES. INFORMATION MAY NEED TO BE SUBMITTED DESCRIBING THE USE AND SETTING OF THE DRUG TO MAKE THE DETERMINATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# LIDOCAINE SOLUTION

---

## Products Affected

- *lidocaine hcl mucous membrane solution 4 % (40 mg/ml)*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# LOMITAPIDE

---

## Products Affected

- JUXTAPID ORAL CAPSULE 10 MG, 20 MG, 30 MG, 40 MG, 5 MG, 60 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST, ENDOCRINOLOGIST OR LIPIDOLOGIST.
<b>Coverage Duration</b>	12 MONTHS

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>1) DIAGNOSIS DETERMINED BY A) DEFINITE SIMON BROOME DIAGNOSTIC CRITERIA, OR B) DUTCH LIPID NETWORK CRITERIA SCORE OF 8 OR GREATER, OR C) CLINICAL DIAGNOSIS BASED ON A HISTORY OF AN UNTREATED LDL-C CONCENTRATION GREATER THAN 500 MG/DL TOGETHER WITH EITHER XANTHOMA BEFORE 10 YEARS OF AGE, OR EVIDENCE OF HEFH IN BOTH PARENTS. 2) LDL-C LEVEL GREATER THAN OR EQUAL TO 70MG/DL WHILE ON MAXIMAL DRUG TREATMENT. 3) TRIAL OF EVOLOCUMAB UNLESS THE PATIENT HAS NON-FUNCTIONING LDL RECEPTORS. 4) MEETS ONE OF THE FOLLOWING: A) TAKING A HIGH-INTENSITY STATIN (I.E., ATORVASTATIN 40-80MG DAILY, ROSUVASTATIN 20-40MG DAILY) FOR A DURATION OF AT LEAST 8 WEEKS, B) TAKING A MAXIMALLY TOLERATED DOSE OF ANY STATIN FOR A DURATION OF AT LEAST 8 WEEKS GIVEN THAT THE PATIENT CANNOT TOLERATE A HIGH-INTENSITY STATIN, C) ABSOLUTE CONTRAINDICATION TO STATIN THERAPY (E.G., ACTIVE DECOMPENSATED LIVER DISEASE, NURSING FEMALE, PREGNANCY OR PLANS TO BECOME PREGNANT, HYPERSENSITIVITY REACTIONS), D) STATIN INTOLERANCE, OR E) TRIAL OF ROSUVASTATIN, ATORVASTATIN, OR STATIN THERAPY AT ANY DOSE AND HAS EXPERIENCED SKELETAL-MUSCLE RELATED SYMPTOMS (E.G., MYOPATHY).</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# LONCASTUXIMAB TESIRINE-LPYL

---

## Products Affected

- ZYNLONTA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# LORLATINIB

---

## Products Affected

- LORBRENA ORAL TABLET 100 MG,  
25 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# LUMACAF TOR-IVACAF TOR

## Products Affected

- ORKAMBI ORAL GRANULES IN PACKET
- ORKAMBI ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: CYSTIC FIBROSIS (CF): CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CF.
Age Restrictions	
Prescriber Restrictions	CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CF EXPERT.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: LIFETIME.
Other Criteria	CF: RENEWAL: MAINTAINED, IMPROVED, OR DEMONSTRATED LESS THAN EXPECTED DECLINE IN FEV1 OR BODY MASS INDEX (BMI), OR REDUCTION IN NUMBER OF PULMONARY EXACERBATIONS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# LUMASIRAN

---

## Products Affected

- OXLUMO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# MACITENTAN

---

## Products Affected

- OPSUMIT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# MARGETUXIMAB-CMKB

---

## Products Affected

- MARGENZA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# MECHLORETHAMINE

---

## Products Affected

- VALCHLOR

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# MEPOLIZUMAB

## Products Affected

- NUCALA SUBCUTANEOUS AUTO-INJECTOR
- NUCALA SUBCUTANEOUS RECON SOLN
- NUCALA SUBCUTANEOUS SYRINGE 100 MG/ML, 40 MG/0.4 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: ASTHMA: BLOOD EOSINOPHIL LEVEL GREATER THAN OR EQUAL TO 150 CELLS/MCL WITHIN THE PAST 12 MONTHS.
Age Restrictions	
Prescriber Restrictions	INITIAL: ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN PULMONARY OR ALLERGY MEDICINE. NASAL POLYPS: PRESCRIBED BY OR IN CONSULTATION WITH AN OTOLARYNGOLOGIST, ALLERGIST OR IMMUNOLOGIST.
Coverage Duration	INITIAL: ASTHMA: 4 MO. NASAL POLYPS: 6 MO. OTHERS: 12 MO. RENEWAL: NASAL POLYPS, ASTHMA: 12 MO.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY TOLERATED DOSE OF AN INHALED CORTICOSTEROID AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 2) ONE OF THE FOLLOWING: (A) AT LEAST ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR (B) POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA, AND 3) NOT CONCURRENTLY RECEIVING XOLAIR, DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA. NASAL POLYPS: PREVIOUS 56 DAY TRIAL OF ONE TOPICAL NASAL CORTICOSTEROID. RENEWAL: ASTHMA: 1) NOT CONCURRENTLY RECEIVING XOLAIR, DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA, 2) CONTINUED USE OF ICS AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY ONE OF THE FOLLOWING: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS, OR (D) INCREASE IN PERCENT PREDICTED FEV1 FROM</p>
	<p>PRETREATMENT BASELINE. NASAL POLYPS: CLINICAL BENEFIT COMPARED TO BASELINE.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	



<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No

# MIDOSTAURIN

---

## Products Affected

- RYDAPT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	ACUTE MYELOID LEUKEMIA: 6 MONTHS. ADVANCED SYSTEMIC MASTOCYTOSIS: 12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# MIFEPRISTONE

## Products Affected

- KORLYM
- *mifepristone oral tablet 300 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	CUSHINGS SYNDROME (CS): INITIAL: DIAGNOSIS CONFIRMED BY ONE OF THE FOLLOWING: 1) 24-HR URINE FREE CORTISOL (2 OR MORE TESTS TO CONFIRM), 2) OVERNIGHT 1MG DEXAMETHASONE TEST, OR 3) LATE NIGHT SALIVARY CORTISOL (2 OR MORE TESTS TO CONFIRM).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	CS: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
<b>Coverage Duration</b>	INITIAL AND RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	CS: INITIAL: HYPERCORTISOLISM IS NOT A RESULT OF CHRONIC GLUCOCORTICOIDS. RENEWAL: 1) CONTINUES TO HAVE IMPROVEMENT OF GLUCOSE TOLERANCE OR STABLE GLUCOSE TOLERANCE (E.G., REDUCED A1C, IMPROVED FASTING GLUCOSE, ETC.), 2) CONTINUES TO HAVE TOLERABILITY TO KORLYM, AND 3) CONTINUES TO NOT BE A CANDIDATE FOR SURGICAL TREATMENT OR HAS FAILED SURGERY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# MIGALASTAT

---

## Products Affected

- GALAFOLD

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	FABRY DISEASE: INITIAL SYMPTOMATIC OR EVIDENCE OF INJURY FROM GL-3 TO THE KIDNEY, HEART, OR CENTRAL NERVOUS SYSTEM RECOGNIZED BY LABORATORY, HISTOLOGICAL, OR IMAGING FINDINGS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	FABRY DISEASE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEPHROLOGIST, CARDIOLOGIST, OR SPECIALIST IN GENETICS OR INHERITED METABOLIC DISORDERS.
<b>Coverage Duration</b>	INITIAL: 6 MOS. RENEWAL: 12 MOS.
<b>Other Criteria</b>	FABRY DISEASE: INITIAL: NOT CONCURRENTLY USING ENZYME REPLACEMENT THERAPY (I.E. FABRAZYME). RENEWAL: 1) IMPROVEMENT OR STABILIZATION, AND 2) NOT CONCURRENTLY USING ENZYME REPLACEMENT THERAPY (I.E. FABRAZYME).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# MIGLUSTAT

---

## Products Affected

- *miglustat*
- *yargesa*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# MILTEFOSINE

---

## Products Affected

- IMPAVIDO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# MOBOCERTINIB

---

## Products Affected

- EXKIVITY

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# MOMELOTINIB

---

## Products Affected

- OJJAARA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# MOSUNETUZUMAB-AXGB

---

## Products Affected

- LUNSUMIO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	RELAPSED OR REFRACTORY FOLLICULAR LYMPHOMA: INITIAL: 6 MONTHS. RENEWAL: 7 MONTHS.
Other Criteria	RELAPSED OR REFRACTORY FOLLICULAR LYMPHOMA: RENEWAL: 1) HAS ACHIEVED A PARTIAL RESPONSE TO TREATMENT, AND 2) HAS NOT PREVIOUSLY RECEIVED MORE THAN 17 CYCLES OF TREATMENT. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# NAFARELIN

## Products Affected

- SYNAREL

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: ENDOMETRIOSIS: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS. CENTRAL PRECOCIOUS PUBERTY (CPP): FEMALES: ELEVATED LEVELS OF FOLLICLE-STIMULATING HORMONE (FSH) GREATER THAN 4.0 MIU/ML AND LUTEINIZING HORMONE (LH) LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS. MALES: ELEVATED LEVELS OF FSH GREATER THAN 5.0 MIU/ML AND LH LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: ENDOMETRIOSIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST. CPP: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
<b>Coverage Duration</b>	ENDOMETRIOSIS: 6 MONTHS. CPP: INITIAL/RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: ENDOMETRIOSIS: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, 2) TRIAL OF OR CONTRAINDICATION TO NSAID AND PROGESTIN-CONTAINING PREPARATION, AND 3) HAS NOT RECEIVED A TOTAL OF 6 MONTHS OF TREATMENT PER LIFETIME.</p> <p>CPP: FEMALES: 1) YOUNGER THAN 8 YEARS OF AGE AT ONSET OF CPP, AND 2) AT TANNER STAGE 2 OR ABOVE FOR BREAST DEVELOPMENT AND PUBIC HAIR GROWTH.</p> <p>MALES: 1) YOUNGER THAN 9 YEARS OF AGE AT ONSET OF CPP, AND 2) AT TANNER STAGE 2 OR ABOVE FOR GENITAL DEVELOPMENT AND PUBIC HAIR GROWTH.</p> <p>RENEWAL: CPP: 1) TANNER STAGING AT INITIAL DIAGNOSIS HAS STABILIZED OR REGRESSED DURING THREE SEPARATE MEDICAL VISITS IN THE PREVIOUS YEAR, AND 2) HAS NOT REACHED ACTUAL AGE WHICH CORRESPONDS TO CURRENT PUBERTAL AGE.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# NARCOLEPSY AGENTS

---

## Products Affected

- *armodafinil*
- *modafinil oral tablet 100 mg, 200 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# NATALIZUMAB

## Products Affected

- TYSABRI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: CROHNS DISEASE (CD): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	MULTIPLE SCLEROSIS (MS): 12 MOS. CD: INITIAL: 6 MOS, RENEWAL: 12 MOS.
Other Criteria	INITIAL: MS: TRIAL OF TWO AGENTS INDICATED FOR THE TREATMENT OF MS. CD: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, SKYRIZI, RINVOQ. RENEWAL: CD: 1) RECEIVED AT LEAST 12 MONTHS OF THERAPY WITH TYSABRI AND HAS NOT REQUIRED MORE THAN 3 MONTHS OF CORTICOSTEROID USE WITHIN THE PAST 12 MONTHS TO CONTROL CROHNS DISEASE WHILE ON TYSABRI, OR 2) HAS ONLY RECEIVED 6 MONTHS OF THERAPY WITH TYSABRI AND HAS TAPERED OFF CORTICOSTEROIDS DURING THE FIRST 24 WEEKS OF TYSABRI THERAPY. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# NAXITAMAB-GQGK

---

## Products Affected

- DANYELZA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# NERATINIB

---

## Products Affected

- NERLYNX

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	EARLY-STAGE (STAGE I-III) BREAST CANCER: MEDICATION IS BEING REQUESTED WITHIN 2 YEARS OF COMPLETING THE LAST TRASTUZUMAB DOSE. ALL OTHER FDA APPROVED INDICATIONS ARE COVERED WITHOUT ADDITIONAL CRITERIA, EXCEPT THOSE CRITERIA IN THE FDA APPROVED LABEL.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# NILOTINIB

---

## Products Affected

- TASIGNA ORAL CAPSULE 150 MG, 200 MG, 50 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	PREVIOUSLY TREATED CML: MUTATIONAL ANALYSIS PRIOR TO INITIATION AND TASIGNA IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# NINTEDANIB

## Products Affected

- OFEV

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	<p>INITIAL: IDIOPATHIC PULMONARY FIBROSIS (IPF): 1) A USUAL INTERSTITIAL PNEUMONIA (UIP) PATTERN AS EVIDENCED BY HIGH-RESOLUTION COMPUTED TOMOGRAPHY (HRCT) ALONE OR VIA A COMBINATION OF SURGICAL LUNG BIOPSY AND HRCT, AND 2) BASELINE FORCED VITAL CAPACITY (FVC) AT LEAST 50% OF PREDICTED VALUE. SYSTEMIC SCLEROSIS-ASSOCIATED INTERSTITIAL LUNG DISEASE (SSC-ILD): 1) AT LEAST 10% FIBROSIS ON A CHEST HRCT, AND 2) BASELINE FVC AT LEAST 40% OF PREDICTED VALUE. CHRONIC FIBROSING INTERSTITIAL LUNG DISEASE WITH A PROGRESSIVE PHENOTYPE (PF-ILD): 1) AT LEAST 10% FIBROSIS ON A CHEST HRCT, AND 2) BASELINE FVC AT LEAST 45% OF PREDICTED VALUE.</p>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	<p>INITIAL: IPF: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST. SSC-ILD, PF-ILD: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR RHEUMATOLOGIST.</p>
<b>Coverage Duration</b>	<p>INITIAL: SSC-ILD: 6 MOS. IPF, PF-ILD: 12 MOS. RENEWAL (ALL INDICATIONS): 12 MOS.</p>

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: IPF: 1) DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., CONNECTIVE TISSUE DISEASE, DRUG TOXICITY, ASBESTOS OR BERYLLIUM EXPOSURE, HYPERSENSITIVITY PNEUMONITIS), AND 2) TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: ESBRIET (PIRFENIDONE). SSC-ILD: 1) DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., HEART FAILURE/FLUID OVERLOAD, DRUG-INDUCED LUNG TOXICITY, RECURRENT ASPIRATION), AND 2) TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: ACTEMRA SUBQ. PF-ILD: LUNG FUNCTION AND RESPIRATORY SYMPTOMS OR CHEST IMAGING HAVE WORSENERD/PROGRESSED DESPITE TREATMENT WITH MEDICATIONS USED IN CLINICAL PRACTICE FOR ILD (NOT ATTRIBUTABLE TO COMORBIDITIES SUCH AS INFECTION, HEART FAILURE). RENEWAL: IPF, SSC-ILD, PF-ILD: CLINICAL MEANINGFUL IMPROVEMENT OR MAINTENANCE IN ANNUAL RATE OF DECLINE.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# NIRAPARIB

---

## Products Affected

- ZEJULA ORAL CAPSULE
- ZEJULA ORAL TABLET

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	RECURRENT EPITHELIAL OVARIAN, FALLOPIAN TUBE, OR PRIMARY PERITONEAL CANCER: 1) ZEJULA WILL BE USED AS MONOTHERAPY, AND 2) ZEJULA IS STARTED NO LATER THAN 8 WEEKS AFTER THE MOST RECENT PLATINUM-CONTAINING REGIMEN.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# NIRAPARIB/ABIRATERONE

---

## Products Affected

- AKEEGA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (MCRPC): 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# NIROGACESTAT

---

## Products Affected

- OGSIVEO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# NITISINONE

## Products Affected

- *nitisinone*
- ORFADIN ORAL SUSPENSION

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	HEREDITARY TYROSINEMIA TYPE 1 (HT-1): INITIAL: DIAGNOSIS CONFIRMED BY ELEVATED URINARY OR PLASMA SUCCINYLACETONE LEVELS OR A MUTATION IN THE FUMARYLACETOACETATE HYDROLASE GENE. RENEWAL: URINARY OR PLASMA SUCCINYLACETONE LEVELS HAVE DECREASED FROM BASELINE WHILE ON TREATMENT WITH NITISINONE.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	HT-1: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PRESCRIBER SPECIALIZING IN INHERITED METABOLIC DISEASES.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	HT-1: INITIAL: ORFADIN SUSPENSION: TRIAL OF OR CONTRAINDICATION TO PREFERRED NITISINONE TABLETS OR CAPSULES.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# NIVOLUMAB

---

## Products Affected

- OPDIVO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	UNRESECTABLE OR METASTATIC MELANOMA: NO CONCURRENT USE WITH TARGETED THERAPY (I.E., BRAF INHIBITORS, MEK INHIBITORS, AND NTRK INHIBITORS).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# NIVOLUMAB-RELATLIMAB-RMBW

---

## Products Affected

- OPDUALAG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# OBETICHOLIC ACID

## Products Affected

- OCALIVA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	PRIMARY BILIARY CHOLANGITIS (PBC): INITIAL/RENEWAL: COMPLETE BILIARY OBSTRUCTION.
<b>Required Medical Information</b>	PBC: INITIAL: DIAGNOSIS CONFIRMED BY TWO OF THE FOLLOWING: 1) ALKALINE PHOSPHATASE LEVEL OF AT LEAST 1.5 TIMES THE UPPER LIMIT OF NORMAL, 2) PRESENCE OF ANTIMITOCHONDRIAL ANTIBODIES AT A TITER OF 1:40 OR HIGHER, OR 3) HISTOLOGIC EVIDENCE OF NON-SUPPURATIVE DESTRUCTIVE CHOLANGITIS AND DESTRUCTION OF INTERLOBULAR BILE DUCTS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PBC: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST OR HEPATOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	PBC: INITIAL: USED IN COMBINATION WITH URSODEOXYCHOLIC ACID IN A PATIENT WITH AN INADEQUATE RESPONSE TO URSODEOXYCHOLIC ACID, OR AS MONOTHERAPY IN A PATIENT WHO IS UNABLE TO TOLERATE URSODEOXYCHOLIC ACID. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# OCRELIZUMAB

---

## Products Affected

- OCREVUS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	RELAPSING FORM OF MULTIPLE SCLEROSIS (MS): TRIAL OF TWO AGENTS INDICATED FOR THE TREATMENT OF RELAPSING FORMS OF MS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# OFATUMUMAB-SQ

---

## Products Affected

- KESIMPTA PEN

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# OLANZAPINE/SAMIDORPHAN

## Products Affected

- LYBALVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	SCHIZOPHRENIA/BIPOLAR I: PRESCRIBED BY OR IN CONSULTATION WITH A PSYCHIATRIST
Coverage Duration	12 MONTHS
Other Criteria	SCHIZOPHRENIA: (1) PATIENT IS AT HIGH RISK OF WEIGHT GAIN AND (2) TRIAL OF OR CONTRAINDICATION TO LATUDA OR ONE OF THE FOLLOWING ORAL ANTIPSYCHOTICS: RISPERIDONE, CLOZAPINE TABLET, OLANZAPINE, IMMEDIATE RELEASE QUETIAPINE FUMARATE, ZIPRASIDONE, ARIPIPRAZOLE. BIPOLAR I: (1) PATIENT IS AT HIGH RISK OF WEIGHT GAIN AND (2) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING ORAL ANTIPSYCHOTICS: RISPERIDONE, OLANZAPINE, IMMEDIATE RELEASE QUETIAPINE FUMARATE, ZIPRASIDONE, ARIPIPRAZOLE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# OLAPARIB

## Products Affected

- LYNPARZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	RECURRENT EPITHELIAL OVARIAN, FALLOPIAN TUBE OR PRIMARY PERITONEAL CANCER: MEDICATION WILL BE USED AS MONOTHERAPY. METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: 1) PREVIOUSLY RECEIVED A BILATERAL ORCHIECTOMY, OR 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. ALL OTHER FDA APPROVED INDICATIONS ARE COVERED WITHOUT ADDITIONAL CRITERIA, EXCEPT THOSE CRITERIA IN THE FDA APPROVED LABEL.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# OLUTASIDENIB

---

## Products Affected

- REZLIDHIA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# OMACETAXINE

---

## Products Affected

- SYNRIPO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# OMALIZUMAB

## Products Affected

- XOLAIR SUBCUTANEOUS AUTO-INJECTOR 300 MG/2 ML
- XOLAIR SUBCUTANEOUS RECON SOLN
- XOLAIR SUBCUTANEOUS SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: ASTHMA: POSITIVE SKIN PRICK OR BLOOD TEST (E.G., ELISA, FEIA) TO A PERENNIAL AEROALLERGEN AND A BASELINE IGE SERUM LEVEL GREATER THAN OR EQUAL TO 30 IU/ML.
Age Restrictions	
Prescriber Restrictions	INITIAL AND RENEWAL: CHRONIC SPONTANEOUS URTICARIA (CSU): PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE, DERMATOLOGY OR IMMUNOLOGY. INITIAL: NASAL POLYPS: PRESCRIBED BY OR IN CONSULTATION WITH AN OTOLARYNGOLOGIST, ALLERGIST OR IMMUNOLOGIST. ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE.
Coverage Duration	INITIAL: ASTHMA: 4 MO. CSU, NASAL POLYPS: 6 MO. RENEWAL: ASTHMA, NASAL POLYPS: 12 MO. CSU: 6 MO.



PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: CSU: TRIAL OF OR CONTRAINDICATION TO A MAXIMALLY TOLERATED DOSE OF AN H1 ANTI-HISTAMINE AND STILL EXPERIENCES HIVES ON MOST DAYS OF THE WEEK. NASAL POLYPS: 1) PREVIOUS 56 DAY TRIAL OF ONE TOPICAL NASAL CORTICOSTEROID, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE PREFERRED AGENT: NUCALA, DUPIXENT. ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY TOLERATED DOSE OF AN INHALED CORTICOSTEROID AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, 2) ONE OF THE FOLLOWING: (A) PATIENT EXPERIENCED AT LEAST ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR (B) PATIENT HAS POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA, 3) NOT CONCURRENTLY RECEIVING DUPIXENT OR ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA. RENEWAL: CSU: DIAGNOSIS OF CSU. NASAL POLYPS: CLINICAL BENEFIT COMPARED TO BASELINE. ASTHMA: 1) NOT CONCURRENTLY RECEIVING DUPIXENT OR ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA, 2) CONTINUED USE OF ICS AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY ONE OF THE FOLLOWING: (A) REDUCTION IN ASTHMA</p>
	<p>EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS, OR (D) INCREASE IN PERCENT PREDICTED FEV1 FROM PRETREATMENT BASELINE.</p>
<b>Indications</b>	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# OSIMERTINIB

---

## Products Affected

- TAGRISSO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	EGFR EXON 19 DELETIONS OR EXON 21 L858R MUTATIONS NON-SMALL CELL LUNG CANCER (NSCLC) AND METASTATIC NSCLC WITH EGFR T790M MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# OXANDROLONE

---

## Products Affected

- *oxandrolone*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	PROTEIN CATABOLISM, BONE PAIN: 1) MONITORED FOR PELIOSIS HEPATIS, LIVER CELL TUMORS, AND BLOOD LIPID CHANGES, 2) DOES NOT HAVE KNOWN OR SUSPECTED: CARCINOMA OF THE PROSTATE OR BREAST IN MALE PATIENTS, CARCINOMA OF THE BREAST IN FEMALES WITH HYPERCALCEMIA, NEPHROSIS (THE NEPHROTIC PHASE OF NEPHRITIS), OR HYPERCALCEMIA, AND 3) DOES NOT HAVE SEVERE HEPATIC DYSFUNCTION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PACRITINIB

---

## Products Affected

- VONJO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
<b>Other Criteria</b>	MYELOFIBROSIS: RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PALBOCICLIB

---

## Products Affected

- IBRANCE

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	ADVANCED OR METASTATIC BREAST CANCER: TRIAL OF OR CONTRAINDICATION TO ONE OF THE PREFERRED AGENTS, WHERE INDICATIONS ALIGN: KISQALI, VERZENIO.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PANOBINOSTAT

---

## Products Affected

- FARYDAK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	MULTIPLE MYELOMA: RENEWAL: TOLERATED THE FIRST 8 CYCLES OF THERAPY WITHOUT UNRESOLVED SEVERE OR MEDICALLY SIGNIFICANT TOXICITY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PARATHYROID HORMONE

---

## Products Affected

- NATPARA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	HYPOCALCEMIA SECONDARY TO HYPOPARATHYROIDISM: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	HYPOCALCEMIA SECONDARY TO HYPOPARATHYROIDISM: 1) TRIAL OF OR CONTRAINDICATION TO CALCITRIOL, 2) HYPOPARATHYROIDISM IS NOT DUE TO A CALCIUM SENSING RECEPTOR (CSR) MUTATION, AND 3) HYPOPARATHYROIDISM IS NOT CONSIDERED ACUTE POST-SURGICAL HYPOPARATHYROIDISM.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# PASIREOTIDE DIASPARTATE

---

## Products Affected

- SIGNIFOR

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	CUSHINGS DISEASE (CD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	CD: RENEWAL: 1) CONTINUED IMPROVEMENT OF CUSHINGS DISEASE, AND 2) MAINTAINED TOLERABILITY TO SIGNIFOR.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PAZOPANIB

---

## Products Affected

- *pazopanib*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ADVANCED SOFT TISSUE SARCOMA (STS): NOT USED FOR ADIPOCYTIC STS OR GASTROINTESTINAL STROMAL TUMORS (GIST)
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PEGFILGRASTIM - APGF

---

## Products Affected

- NYVEPRIA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	NON MYELOID MALIGNANCY: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PEGFILGRASTIM - CBQV

---

## Products Affected

- UDENYCA
- UDENYCA AUTOINJECTOR
- UDENYCA ONBODY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NON MYELOID MALIGNANCY, ACUTE RADIATION EXPOSURE: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	NON MYELOID MALIGNANCY: TRIAL OF OR CONTRAINDICATION TO NYVEPRIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PEGFILGRASTIM - JMDB

---

## Products Affected

- FULPHILA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	NON MYELOID MALIGNANCY: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	NON MYELOID MALIGNANCY: TRIAL OF OR CONTRAINDICATION TO NYVEPRIA
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PEGFILGRASTIM-NEULASTA ONPRO

---

## Products Affected

- NEULASTA ONPRO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	NON MYELOID MALIGNANCY, ACUTE RADIATION EXPOSURE: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PEGINTERFERON ALFA-2A

---

## Products Affected

- PEGASYS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	CHRONIC HEPATITIS C VIRUS INFECTION. CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	HEPATITIS B: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST, INFECTIOUS DISEASE SPECIALIST, OR PHYSICIAN SPECIALIZING IN THE TREATMENT OF HEPATITIS (E.G., HEPATOLOGIST).
<b>Coverage Duration</b>	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
<b>Other Criteria</b>	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PEGVALIASE-PQPZ

---

## Products Affected

- PALYNZIQ

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	PHENYLKETONURIA (PKU): INITIAL: NOT ON CONCURRENT TREATMENT WITH KUVAN. RENEWAL: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NOT ON CONCURRENT TREATMENT WITH KUVAN.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# PEGVISOMANT

---

## Products Affected

- SOMAVERT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PEMBROLIZUMAB

---

## Products Affected

- KEYTRUDA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	UNRESECTABLE OR METASTATIC MELANOMA: NO CONCURRENT USE WITH TARGETED THERAPY (I.E., BRAF INHIBITORS, MEK INHIBITORS, AND NTRK INHIBITORS).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PEMIGATINIB

---

## Products Affected

- PEMAZYRE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	CHOLANGIOCARCINOMA, MYELOID/LYMPHOID NEOPLASMS: COMPREHENSIVE OPHTHALMOLOGICAL EXAMINATION, INCLUDING OPTICAL COHERENCE TOMOGRAPHY (OCT), WILL BE COMPLETED PRIOR TO INITIATION OF THERAPY AND AT THE RECOMMENDED SCHEDULED INTERVALS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PENICILLAMINE TABLET

---

## Products Affected

- *penicillamine oral tablet*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: CYSTINURIA: HAS NEPHROLITHIASIS AND ONE OF THE FOLLOWING: 1) STONE ANALYSIS SHOWING PRESENCE OF CYSTINE, 2) PRESENCE OF PATHOGNOMONIC HEXAGONAL CYSTINE CRYSTALS ON URINALYSIS, OR 3) FAMILY HISTORY OF CYSTINURIA AND POSITIVE CYANIDE-NITROPRUSSIDE SCREENING.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: WILSONS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A HEPATOLOGIST OR GASTROENTEROLOGIST. CYSTINURIA: PRESCRIBED BY OR IN CONSULTATION WITH A NEPHROLOGIST. RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
<b>Coverage Duration</b>	INITIAL: 12 MONTHS, RENEWAL: LIFETIME.

PA Criteria	Criteria Details
<b>Other Criteria</b>	INITIAL: WILSONS DISEASE: 1) LEIPZIG SCORE OF 4 OR GREATER. RA: 1) NO HISTORY OR OTHER EVIDENCE OF RENAL INSUFFICIENCY, AND 2) TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. RENEWAL: RA: 1) NO HISTORY OR OTHER EVIDENCE OF RENAL INSUFFICIENCY, AND 2) EXPERIENCED OR MAINTAINED IMPROVEMENT IN TENDER JOINT COUNT OR SWOLLEN JOINT COUNT COMPARED TO BASELINE. WILSONS DISEASE, CYSTINURIA: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PEXIDARTINIB

---

## Products Affected

- TURALIO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PIMAVANSERIN

---

## Products Affected

- NUPLAZID

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	PSYCHOSIS IN PARKINSONS DISEASE (PD): INITIAL: 18 YEARS OR OLDER
<b>Prescriber Restrictions</b>	PSYCHOSIS IN PD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, GERIATRICIAN, OR A BEHAVIORAL HEALTH SPECIALIST (E.G., PSYCHIATRIST).
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	PSYCHOSIS IN PD: RENEWAL: IMPROVEMENT IN PSYCHOSIS SYMPTOMS FROM BASELINE AND DEMONSTRATES A CONTINUED NEED FOR TREATMENT.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PIRFENIDONE

## Products Affected

- *pirfenidone oral capsule*
- *pirfenidone oral tablet 267 mg, 534 mg, 801 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	IDIOPATHIC PULMONARY FIBROSIS (IPF): INITIAL: 1) A USUAL INTERSTITIAL PNEUMONIA (UIP) PATTERN AS EVIDENCED BY HIGH-RESOLUTION COMPUTED TOMOGRAPHY (HRCT) ALONE OR VIA A COMBINATION OF SURGICAL LUNG BIOPSY AND HRCT, AND 2) PREDICTED FORCED VITAL CAPACITY (FVC) OF AT LEAST 50% AT BASELINE.
<b>Age Restrictions</b>	IPF: INITIAL: 18 YEARS OR OLDER.
<b>Prescriber Restrictions</b>	IPF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	IPF: INITIAL: 1) DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., CONNECTIVE TISSUE DISEASE, DRUG TOXICITY, ASBESTOS OR BERYLLIUM EXPOSURE, HYPERSENSITIVITY PNEUMONITIS, SYSTEMIC SCLEROSIS, RHEUMATOID ARTHRITIS, RADIATION, SARCOIDOSIS, BRONCHIOLITIS OBLITERANS ORGANIZING PNEUMONIA, HUMAN IMMUNODEFICIENCY VIRUS (HIV) INFECTION, VIRAL HEPATITIS, OR CANCER). RENEWAL: CLINICAL MEANINGFUL IMPROVEMENT OR MAINTENANCE IN ANNUAL RATE OF DECLINE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	



<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No

# PIRTOBRUTINIB

---

## Products Affected

- JAYPIRCA ORAL TABLET 100 MG, 50 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# POMALIDOMIDE

---

## Products Affected

- POMALYST

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PONATINIB

---

## Products Affected

- ICLUSIG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	CML: MUTATIONAL ANALYSIS PRIOR TO INITIATION AND ICLUSIG IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# POSACONAZOLE

## Products Affected

- *posaconazole oral*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	OROPHARYNGEAL CANDIDIASIS (OPC): 3 MONTHS. PROPHYLAXIS: 6 MONTHS. TREATMENT: 12 WEEKS.
<b>Other Criteria</b>	POSACONAZOLE SUSPENSION ONLY: 1) OPC: TRIAL OF OR CONTRAINDICATION TO FLUCONAZOLE OR ITRACONAZOLE. 2) PROPHYLAXIS OF INVASIVE ASPERGILLUS AND CANDIDA INFECTION: INABILITY TO SWALLOW TABLETS. POSACONAZOLE TABLETS ONLY: 1) TREATMENT OF INVASIVE ASPERGILLOSIS, 2) PROPHYLAXIS OF INVASIVE ASPERGILLUS AND CANDIDA INFECTION: AT HIGH RISK OF DEVELOPING THESE INFECTIONS DUE TO BEING SEVERELY IMMUNOCOMPROMISED. CONTINUATION OF THERAPY AFTER HOSPITAL DISCHARGE REQUIRES NO EXTRA CRITERIA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# POSACONAZOLE-POWDERMIX

---

## Products Affected

- NOXAFIL ORAL SUSP,DELAYED  
RELEASE FOR RECON

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	PROPHYLAXIS OF INVASIVE ASPERGILLUS AND CANDIDA INFECTION: INABILITY TO SWALLOW TABLETS. CONTINUATION OF THERAPY AFTER HOSPITAL DISCHARGE REQUIRES NO EXTRA CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PRALSETINIB

---

## Products Affected

- GAVRETO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PRAMLINTIDE

---

## Products Affected

- SYMLINPEN 120
- SYMLINPEN 60

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	TYPE I OR TYPE II DIABETES: REQUIRING INSULIN OR CONTINUOUS INSULIN INFUSION (INSULIN PUMP) FOR GLYCEMIC CONTROL
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# PYRIMETHAMINE

## Products Affected

- *pyrimethamine*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	TOXOPLASMOSIS: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN INFECTIOUS DISEASE SPECIALIST.
<b>Coverage Duration</b>	TOXOPLASMOSIS: INITIAL: 8 WEEKS, RENEWAL: 6 MOS.
<b>Other Criteria</b>	TOXOPLASMOSIS: RENEWAL: ONE OF THE FOLLOWING: (1) PERSISTENT CLINICAL DISEASE (HEADACHE, NEUROLOGICAL SYMPTOMS, OR FEVER) AND PERSISTENT RADIOGRAPHIC DISEASE (ONE OR MORE MASS LESIONS ON BRAIN IMAGING), OR (2) CD4 COUNT LESS THAN 200 CELLS/MM3 AND CURRENTLY TAKING AN ANTI-RETROVIRAL THERAPY IF HIV POSITIVE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# QUININE

---

## Products Affected

- *quinine sulfate*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# QUIZARTINIB

---

## Products Affected

- VANFLYTA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RAMUCIRUMAB

---

## Products Affected

- CYRAMZA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# REGORAFENIB

---

## Products Affected

- STIVARGA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RELUGOLIX

---

## Products Affected

- ORGOVYX

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# REPOTRECTINIB

---

## Products Affected

- AUGTYRO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RESLIZUMAB

---

## Products Affected

- CINQAIR

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	ASTHMA: INITIAL: BLOOD EOSINOPHIL LEVEL GREATER THAN OR EQUAL TO 150 CELLS/MCL WITHIN THE PAST 12 MONTHS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	ASTHMA: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE.
<b>Coverage Duration</b>	ASTHMA: INITIAL: 4 MONTHS. RENEWAL: 12 MONTHS



PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>ASTHMA: INITIAL: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND ONE OTHER MAINTENANCE MEDICATION, AND 2) ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS, OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA, 3) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: FASENRA, NUCALA, DUPIXENT, AND 4) NO CONCURRENT USE WITH XOLAIR, DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA. RENEWAL: 1) NO CONCURRENT USE WITH XOLAIR, DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA, 2) CONTINUED USE OF ICS AND ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS, OR (D) INCREASE IN PERCENT PREDICTED FEV1 FROM PRETREATMENT BASELINE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RETIFANLIMAB-DLWR

---

## Products Affected

- ZYNYZ

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RIBOCICLIB

---

## Products Affected

- KISQALI ORAL TABLET 200 MG/DAY (200 MG X 1), 400 MG/DAY (200 MG X 2), 600 MG/DAY (200 MG X 3)

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RIBOCICLIB-LETROZOLE

---

## Products Affected

- KISQALI FEMARA CO-PACK ORAL  
TABLET 200 MG/DAY(200 MG X 1)-2.5  
MG, 400 MG/DAY(200 MG X 2)-2.5  
MG, 600 MG/DAY(200 MG X 3)-2.5 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RIFAXIMIN

---

## Products Affected

- XIFAXAN ORAL TABLET 200 MG, 550 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	TRAVELERS DIARRHEA/HE: 12 MOS. IBS-D: 8 WKS.
<b>Other Criteria</b>	RIFAXIMIN 550 MG TABLETS: HE: TRIAL OF OR CONTRAINDICATION TO LACTULOSE OR CONCURRENT LACTULOSE THERAPY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RIMEGEPANT

## Products Affected

- NURTEC ODT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: ACUTE MIGRAINE TREATMENT: TRIAL OF OR CONTRAINDICATION TO ONE TRIPTAN (E.G., SUMATRIPTAN, RIZATRIPTAN). EPISODIC MIGRAINE PREVENTION: 1) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREVENTIVE MIGRAINE TREATMENTS: DIVALPROEX SODIUM, TOPIRAMATE, PROPRANOLOL, TIMOLOL. RENEWAL: ACUTE MIGRAINE TREATMENT: 1) IMPROVEMENT FROM BASELINE IN A VALIDATED ACUTE TREATMENT PATIENT-REPORTED OUTCOME QUESTIONNAIRE, OR 2) THERAPY WORKS CONSISTENTLY IN MAJORITY OF MIGRAINE ATTACKS. EPISODIC MIGRAINE PREVENTION: 1) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION, AND 2) REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY.
<b>Indications</b>	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RIOCIQUAT

---

## Products Affected

- ADEMPAS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PULMONARY ARTERIAL HYPERTENSION (PAH): DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS. PERSISTENT/RECURRENT CHRONIC THROMBOEMBOLIC PULMONARY HYPERTENSION (CTEPH) (WHO GROUP 4): WHO FUNCTIONAL CLASS II-IV SYMPTOMS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PAH, CTEPH: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.



PA Criteria	Criteria Details
<b>Other Criteria</b>	INITIAL: PAH: NOT CONCURRENTLY TAKING NITRATES, NITRIC OXIDE DONORS, PHOSPHODIESTERASE INHIBITORS, OR NON-SPECIFIC PDE INHIBITORS. CTEPH: 1) NOT CONCURRENTLY TAKING NITRATES, NITRIC OXIDE DONORS, PHOSPHODIESTERASE INHIBITORS, OR NON-SPECIFIC PDE INHIBITORS, AND 2) NOT A CANDIDATE FOR SURGERY OR HAS INOPERABLE CTEPH OR HAS PERSISTENT OR RECURRENT DISEASE AFTER SURGICAL TREATMENT. RENEWAL: PAH, CTEPH: NOT CONCURRENTLY TAKING NITRATES, NITRIC OXIDE DONORS, PHOSPHODIESTERASE INHIBITORS, OR NON-SPECIFIC PDE INHIBITORS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RIPRETINIB

---

## Products Affected

- QINLOCK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RISANKIZUMAB-RZAA

## Products Affected

- SKYRIZI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PLAQUE PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE OR GENITAL AREA.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. CROHNS DISEASE (CD): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
<b>Other Criteria</b>	INITIAL: PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG). CD: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPYRINE, METHOTREXATE, OR MESALAMINE. RENEWAL: PSO, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RISDIPLAM

## Products Affected

- EVRYSDI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	SPINAL MUSCULAR ATROPHY (SMA): INITIAL: GENE MUTATION ANALYSIS INDICATING MUTATIONS OR DELETIONS OF BOTH ALLELES OF THE SURVIVAL MOTOR NEURON 1 (SMN1) GENE. FOR PRESYMPTOMATIC PATIENTS: UP TO THREE COPIES OF SURVIVAL MOTOR NEURON 2 (SMN2) BASED ON NEWBORN SCREENING.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	SMA: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROMUSCULAR SPECIALIST OR SMA SPECIALIST AT A SMA SPECIALTY CENTER.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	SMA: INITIAL: FOR SYMPTOMATIC PATIENTS: (1) BASELINE MOTOR FUNCTION ASSESSMENT BY A NEUROMUSCULAR SPECIALIST OR SMA SPECIALIST, AND (2) IF PATIENT RECEIVED GENE THERAPY, THE PATIENT HAD LESS THAN EXPECTED CLINICAL BENEFIT. RENEWAL: IMPROVED, MAINTAINED, OR DEMONSTRATED LESS THAN EXPECTED DECLINE IN: (1) MOTOR FUNCTION ASSESSMENTS COMPARED TO BASELINE, OR (2) OTHER MUSCLE FUNCTION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RITUXIMAB AND HYALURONIDASE HUMAN-SQ

---

## Products Affected

- RITUXAN HYCELA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	FOLLICULAR LYMPHOMA (FL), DIFFUSE LARGE B-CELL LYMPHOMA (DLBCL), CHRONIC LYMPHOCYTIC LEUKEMIA (CLL): HAS RECEIVED OR WILL RECEIVE AT LEAST ONE FULL DOSE OF A RITUXIMAB PRODUCT BY INTRAVENOUS INFUSION PRIOR TO INITIATION OF RITUXIMAB AND HYALURONIDASE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# RITUXIMAB-ABBS

## Products Affected

- TRUXIMA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. NHL, CLL: PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST.
<b>Coverage Duration</b>	RA: INITIAL: 6 MO, RENEWAL: 12 MO. NHL, GPA, MPA: 12 MO. CLL: 6 MO.
<b>Other Criteria</b>	RA: INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RITUXIMAB-ARRX

## Products Affected

- RIABNI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	RHEUMATOID ARTHRITIS (RA): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. NHL, CLL: PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST.
<b>Coverage Duration</b>	RA: INITIAL: 6 MO, RENEWAL: 12 MO. NHL, WG, MPA: 12 MO. CLL: 6 MO.
<b>Other Criteria</b>	RA: INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# RITUXIMAB-PVVR

## Products Affected

- RUXIENCE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	RA: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. NHL, CLL: PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST.
Coverage Duration	NHL, GPA, MPA: 12 MONTHS. CLL: 6 MONTHS. RA: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	RA: INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO TWO PREFERRED AGENTS: HUMIRA, ENBREL, RINVOQ, XELJANZ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ROPEGINTERFERON ALFA-2B-NJFT

---

## Products Affected

- BESREMI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RUCAPARIB

---

## Products Affected

- RUBRACA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: ONE OF THE FOLLOWING: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RUXOLITINIB

---

## Products Affected

- JAKAFI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	MYELOFIBROSIS: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. POLYCYTHEMIA VERA, GVHD: 12 MONTHS.
<b>Other Criteria</b>	MYELOFIBROSIS: RENEWAL: PATIENT CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# SAPROPTERIN

---

## Products Affected

- *javygtor oral tablet, soluble*
- *sapropterin oral tablet, soluble*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL: 1 MONTH, RENEWAL 12 MONTHS.
<b>Other Criteria</b>	HYPERPHENYLALANINEMIA (HPA): INITIAL: NOT CONCURRENTLY USING PALYNZIQ. RENEWAL: 1) CONTINUES TO BENEFIT FROM TREATMENT, AND 2) NOT CONCURRENTLY USING PALYNZIQ.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# SARILUMAB

## Products Affected

- KEVZARA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	RHEUMATOID ARTHRITIS (RA): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	RA: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. POLYMYALGIA RHEUMATICA (PMR): 12 MONTHS.
Other Criteria	RA: INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# SECUKINUMAB

## Products Affected

- COSENTYX (2 SYRINGES)
- COSENTYX PEN (2 PENS)
- COSENTYX SUBCUTANEOUS SYRINGE 75 MG/0.5 ML
- COSENTYX UNOREADY PEN

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, OR FACE. NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN (CRP) LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PSO, HIDRADENITIS SUPPURATIVA (HS): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR A DERMATOLOGIST. ANKYLOSING SPONDYLITIS (AS), NR-AXSPA, ENTHESITIS-RELATED ARTHRITIS (ERA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
<b>Coverage Duration</b>	INITIAL: HS: 4 MONTHS, ALL OTHER INDICATIONS: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTI-RHEUMATIC DRUG). NR-AXSPA: TRIAL OF OR CONTRAINDICATION TO AN NSAID (NON-STEROIDAL ANTI-INFLAMMATORY DRUG). ERA: TRIAL OF OR CONTRAINDICATION TO ONE NSAID, SULFASALAZINE, OR METHOTREXATE. HS: NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGICS FOR HS OR OTHER IL-17 INHIBITORS FOR ANY INDICATION. RENEWAL: PSO, PSA, AS, NR-AXSPA, ERA: CONTINUES TO BENEFIT FROM THE MEDICATION. HS: 1) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGICS FOR HS OR OTHER IL-17 INHIBITORS FOR ANY INDICATION, AND 2) CONTINUES TO BENEFIT FROM THE MEDICATION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# SECUKINUMAB IV

## Products Affected

- COSENTYX INTRAVENOUS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN (CRP) LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR A DERMATOLOGIST. ANKYLOSING SPONDYLITIS (AS), NR-AXSPA: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
<b>Other Criteria</b>	INITIAL: PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTI-RHEUMATIC DRUG). NR-AXSPA: TRIAL OF OR CONTRAINDICATION TO AN NSAID (NON-STEROIDAL ANTI-INFLAMMATORY DRUG). RENEWAL: PSA, AS, NR-AXSPA: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# SELEXIPAG

## Products Affected

- UPTRAVI INTRAVENOUS
- UPTRAVI ORAL TABLET 1,000 MCG, 1,200 MCG, 1,400 MCG, 1,600 MCG, 200 MCG, 400 MCG, 600 MCG, 800 MCG
- UPTRAVI ORAL TABLETS, DOSE PACK

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	PAH: INITIAL: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING AGENTS FROM DIFFERENT DRUG CLASSES: 1) FORMULARY VERSION OF AN ORAL ENDOTHELIN RECEPTOR ANTAGONIST, 2) FORMULARY VERSION OF AN ORAL PHOSPHODIESTERASE TYPE-5 INHIBITOR, OR 3) FORMULARY VERSION OF AN ORAL CGMP STIMULATOR.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# SELINEXOR

---

## Products Affected

- XPOVIO ORAL TABLET 100 MG/WEEK (50 MG X 2), 40 MG/WEEK (40 MG X 1), 40MG TWICE WEEK (40 MG X 2), 60 MG/WEEK (60 MG X 1), 60MG TWICE WEEK (120 MG/WEEK), 80 MG/WEEK (40 MG X 2), 80MG TWICE WEEK (160 MG/WEEK)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# SELPERCATINIB

---

## Products Affected

- RETEVMO ORAL CAPSULE 40 MG,  
80 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# SELUMETINIB

---

## Products Affected

- KOSELUGO ORAL CAPSULE 10 MG,  
25 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# SILDENAFIL TABLET

## Products Affected

- *sildenafil (pulm.hypertension) oral tablet*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: AGES 18 YEARS OR OLDER: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS. AGES 1 TO 17 YEARS: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PAP GREATER THAN 20 MMHG, 2) PCWP OF 15 MMHG OR LESS, AND 3) PVR OF 3 WOOD UNITS OR GREATER.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	PAH: INITIAL/RENEWAL: NOT CONCURRENTLY OR INTERMITTENTLY TAKING ORAL ERECTILE DYSFUNCTION AGENTS (E.G. CIALIS, VIAGRA), ANY ORGANIC NITRATES IN ANY FORM, OR GUANYLATE CYCLASE STIMULATORS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No

# SIPONIMOD

## Products Affected

- MAYZENT ORAL TABLET 0.25 MG, 1 MG, 2 MG
- MAYZENT STARTER(FOR 1MG MAINT)
- MAYZENT STARTER(FOR 2MG MAINT)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	MULTIPLE SCLEROSIS: RENEWAL: 1) DEMONSTRATION OF CLINICAL BENEFIT COMPARED TO PRE-TREATMENT BASELINE AND 2) DOES NOT HAVE LYMPHOPENIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# SIROLIMUS PROTEIN-BOUND

---

## Products Affected

- FYARRO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# SODIUM OXYBATE-XYREM

## Products Affected

- *sodium oxybate*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: ALL INDICATIONS: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR SPECIALIST IN SLEEP MEDICINE
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
<b>Other Criteria</b>	INITIAL: EXCESSIVE DAYTIME SLEEPINESS (EDS) IN NARCOLEPSY: 1) NOT CURRENTLY TAKING A SEDATIVE HYPNOTIC AGENT, 2) FOR PATIENTS 18 YEARS OR OLDER: TRIAL OF OR CONTRAINDICATION TO A FORMULARY VERSION OF MODAFINIL, ARMODAFINIL, OR SOLRIAMFETOL AND ONE OTHER GENERIC STIMULANT INDICATED FOR EDS IN NARCOLEPSY, AND 3) FOR PATIENTS 7 TO 17 YEARS OF AGE: TRIAL OF OR CONTRAINDICATION TO ONE GENERIC STIMULANT INDICATED FOR EDS IN NARCOLEPSY. CATAPLEXY IN NARCOLEPSY: NOT CURRENTLY TAKING A SEDATIVE HYPNOTIC AGENT. RENEWAL (ALL INDICATIONS): 1) SUSTAINED IMPROVEMENT OF SYMPTOMS COMPARED TO BASELINE, AND 2) NOT CURRENTLY TAKING A SEDATIVE HYPNOTIC AGENT.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No

# SODIUM PHENYL BUTYRATE TABLETS

---

## Products Affected

- *sodium phenylbutyrate oral tablet*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	UREA CYCLE DISORDER (UCD): INITIAL: UCD IS CONFIRMED VIA ENZYMATIC, BIOCHEMICAL OR GENETIC TESTING.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	UCD: RENEWAL: CLINICAL BENEFIT FROM BASELINE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# SOFOSBUVIR/VELPATASVIR

## Products Affected

- EPCLUSA ORAL PELLETS IN PACKET 150-37.5 MG, 200-50 MG
- EPCLUSA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HCV RNA LEVEL WITHIN PAST 6 MONTHS.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
Other Criteria	1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, 2) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING MEDICATIONS: AMIODARONE, CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, HIV REGIMEN THAT CONTAINS EFAVIRENZ, ROSUVASTATIN AT DOSES ABOVE 10MG, TIPRANA VIR/RITONAVIR, TOPOTECAN, SOVALDI (AS A SINGLE AGENT), HARVONI, ZEPATIER, MAVYRET, OR VOSEVI, AND 3) PATIENTS WITH DECOMPENSATED CIRRHOSIS REQUIRE CONCURRENT RIBAVIRIN UNLESS RIBAVIRIN INELIGIBLE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# SOFOSBUVIR/VELPATASVIR/VOXILAPREVIR

## Products Affected

- VOSEVI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	HCV RNA LEVEL WITHIN PAST 6 MONTHS
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
<b>Other Criteria</b>	1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, 2) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING MEDICATIONS: AMIODARONE, CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, CYCLOSPORINE, PITAVASTATIN, PRAVASTATIN (DOSES ABOVE 40MG), ROSUVASTATIN, METHOTREXATE, MITOXANTRONE, IMATINIB, IRINOTECAN, LAPATINIB, SULFASALAZINE, TOPOTECAN, OR HIV REGIMEN THAT CONTAINS EFAVIRENZ, ATAZANAVIR, LOPINAVIR, TIPRANAVIR/RITONAVIR, SOVALDI (AS A SINGLE AGENT), EPCLUSA, HARVONI, ZEPATIER, OR MAVYRET, AND 3) DOES NOT HAVE MODERATE OR SEVERE HEPATIC IMPAIRMENT (CHILD-PUGH B OR C).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No

# SOLRIAMFETOL

---

## Products Affected

- SUNOSI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: EXCESSIVE DAYTIME SLEEPINESS (EDS) IN NARCOLEPSY: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR SPECIALIST IN SLEEP MEDICINE.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: EDS IN NARCOLEPSY: TRIED THE FORMULARY VERSION OF MODAFINIL OR ARMODAFINIL, AND ONE OTHER GENERIC STIMULANT INDICATED FOR EDS IN NARCOLEPSY. EDS IN OBSTRUCTIVE SLEEP APNEA (OSA): TRIED THE FORMULARY VERSION OF MODAFINIL OR ARMODAFINIL. RENEWAL: EDS IN NARCOLEPSY OR OSA: SUSTAINED IMPROVEMENT OF SYMPTOMS COMPARED TO BASELINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# SOMATROPIN - NORDITROPIN

---

## Products Affected

- NORDITROPIN FLEXPPO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	INITIAL/RENEWAL: ATHLETIC ENHANCEMENT, ANTI-AGING PURPOSES.
<b>Required Medical Information</b>	INITIAL: PEDIATRIC GROWTH HORMONE DEFICIENCY (GHD), IDIOPATHIC SHORT STATURE (ISS), SMALL FOR GESTATIONAL AGE (SGA), TURNER SYNDROME (TS), NOONAN SYNDROME: HEIGHT AT LEAST 2 STANDARD DEVIATIONS (SD) BELOW THE MEAN HEIGHT FOR NORMAL CHILDREN OF THE SAME AGE AND GENDER. PRADER WILLI SYNDROME (PWS): CONFIRMED GENETIC DIAGNOSIS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL/RENEWAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: ADULT GHD: GROWTH HORMONE DEFICIENCY ALONE OR ASSOCIATED WITH MULTIPLE HORMONE DEFICIENCIES (HYPOPITUITARISM), AS A RESULT OF PITUITARY DISEASES, HYPOTHALAMIC DISEASE, SURGERY, RADIATION THERAPY, TRAUMA, OR CONTINUATION OF THERAPY FROM CHILDHOOD ONSET GROWTH HORMONE DEFICIENCY. PEDIATRIC GHD, ISS, SGA, TS, NOONAN SYNDROME: OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF THE WRIST AND HAND. RENEWAL: PEDIATRIC GHD: 1) IMPROVEMENT WHILE ON THERAPY (I.E., INCREASED HEIGHT OR INCREASED GROWTH VELOCITY), AND 2) OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF THE WRIST AND HAND OR PATIENT HAS NOT COMPLETED PREPUBERTALGROWTH. ISS, SGA, TS, NOONAN SYNDROME: 1) IMPROVEMENT WHILE ON THERAPY (I.E., INCREASED HEIGHT OR INCREASED GROWTH VELOCITY), AND 2) OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF THE WRIST AND HAND. PWS: IMPROVEMENT IN BODY COMPOSITION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# SOMATROPIN - SEROSTIM

## Products Affected

- SEROSTIM SUBCUTANEOUS  
RECON SOLN 4 MG, 5 MG, 6 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	INITIAL/RENEWAL: ATHLETIC ENHANCEMENT, ANTI-AGING PURPOSES
<b>Required Medical Information</b>	INITIAL: HIV/WASTING: MEETS ONE OF THE FOLLOWING CRITERIA FOR WEIGHT LOSS: 1) 10% UNINTENTIONAL WEIGHT LOSS OVER 12 MONTHS, 2) 7.5% UNINTENTIONAL WEIGHT LOSS OVER 6 MONTHS, 3) 5% BODY CELL MASS (BCM) LOSS WITHIN 6 MONTHS, 4) BODY CELL MASS (BCM) LESS THAN 35% (MEN) OF TOTAL BODY WEIGHT AND BODY MASS INDEX (BMI) LESS THAN 27 KG PER METER SQUARED, 5) BCM LESS THAN 23% (WOMEN) OF TOTAL BODY WEIGHT AND A BODY MASS INDEX (BMI) LESS THAN 27 KG PER METER SQUARED, OR 6) BMI LESS THAN 18.5 KG PER METER SQUARED.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	HIV/WASTING: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST, NUTRITIONAL SUPPORT SPECIALIST, OR INFECTIOUS DISEASE SPECIALIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 3 MONTHS.
<b>Other Criteria</b>	HIV/WASTING: INITIAL: 1) CURRENTLY ON HIV ANTIRETROVIRAL THERAPY, AND 2) INADEQUATE RESPONSE TO ONE PREVIOUS THERAPY (E.G., MEGACE, APPETITE STIMULANTS, ANABOLIC STEROIDS). RENEWAL: 1) CURRENTLY ON HIV ANTIRETROVIRAL THERAPY, AND 2) CLINICAL BENEFIT IN MUSCLE MASS AND WEIGHT.
<b>Indications</b>	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# SONIDEGIB

---

## Products Affected

- ODOMZO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	LOCALLY ADVANCED BASAL CELL CARCINOMA (BCC); BASELINE SERUM CREATINE KINASE (CK) AND SERUM CREATININE LEVELS
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# SORAFENIB

---

## Products Affected

- *sorafenib*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# SOTORASIB

---

## Products Affected

- LUMAKRAS ORAL TABLET 120 MG,  
320 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# STIRIPENTOL

---

## Products Affected

- DIACOMIT ORAL CAPSULE 250 MG, 500 MG
- DIACOMIT ORAL POWDER IN PACKET 250 MG, 500 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	DRAVET SYNDROME: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# SUNITINIB

---

## Products Affected

- *sunitinib malate*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	GASTROINTESTINAL STROMAL TUMORS (GIST): TRIAL OF OR CONTRAINDICATION TO IMATINIB (GLEEVEC).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TADALAFIL - ADCIRCA, ALYQ

## Products Affected

- *alyq*
- *tadalafil (pulm. hypertension)*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	PAH: INITIAL/RENEWAL: NOT CONCURRENTLY OR INTERMITTENTLY TAKING ORAL ERECTILE DYSFUNCTION AGENTS (E.G. CIALIS, VIAGRA), ANY ORGANIC NITRATES IN ANY FORM, OR GUANYLATE CYCLASE STIMULATORS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TALAZOPARIB

## Products Affected

- TALZENNA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ADVANCED OR METASTATIC BREAST CANCER: PATIENT HAS BEEN TREATED WITH CHEMOTHERAPY IN THE NEOADJUVANT, ADJUVANT, OR METASTATIC SETTING. PATIENTS WITH HORMONE RECEPTOR (HR)-POSITIVE BREAST CANCER MUST HAVE ADDITIONAL PRIOR TREATMENT WITH ENDOCRINE THERAPY OR BE CONSIDERED INAPPROPRIATE FOR ENDOCRINE THERAPY. METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TALIMOGENE

## Products Affected

- IMLYGIC INJECTION SUSPENSION  
10EXP6 (1 MILLION) PFU/ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	UNRESECTABLE MELANOMA: 1) IMLYGIC TO BE INJECTED INTO CUTANEOUS, SUBCUTANEOUS, AND/OR NODAL LESIONS THAT ARE VISIBLE, PALPABLE, OR DETECTABLE BY ULTRASOUND GUIDANCE, 2) NOT CURRENTLY RECEIVING IMMUNOSUPPRESSIVE THERAPY, 3) NO HISTORY OF PRIMARY OR ACQUIRED IMMUNODEFICIENT STATES, LEUKEMIA, LYMPHOMA, OR AIDS, AND 4) NO CONCURRENT USE WITH PEMBROLIZUMAB, NIVOLUMAB, IPILIMUMAB, DABRAFENIB, TRAMETINIB, VEMURAFENIB, INTERLEUKIN-2, INTERFERON, DACARBAZINE, TEMOZOLOMIDE, PACLITAXEL, CARBOPLATIN, IMATINIB, MELPHALAN, IMIQUIMOD, OR RADIATION THERAPY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TALQUETAMAB-TGVS

---

## Products Affected

- TALVEY

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TASIMELTEON

---

## Products Affected

- *tasimelteon*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	LIFETIME
<b>Other Criteria</b>	NON-24 HOUR SLEEP-WAKE DISORDER: PATIENT IS LIGHT-INSENSITIVE OR HAS TOTAL BLINDNESS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TAZEMETOSTAT

---

## Products Affected

- TAZVERIK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TBO-FILGRASTIM

---

## Products Affected

- GRANIX

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	NON-MYELOID MALIGNANCIES: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	NON-MYELOID MALIGNANCIES: TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: NIVESTYM.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# TEBENTAFUSP-TEBN

---

## Products Affected

- KIMMTRAK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TECLISTAMAB-CQYV

---

## Products Affected

- TECVAYLI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TEDUGLUTIDE

---

## Products Affected

- GATTEX 30-VIAL

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	SHORT BOWEL SYNDROME (SBS): INITIAL/RENEWAL: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST
<b>Coverage Duration</b>	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS
<b>Other Criteria</b>	SBS: INITIAL: PATIENT IS DEPENDENT ON INTRAVENOUS PARENTERAL NUTRITION DEFINED AS REQUIRING PARENTERAL NUTRITION AT LEAST THREE TIMES PER WEEK. RENEWAL: ACHIEVED OR MAINTAINED A DECREASED NEED FOR PARENTERAL SUPPORT COMPARED TO BASELINE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TELOTRISTAT

---

## Products Affected

- XERMELO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	CARCINOID SYNDROME DIARRHEA: PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST OR GASTROENTEROLOGIST
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TEPOTINIB

---

## Products Affected

- TEPMETKO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TERIFLUNOMIDE

---

## Products Affected

- *teriflunomide*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TESAMORELIN

---

## Products Affected

- EGRIFTA SV

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	3 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TESTOSTERONE

## Products Affected

- *testosterone transdermal gel in metered-dose pump 12.5 mg/1.25 gram (1%), 20.25 mg/1.25 gram (1.62%)*
- *testosterone transdermal gel in packet 1% (25 mg/2.5gram), 1% (50 mg/5 gram)*
- *testosterone transdermal solution in metered pump w/lapp*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	MALE HYPOGONADISM: INITIAL: CONFIRMED BY: 1) AT LEAST TWO TOTAL SERUM TESTOSTERONE LEVELS OF LESS THAN 300 NG/DL TAKEN ON SEPARATE OCCASIONS, OR 2) FREE SERUM TESTOSTERONE LEVEL OF LESS THAN 5 NG/DL.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	MALE HYPOGONADISM: INITIAL: 1) 40 YEARS OR OLDER: PROSTATE SPECIFIC ANTIGEN HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING. RENEWAL: 1) 40 YEARS OR OLDER: PROSTATE SPECIFIC ANTIGEN HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING, AND 2) IMPROVED SYMPTOMS COMPARED TO BASELINE AND TOLERANCE TO TREATMENT.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# TESTOSTERONE CYPIONATE

---

## Products Affected

- *testosterone cypionate*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	MALE HYPOGONADISM: INITIAL: CONFIRMED BY: 1) AT LEAST TWO TOTAL SERUM TESTOSTERONE LEVELS OF LESS THAN 300 NG/DL TAKEN ON SEPARATE OCCASIONS, OR 2) FREE SERUM TESTOSTERONE LEVEL OF LESS THAN 5 NG/DL.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	MALE HYPOGONADISM: INITIAL: 1) 40 YEARS OR OLDER: PROSTATE SPECIFIC ANTIGEN HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING. RENEWAL: 1) 40 YEARS OR OLDER: PROSTATE SPECIFIC ANTIGEN HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING, AND 2) IMPROVED SYMPTOMS COMPARED TO BASELINE AND TOLERANCE TO TREATMENT.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TESTOSTERONE ENANTHATE

## Products Affected

- *testosterone enanthate*
- XYOSTED

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	MALE HYPOGONADISM: INITIAL: CONFIRMED BY: 1) AT LEAST TWO TOTAL SERUM TESTOSTERONE LEVELS OF LESS THAN 300 NG/DL TAKEN ON SEPARATE OCCASIONS, OR 2) FREE SERUM TESTOSTERONE LEVEL OF LESS THAN 5 NG/DL.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	MALE HYPOGONADISM: INITIAL/RENEWAL: 12 MO. ALL OTHER INDICATIONS: LIFETIME OF MEMBERSHIP IN PLAN.
<b>Other Criteria</b>	MALE HYPOGONADISM: INITIAL: 1) 40 YEARS OR OLDER: PROSTATE SPECIFIC ANTIGEN HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING. RENEWAL: 1) 40 YEARS OR OLDER: PROSTATE SPECIFIC ANTIGEN HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING, AND 2) IMPROVED SYMPTOMS COMPARED TO BASELINE AND TOLERANCE TO TREATMENT.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TETRABENAZINE

---

## Products Affected

- *tetrabenazine*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	HUNTINGTONS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR MOVEMENT DISORDER SPECIALIST
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TEZACAFTOR/IVACAFTOR

## Products Affected

- SYMDEKO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	CYSTIC FIBROSIS (CF); INITIAL: CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CYSTIC FIBROSIS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CYSTIC FIBROSIS EXPERT
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: LIFETIME
<b>Other Criteria</b>	CF: RENEWAL: MAINTAINED, IMPROVED, OR DEMONSTRATED LESS THAN EXPECTED DECLINE IN FEV1 OR BODY MASS INDEX (BMI), OR REDUCTION IN NUMBER OF PULMONARY EXACERBATIONS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# THALIDOMIDE

---

## Products Affected

- THALOMID

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TISOTUMAB VEDOTIN-TFTV

---

## Products Affected

- TIVDAK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TIVOZANIB

---

## Products Affected

- FOTIVDA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TOCILIZUMAB IV

## Products Affected

- ACTEMRA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	CORONAVIRUS DISEASE 2019 (COVID-19) IN HOSPITALIZED ADULTS
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS (SJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, OR IMMUNOLOGIST.
<b>Coverage Duration</b>	INITIAL: RA, PJIA, SJIA, GCA: 6 MONTHS. CRS: 1 MONTH. RENEWAL: RA, PJIA, SJIA, GCA: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: RA: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PJIA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ IR. SJIA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG). RENEWAL: RA, PJIA, SJIA: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.



<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TOCILIZUMAB SQ

---

## Products Affected

- ACTEMRA
- ACTEMRA ACTPEN

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS (SJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, OR IMMUNOLOGIST. SYSTEMIC SCLEROSIS-ASSOCIATED INTERSTITIAL LUNG DISEASE (SSC-ILD): PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR RHEUMATOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: RA: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PJIA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ IR. SJIA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG). SSC-ILD: DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., CONNECTIVE TISSUE DISEASE, DRUG TOXICITY, ASBESTOS OR BERYLLIUM EXPOSURE, HYPERSENSITIVITY PNEUMONITIS). RENEWAL: RA, PJIA, SJIA: CONTINUES TO BENEFIT FROM THE MEDICATION. SSC-ILD: CLINICAL MEANINGFUL IMPROVEMENT OR MAINTENANCE IN ANNUAL RATE OF DECLINE.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TOFACITINIB

## Products Affected

- XELJANZ
- XELJANZ XR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS), POLYARTICULAR COURSE JUVENILE IDIOPATHIC ARTHRITIS (PCJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF A PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PSA, PCJIA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD. UC: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPYRINE, METHOTREXATE, OR MESALAMINE. RENEWAL: RA, PSA, AS, PCJIA: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TOPICAL TRETINOIN

---

## Products Affected

- ALTRENO
- *tretinoin*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	COSMETIC INDICATIONS SUCH AS WRINKLES, PHOTOAGING, MELASMA.
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	ACNE VULGARIS: BRAND TOPICAL TRETINOIN REQUIRES TRIAL OF OR CONTRAINDICATION TO A GENERIC TOPICAL TRETINOIN PRODUCT.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TORIPALIMAB-TPZI

---

## Products Affected

- LOQTORZI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	NASOPHARYNGEAL CARCINOMA (NPC): FIRST LINE TREATMENT: 24 MOS, PREVIOUSLY TREATED: LIFETIME.
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TRAMETINIB

---

## Products Affected

- MEKINIST ORAL TABLET 0.5 MG, 2 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# TRAMETINIB SOLUTION

---

## Products Affected

- MEKINIST ORAL RECON SOLN

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	UNABLE TO SWALLOW MEKINIST TABLETS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TRASTUZUMAB HYALURONIDASE

---

## Products Affected

- HERCEPTIN HYLECTA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	ADJUVANT BREAST CANCER, METASTATIC BREAST CANCER: TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREFERRED AGENTS: HERZUMA, OGIVRI, ONTRUZANT, TRAZIMERA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TRASTUZUMAB-DKST

---

## Products Affected

- OGIVRI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TRASTUZUMAB-DTTB

---

## Products Affected

- ONTRUZANT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TRASTUZUMAB-PKRB

---

## Products Affected

- HERZUMA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TRASTUZUMAB-QYYP

---

## Products Affected

- TRAZIMERA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TREMELIMUMAB-ACTL

---

## Products Affected

- IMJUDO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	UHCC: 30 DAYS. METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC): 5 MONTHS.
<b>Other Criteria</b>	UNRESECTABLE HEPATOCELLULAR CARCINOMA (UHCC): HAS NOT RECEIVED PRIOR TREATMENT WITH IMJUDO.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TRIENTINE CAPSULE

---

## Products Affected

- *trientine oral capsule 250 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	WILSONS DISEASE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEPATOLOGIST OR GASTROENTEROLOGIST.
Coverage Duration	INITIAL: 12 MONTHS, RENEWAL: LIFETIME.
Other Criteria	WILSONS DISEASE: INITIAL: 1) LEIPZIG SCORE OF 4 OR GREATER, AND 2) TRIAL OF OR CONTRAINDICATION TO FORMULARY VERSION OF PENICILLAMINE TABLET. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# TRIFLURIDINE/TIPIRACIL

---

## Products Affected

- LONSURF ORAL TABLET 15-6.14 MG, 20-8.19 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TRIPTORELIN-TRELSTAR

---

## Products Affected

- TRELSTAR INTRAMUSCULAR  
SUSPENSION FOR  
RECONSTITUTION

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS.
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TUCATINIB

---

## Products Affected

- TUKYSA ORAL TABLET 150 MG, 50 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# UBROGEPANT

---

## Products Affected

- UBRELVY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	ACUTE MIGRAINE TREATMENT: INITIAL: TRIAL OF OR CONTRAINDICATION TO ONE TRIPTAN (E.G., SUMATRIPTAN, RIZATRIPTAN). RENEWAL: 1) IMPROVEMENT FROM BASELINE IN A VALIDATED ACUTE TREATMENT PATIENT-REPORTED OUTCOME QUESTIONNAIRE, OR 2) THERAPY WORKS CONSISTENTLY IN MAJORITY OF MIGRAINE ATTACKS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# UPADACITINIB

---

## Products Affected

- RINVOQ

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RA, AS, NR-AXSPA: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSA: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. AD: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST, ALLERGIST, OR IMMUNOLOGIST. UC, CD: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF A PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD. ATOPIC DERMATITIS: 1) ATOPIC DERMATITIS COVERING AT LEAST 10 PERCENT OF BODY SURFACE AREA OR ATOPIC DERMATITIS AFFECTING THE FACE, HEAD, NECK, HANDS, FEET, GROIN, OR INTERTRIGINOUS AREAS, 2) INTRACTABLE PRURITUS OR CRACKING/OOZING/BLEEDING OF AFFECTED SKIN, 3) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING: TOPICAL CORTICOSTEROID, TOPICAL CALCINEURIN INHIBITOR, TOPICAL PDE4 INHIBITOR, OR TOPICAL JAK INHIBITOR, AND 4) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGIC/JAK INHIBITOR FOR THE TREATMENT OF ATOPIC DERMATITIS. UC: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPYRINE, METHOTREXATE, OR MESALAMINE. NR-AXSPA: TRIAL OF OR CONTRAINDICATION TO AN NSAID (NON-STEROIDAL ANTI-INFLAMMATORY DRUG). CD: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPYRINE, METHOTREXATE, OR MESALAMINE. RENEWAL: RA, PSA, AS, NR-AXSPA: CONTINUES TO BENEFIT FROM THE MEDICATION. ATOPIC DERMATITIS: 1) IMPROVEMENT WHILE ON THERAPY, AND 2) NO CONCURRENT USE WITH OTHER</p>
	SYSTEMIC BIOLOGIC/JAK INHIBITOR FOR THE TREATMENT OF ATOPIC DERMATITIS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No

# USTEKINUMAB

## Products Affected

- STELARA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, OR FACE.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG). PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. CD, UC: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPYRINE, METHOTREXATE, OR MESALAMINE. RENEWAL: PSA, PSO: CONTINUES TO BENEFIT FROM THE MEDICATION.



<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# USTEKINUMAB IV

---

## Products Affected

- STELARA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	2 MONTHS
Other Criteria	CD, UC: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPYRINE, METHOTREXATE, OR MESALAMINE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# VANDETANIB

---

## Products Affected

- CAPRELSA ORAL TABLET 100 MG,  
300 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	CURRENTLY STABLE ON CAPRELSA REQUIRES NO EXTRA CRITERIA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# VEMURAFENIB

---

## Products Affected

- ZELBORAF

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	MELANOMA: ZELBORAF WILL BE USED ALONE OR IN COMBINATION WITH COTELLIC
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# VENETOCLAX

---

## Products Affected

- VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG
- VENCLEXTA STARTING PACK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# VERICIGUAT

## Products Affected

- VERQUVO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL/RENEWAL:12 MONTHS.
<b>Other Criteria</b>	HEART FAILURE (HF): INITIAL: TRIAL OF OR CONTRAINDICATION TO 1) ONE AGENT FROM ANY OF THE FOLLOWING STANDARD OF CARE CLASSES: A) ACE INHIBITOR, ARB, OR ARNI, B) BETA BLOCKER (I.E., BISOPROLOL, CARVEDILOL, METOPROLOL SUCCINATE), OR C) ALDOSTERONE ANTAGONIST (I.E., SPIRONOLACTONE, EPLERENONE), AND 2) ONE PREFERRED SGLT-2 INHIBITOR (E.G., FARXIGA, XIGDUO XR, JARDIANCE). INITIAL/RENEWAL: NOT CONCURRENTLY TAKING LONG-ACTING NITRATES OR NITRIC OXIDE DONORS, RIOCIGUAT, OR PDE-5 INHIBITORS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# VIGABATRIN

---

## Products Affected

- *vigabatrin*
- *vigadrone*
- *vigpoder*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	REFRACTORY COMPLEX PARTIAL SEIZURES (CPS), INFANTILE SPASMS: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	CPS: TRIAL OF OR CONTRAINDICATION TO TWO ANTIEPILEPTIC AGENTS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# VISMODEGIB

---

## Products Affected

- ERIVEDGE

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# VORICONAZOLE SUSPENSION

---

## Products Affected

- *voriconazole oral suspension for reconstitution*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	CANDIDA INFECTIONS: 3 MOS. ALL OTHER INDICATIONS: 6 MOS.
<b>Other Criteria</b>	CANDIDA INFECTIONS: TRIAL OF OR CONTRAINDICATION TO FLUCONAZOLE. ALL INDICATIONS: INABILITY TO SWALLOW TABLETS OR AN INDICATION FOR ESOPHAGEAL CANDIDIASIS. CONTINUATION OF THERAPY AFTER HOSPITAL DISCHARGE REQUIRES NO EXTRA CRITERIA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ZANUBRUTINIB

---

## Products Affected

- BRUKINSA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ZURANOLONE

---

## Products Affected

- ZURZUVAE ORAL CAPSULE 20 MG, 25 MG, 30 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	14 DAYS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



## INDEX

<i>abiraterone</i> .....	6	BETASERON SUBCUTANEOUS KIT	
ACTEMRA.....	360, 362	.....	178
ACTEMRA ACTPEN.....	362	<i>bexarotene</i> .....	46
ACTHAR.....	67	<i>bortezomib injection</i> .....	48
ACTIMMUNE.....	180	<i>bosentan</i> .....	49
ADEMPAS.....	296	BOSULIF ORAL CAPSULE 100 MG,	
AJOVY AUTOINJECTOR.....	132	50 MG.....	50
AJOVY SYRINGE.....	132	BOSULIF ORAL TABLET 100 MG,	
AKEEGA.....	236	400 MG, 500 MG.....	50
ALECENSA.....	13	BRAFTOVI.....	107
ALTRENO.....	366	BRUKINSA.....	394
ALUNBRIG ORAL TABLET 180 MG,		<i>butalbital-acetaminophen-caff oral tablet</i>	149
30 MG, 90 MG.....	51	<i>butalbital-aspirin-caffeine</i> .....	149
ALUNBRIG ORAL TABLETS,DOSE		CABLIVI INJECTION KIT.....	58
PACK.....	51	CABOMETYX ORAL TABLET 20	
<i>alyq</i> .....	338	MG, 40 MG, 60 MG.....	55
<i>amabelz</i> .....	154	CALQUENCE (ACALABRUTINIB	
<i>ambrisentan</i> .....	15	MAL).....	8
<i>apomorphine</i> .....	19	CAPRELSA ORAL TABLET 100 MG,	
<i>armodafinil</i> .....	228	300 MG.....	387
<i>ascomp with codeine</i> .....	149	<i>carglumic acid</i> .....	60
AUGTYRO.....	287	CAYSTON.....	32
AUSTEDO ORAL TABLET 12 MG, 6		CERDELGA.....	101
MG, 9 MG.....	84	<i>chlorzoxazone oral tablet 500 mg</i> .....	164
AUSTEDO XR ORAL TABLET		CINQAIR.....	288
EXTENDED RELEASE 24 HR 12		CINRYZE.....	52
MG, 24 MG, 6 MG.....	84	<i>codeine-butalbital-asa-caff</i> .....	149
AUSTEDO XR TITRATION		<i>colchicine oral tablet</i> .....	66
KT(WK1-4).....	84	COMETRIQ ORAL CAPSULE 100	
AVONEX INTRAMUSCULAR PEN		MG/DAY(80 MG X1-20 MG X1), 140	
INJECTOR KIT.....	177	MG/DAY(80 MG X1-20 MG X3), 60	
AVONEX INTRAMUSCULAR		MG/DAY (20 MG X 3/DAY).....	54
SYRINGE KIT.....	177	COPAXONE SUBCUTANEOUS	
AYVAKIT.....	28	SYRINGE 20 MG/ML, 40 MG/ML.....	140
BALVERSA ORAL TABLET 3 MG, 4		COPIKTRA.....	95
MG, 5 MG.....	115	CORTROPHIN GEL.....	67
<i>bendamustine intravenous recon soln</i> .....	39	COSENTYX (2 SYRINGES).....	311
BENDAMUSTINE INTRAVENOUS		COSENTYX INTRAVENOUS.....	313
SOLUTION.....	39	COSENTYX PEN (2 PENS).....	311
BENDEKA.....	39	COSENTYX SUBCUTANEOUS	
BENLYSTA SUBCUTANEOUS.....	36	SYRINGE 75 MG/0.5 ML.....	311
BESREMI.....	306	COSENTYX UNOREADY PEN.....	311
<i>betaine</i> .....	42	COTELLIC.....	65

<i>cyclobenzaprine oral tablet 10 mg, 5 mg</i> ..	164	EMGALITY SYRINGE	
<i>cyproheptadine oral syrup</i> .....	148	SUBCUTANEOUS SYRINGE 120	
CYRAMZA.....	284	MG/ML, 300 MG/3 ML (100 MG/ML	
CYSTARAN.....	71	X 3).....	135
<i>dalfampridine</i> .....	75	ENBREL.....	118
DANYELZA.....	230	ENBREL MINI.....	118
DAURISMO ORAL TABLET 100		ENBREL SURECLICK.....	118
MG, 25 MG.....	139	ENDARI.....	201
<i>deferasirox</i> .....	79	EPCLUSA ORAL PELLETS IN	
<i>deferiprone</i> .....	81	PACKET 150-37.5 MG, 200-50 MG.....	325
DIACOMIT ORAL CAPSULE 250		EPCLUSA ORAL TABLET.....	325
MG, 500 MG.....	336	EPIDIOLEX.....	56
DIACOMIT ORAL POWDER IN		EPKINLY.....	112
PACKET 250 MG, 500 MG.....	336	ERBITUX.....	62
<i>diclofenac sodium topical gel 3 %</i> .....	85	ERIVEDGE.....	392
<i>diclofenac sodium topical solution in</i>		ERLEADA ORAL TABLET 240 MG,	
<i>metered-dose pump</i> .....	86	60 MG.....	18
<i>dimethyl fumarate oral capsule, delayed</i>		<i>erlotinib oral tablet 100 mg, 150 mg, 25</i>	
<i>release (drlec) 120 mg, 120 mg (14)- 240</i>		<i>mg</i> .....	116
<i>mg (46), 240 mg</i> .....	87	<i>estradiol oral</i> .....	153
<i>diphenhydramine hcl oral elixir</i> .....	165	<i>estradiol transdermal patch semiweekly</i> ...	153
<i>diphenoxylate-atropine oral tablet</i> .....	166	<i>estradiol transdermal patch weekly</i> .....	153
<i>dipyridamole oral</i> .....	151	<i>estradiol-norethindrone acet oral tablet</i>	
<i>disopyramide phosphate oral capsule</i> .....	152	<i>0.5-0.1 mg</i> .....	154
DOPTELET (10 TAB PACK).....	29	<i>everolimus (antineoplastic) oral tablet 10</i>	
DOPTELET (15 TAB PACK).....	29	<i>mg, 2.5 mg, 5 mg, 7.5 mg</i> .....	120
DOPTELET (30 TAB PACK).....	29	<i>everolimus (antineoplastic) oral tablet for</i>	
<i>dotti</i> .....	153	<i>suspension</i> .....	121
<i>dronabinol</i> .....	90	EVRYSDI.....	301
<i>droxidopa</i> .....	91	EXKIVITY.....	223
DUAVEE.....	155	FARYDAK.....	255
DUPIXENT PEN.....	92	FASENRA.....	40
DUPIXENT SYRINGE.....	92	FASENRA PEN.....	40
EGRIFTA SV.....	351	<i>fentanyl citrate buccal lozenge on a</i>	
ELIGARD.....	197	<i>handle</i> .....	125
ELIGARD (3 MONTH).....	197	FERRIPROX (2 TIMES A DAY).....	81
ELIGARD (4 MONTH).....	197	FERRIPROX ORAL SOLUTION.....	81
ELIGARD (6 MONTH).....	197	<i>ingolimod</i> .....	130
ELREXFIO 44 MG/1.1 ML VIAL		FINTEPLA.....	124
OUTER, SUV, P/F.....	102	FOTIVDA.....	359
ELREXFIO SUBCUTANEOUS		FRUZAQLA ORAL CAPSULE 1 MG,	
SOLUTION 40 MG/ML.....	102	5 MG.....	133
EMGALITY PEN.....	135	FULPHILA.....	261
		FYARRO.....	321
		<i>fyavolv</i> .....	159

GALAFOLD.....	220	IMBRUVICA ORAL CAPSULE 140	
GAMIFANT.....	104	MG, 70 MG.....	170
GATTEX 30-VIAL.....	347	IMBRUVICA ORAL SUSPENSION...	170
GAVRETO.....	279	IMBRUVICA ORAL TABLET 140	
<i>gefitinib</i> .....	137	MG, 280 MG, 420 MG.....	170
GILENYA ORAL CAPSULE 0.25 MG		IMJUDO.....	375
.....	130	IMLYGIC INJECTION SUSPENSION	
GILOTRIF.....	12	10EXP6 (1 MILLION) PFU/ML.....	340
<i>glatiramer subcutaneous syringe 20</i>		IMPAVIDO.....	222
<i>mg/ml, 40 mg/ml</i> .....	140	INBRIJA INHALATION CAPSULE,	
<i>glatopa subcutaneous syringe 20 mg/ml,</i>		W/INHALATION DEVICE.....	200
<i>40 mg/ml</i> .....	140	<i>indomethacin oral capsule 25 mg, 50 mg..</i>	167
<i>glyburide</i> .....	157	<i>indomethacin oral capsule, extended</i>	
<i>glyburide micronized</i> .....	157	<i>release</i> .....	167
<i>glyburide-metformin</i> .....	157	<i>infliximab</i> .....	175
GRANIX.....	344	INLYTA ORAL TABLET 1 MG, 5	
HAEGARDA SUBCUTANEOUS		MG.....	30
RECON SOLN 2,000 UNIT, 3,000		INQOVI.....	78
UNIT.....	53	INREBIC.....	123
HARVONI ORAL PELLETS IN		IWILFIN.....	96
PACKET 33.75-150 MG, 45-200 MG...	191	JAKAFI.....	308
HARVONI ORAL TABLET.....	191	<i>javygtor oral tablet, soluble</i> .....	309
HERCEPTIN HYLECTA.....	370	JAYPIRCA ORAL TABLET 100 MG,	
HERZUMA.....	373	50 MG.....	274
HUMIRA PEN.....	10	JEMPERLI.....	89
HUMIRA PEN CROHNS-UC-HS		<i>jinteli</i> .....	159
START.....	10	JUXTAPID ORAL CAPSULE 10 MG,	
HUMIRA PEN PSOR-UVEITS-ADOL		20 MG, 30 MG, 40 MG, 5 MG, 60 MG	206
HS.....	10	KALYDECO.....	182
HUMIRA SUBCUTANEOUS		KERENDIA.....	129
SYRINGE KIT 40 MG/0.8 ML.....	10	KESIMPTA PEN.....	243
HUMIRA(CF).....	10	<i>ketorolac oral</i> .....	158
HUMIRA(CF) PEDI CROHNS		KEVZARA.....	310
STARTER.....	10	KEYTRUDA.....	266
HUMIRA(CF) PEN.....	10	KIMMTRAK.....	345
HUMIRA(CF) PEN CROHNS-UC-HS.	10	KINERET.....	17
HUMIRA(CF) PEN PEDIATRIC UC...	10	KISQALI FEMARA CO-PACK ORAL	
HUMIRA(CF) PEN PSOR-UV-ADOL		TABLET 200 MG/DAY(200 MG X 1)-	
HS.....	10	2.5 MG, 400 MG/DAY(200 MG X 2)-	
IBRANCE.....	254	2.5 MG, 600 MG/DAY(200 MG X 3)-	
<i>icatibant</i> .....	171	2.5 MG.....	292
ICLUSIG.....	276	KISQALI ORAL TABLET 200	
IDHIFA.....	106	MG/DAY (200 MG X 1), 400 MG/DAY	
<i>imatinib oral tablet 100 mg, 400 mg</i> .....	173	(200 MG X 2), 600 MG/DAY (200 MG	
		X 3).....	291

KORLYM.....	219	MAYZENT STARTER(FOR 1MG MAINT).....	320
KOSELUGO ORAL CAPSULE 10 MG, 25 MG.....	317	MAYZENT STARTER(FOR 2MG MAINT).....	320
KRAZATI.....	9	<i>megestrol oral suspension 400 mg/10 ml (40 mg/ml)</i> .....	168
KYNMOBI SUBLINGUAL FILM 10 MG, 10-15-20-25-30 MG, 15 MG, 20 MG, 25 MG, 30 MG.....	20	<i>megestrol oral tablet</i> .....	168
<i>lanreotide</i> .....	188	MEKINIST ORAL RECON SOLN.....	369
<i>lapatinib</i> .....	189	MEKINIST ORAL TABLET 0.5 MG, 2 MG.....	368
<i>lenalidomide</i> .....	192	MEKTOVI.....	47
LENVIMA.....	193	<i>methocarbamol oral tablet 500 mg, 750 mg</i> .....	164
<i>leuprolide (3 month)</i> .....	196	<i>mifepristone oral tablet 300 mg</i> .....	219
<i>leuprolide subcutaneous kit</i> .....	195	<i>miglustat</i> .....	221
<i>lidocaine hcl mucous membrane solution 4 % (40 mg/ml)</i> .....	205	<i>mimvey</i> .....	154
<i>lidocaine topical adhesive patch,medicated 5 %</i> .....	203	<i>modafinil oral tablet 100 mg, 200 mg</i> .....	228
<i>lidocaine topical ointment</i> .....	202	<i>morphine concentrate oral solution</i> .....	147
<i>lidocaine-prilocaine topical cream</i> .....	204	MOUNJARO.....	143
LONSURF ORAL TABLET 15-6.14 MG, 20-8.19 MG.....	377	MVASI.....	44
LOQTORZI.....	367	NATPARA.....	256
LORBRENA ORAL TABLET 100 MG, 25 MG.....	209	NERLYNX.....	231
LUMAKRAS ORAL TABLET 120 MG, 320 MG.....	335	NEULASTA ONPRO.....	262
LUNSUMIO.....	225	NINLARO.....	184
LUPRON DEPOT.....	198	<i>nitisinone</i> .....	238
LUPRON DEPOT (3 MONTH).....	198	NIVESTYM.....	126
LUPRON DEPOT (4 MONTH).....	198	NORDITROPIN FLEXPRO.....	329
LUPRON DEPOT (6 MONTH).....	198	<i>norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg</i> .....	159
LYBALVI.....	244	NOXAFIL ORAL SUSP,DELAYED RELEASE FOR RECON.....	278
<i>lyllana</i> .....	153	NUBEQA.....	76
LYNPARZA.....	245	NUCALA SUBCUTANEOUS AUTO-INJECTOR.....	215
LYTGOBI.....	134	NUCALA SUBCUTANEOUS RECON SOLN.....	215
MARGENZA.....	213	NUCALA SUBCUTANEOUS SYRINGE 100 MG/ML, 40 MG/0.4 ML.....	215
MAVENCLAD (10 TABLET PACK).....	63	NUPLAZID.....	271
MAVENCLAD (4 TABLET PACK).....	63	NURTEC ODT.....	294
MAVENCLAD (5 TABLET PACK).....	63	NYVEPRIA.....	259
MAVENCLAD (6 TABLET PACK).....	63	OICALIVA.....	241
MAVENCLAD (7 TABLET PACK).....	63	OCREVUS.....	242
MAVENCLAD (8 TABLET PACK).....	63	ODOMZO.....	333
MAVENCLAD (9 TABLET PACK).....	63		
MAYZENT ORAL TABLET 0.25 MG, 1 MG, 2 MG.....	320		



OFEV .....	233	<i>pirfenidone oral tablet 267 mg, 534 mg,</i>	
OGIVRI .....	371	<i>801 mg</i> .....	272
OGSIVEO .....	237	PLEGRIDY SUBCUTANEOUS PEN	
OJJAARA .....	224	INJECTOR 125 MCG/0.5 ML, 63	
OLUMIANT .....	33	MCG/0.5 ML- 94 MCG/0.5 ML .....	179
ONTRUZANT .....	372	PLEGRIDY SUBCUTANEOUS	
ONUREG .....	31	SYRINGE 125 MCG/0.5 ML, 63	
OPDIVO .....	239	MCG/0.5 ML- 94 MCG/0.5 ML .....	179
OPDUALAG .....	240	POMALYST .....	275
OPSUMIT .....	212	<i>posaconazole oral</i> .....	277
ORENCIA .....	3	PREMARIN ORAL .....	150
ORENCIA (WITH MALTOSE) .....	1	PREMPHASE .....	156
ORENCIA CLICKJECT .....	3	PREMPRO .....	156
ORFADIN ORAL SUSPENSION .....	238	PREVYMIS INTRAVENOUS	
ORGOVYX .....	286	SOLUTION 240 MG/12 ML, 480	
ORILISSA ORAL TABLET 150 MG,		MG/24 ML .....	194
200 MG .....	98	PREVYMIS ORAL .....	194
ORKAMBI ORAL GRANULES IN		PROMACTA ORAL POWDER IN	
PACKET .....	210	PACKET 12.5 MG, 25 MG .....	103
ORKAMBI ORAL TABLET .....	210	PROMACTA ORAL TABLET 12.5	
ORSERDU ORAL TABLET 345 MG,		MG, 25 MG, 50 MG, 75 MG .....	103
86 MG .....	97	<i>promethazine injection solution</i> .....	161
OTEZLA .....	21	<i>promethazine oral</i> .....	161
OTEZLA STARTER .....	21	<i>promethazine rectal</i> .....	161
<i>oxandrolone</i> .....	252	<i>promethegan rectal suppository 12.5 mg,</i>	
OXLUMO .....	211	<i>25 mg</i> .....	161
OZEMPIC SUBCUTANEOUS PEN		<i>pyrimethamine</i> .....	281
INJECTOR 0.25 MG OR 0.5 MG (2		QINLOCK .....	298
MG/3 ML), 0.25 MG OR 0.5 MG(2		<i>quinine sulfate</i> .....	282
MG/1.5 ML), 1 MG/DOSE (4 MG/3		QULIPTA .....	27
ML), 2 MG/DOSE (8 MG/3 ML) .....	142	RAVICTI .....	144
PALYNZIQ .....	264	RELEUKO .....	127
<i>paroxetine hcl oral suspension</i> .....	169	RETACRIT INJECTION SOLUTION	
<i>paroxetine hcl oral tablet</i> .....	169	10,000 UNIT/ML, 2,000 UNIT/ML,	
<i>pazopanib</i> .....	258	20,000 UNIT/2 ML, 20,000 UNIT/ML,	
PEGASYS .....	263	3,000 UNIT/ML, 4,000 UNIT/ML,	
PEMAZYRE .....	267	40,000 UNIT/ML .....	113
<i>penicillamine oral tablet</i> .....	268	RETEVMO ORAL CAPSULE 40 MG,	
<i>phenobarbital</i> .....	160	80 MG .....	316
PIQRAY ORAL TABLET 200		REZLIDHIA .....	246
MG/DAY (200 MG X 1), 250 MG/DAY		REZUROCK .....	37
(200 MG X1-50 MG X1), 300 MG/DAY		RIABNI .....	304
(150 MG X 2) .....	14	RINVOQ .....	381
<i>pirfenidone oral capsule</i> .....	272	RITUXAN HYCELA .....	302

ROZLYTREK ORAL CAPSULE 100 MG, 200 MG.....	108	TAFINLAR ORAL CAPSULE.....	72
ROZLYTREK ORAL PELLETS IN PACKET.....	109	TAFINLAR ORAL TABLET FOR SUSPENSION.....	73
RUBRACA.....	307	TAGRISSE.....	251
RUXIENCE.....	305	TAKHZYRO SUBCUTANEOUS SOLUTION.....	187
RYBELSUS.....	142	TAKHZYRO SUBCUTANEOUS SYRINGE 150 MG/ML, 300 MG/2 ML (150 MG/ML).....	187
RYBREVANT.....	16	TALTZ AUTOINJECTOR.....	185
RYDAPT.....	218	TALTZ SYRINGE.....	185
<i>sajazir</i> .....	171	TALVEY.....	341
<i>sapropterin oral tablet, soluble</i> .....	309	TALZENNA.....	339
SCSEMBLIX ORAL TABLET 20 MG, 40 MG.....	23	TASCENSO ODT.....	131
<i>scopolamine base</i> .....	163	TASIGNA ORAL CAPSULE 150 MG, 200 MG, 50 MG.....	232
SEROSTIM SUBCUTANEOUS RECON SOLN 4 MG, 5 MG, 6 MG....	331	<i>tasimelteon</i> .....	342
SIGNIFOR.....	257	TAZVERIK.....	343
<i>sildenafil (pulm.hypertension) oral tablet</i>	318	TECVAYLI.....	346
SIRTURO.....	35	TEPMETKO.....	349
SKYRIZI.....	299	<i>teriflunomide</i> .....	350
<i>sodium oxybate</i> .....	322	<i>testosterone cypionate</i> .....	353
<i>sodium phenylbutyrate oral tablet</i> .....	324	<i>testosterone enanthate</i> .....	354
SOMATULINE DEPOT SUBCUTANEOUS SYRINGE 120 MG/0.5 ML, 60 MG/0.2 ML, 90 MG/0.3 ML.....	188	<i>testosterone transdermal gel in metered- dose pump 12.5 mg/ 1.25 gram (1%), 20.25 mg/1.25 gram (1.62%)</i> .....	352
SOMAVERT.....	265	<i>testosterone transdermal gel in packet 1% (25 mg/2.5gram), 1% (50 mg/5 gram)</i> ...	352
<i>sorafenib</i> .....	334	<i>testosterone transdermal solution in metered pump w/lapp</i> .....	352
SPRAVATO.....	117	<i>tetrabenazine</i> .....	355
SPRYCEL ORAL TABLET 100 MG, 140 MG, 20 MG, 50 MG, 70 MG, 80 MG.....	77	THALOMID.....	357
STELARA.....	384, 386	TIBSOVO.....	183
STIVARGA.....	285	TIVDAK.....	358
STRENSIQ.....	24	TRACLEER ORAL TABLET FOR SUSPENSION.....	49
<i>sunitinib malate</i> .....	337	TRAZIMERA.....	374
SUNOSI.....	328	TRELSTAR INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION.....	378
SYMDEKO.....	356	TREMFYA.....	146
SYMLINPEN 120.....	280	<i>tretinoin</i> .....	366
SYMLINPEN 60.....	280	<i>trientine oral capsule 250 mg</i> .....	376
SYMPAZAN.....	64	TRIKAFTA ORAL TABLETS, SEQUENTIAL.....	100
SYNAREL.....	226		
SYNRIBO.....	247		
TABRECTA.....	59		
<i>taдалafil (pulm. hypertension)</i> .....	338		

TRULICITY .....	141	XELJANZ .....	364
TRUQAP .....	57	XELJANZ XR .....	364
TRUSELTIQ .....	174	XERMELO .....	348
TRUXIMA .....	303	XGEVA .....	83
TUKYSA ORAL TABLET 150 MG, 50 MG .....	379	XIFAXAN ORAL TABLET 200 MG, 550 MG .....	293
TURALIO .....	270	XOLAIR SUBCUTANEOUS AUTO- INJECTOR 300 MG/2 ML .....	248
TYSABRI .....	229	XOLAIR SUBCUTANEOUS RECON SOLN .....	248
UBRELVY .....	380	XOLAIR SUBCUTANEOUS SYRINGE .....	248
UDENYCA .....	260	XOSPATA .....	138
UDENYCA AUTOINJECTOR .....	260	XPOVIO ORAL TABLET 100 MG/WEEK (50 MG X 2), 40 MG/WEEK (40 MG X 1), 40MG TWICE WEEK (40 MG X 2), 60 MG/WEEK (60 MG X 1), 60MG TWICE WEEK (120 MG/WEEK), 80 MG/WEEK (40 MG X 2), 80MG TWICE WEEK (160 MG/WEEK) .....	315
UDENYCA ONBODY .....	260	XTANDI ORAL CAPSULE .....	110
UPTRAVI INTRAVENOUS .....	314	XTANDI ORAL TABLET 40 MG, 80 MG .....	110
UPTRAVI ORAL TABLET 1,000 MCG, 1,200 MCG, 1,400 MCG, 1,600 MCG, 200 MCG, 400 MCG, 600 MCG, 800 MCG .....	314	XYOSTED .....	354
UPTRAVI ORAL TABLETS,DOSE PACK .....	314	<i>yargesa</i> .....	221
VALCHLOR .....	214	YERVOY .....	181
VANFLYTA .....	283	YONSA .....	7
VEGZELMA .....	43	ZARXIO .....	128
VELCADE .....	48	ZEJULA ORAL CAPSULE .....	235
VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG .....	389	ZEJULA ORAL TABLET .....	235
VENCLEXTA STARTING PACK .....	389	ZELBORAF .....	388
VERQUVO .....	390	ZIRABEV .....	45
VERZENIO .....	5	ZOLADEX .....	145
<i>vigabatrin</i> .....	391	ZTALMY .....	136
<i>vigadrone</i> .....	391	ZTLIDO .....	203
<i>vigpoder</i> .....	391	ZURZUVAE ORAL CAPSULE 20 MG, 25 MG, 30 MG .....	395
VITRAKVI ORAL CAPSULE 100 MG, 25 MG .....	190	ZYDELIG .....	172
VITRAKVI ORAL SOLUTION .....	190	ZYKADIA .....	61
VIZIMPRO .....	74	ZYNLONTA .....	208
VONJO .....	253	ZYNYZ .....	290
<i>voriconazole oral suspension for reconstitution</i> .....	393		
VOSEVI .....	326		
VOWST .....	122		
VUMERITY .....	88		
WELIREG .....	38		
XALKORI ORAL CAPSULE .....	69		
XALKORI ORAL PELLETT 150 MG, 20 MG, 50 MG .....	70		