ABATACEPT IV

Products Affected

• ORENCIA (WITH MALTOSE)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST.
Coverage Duration	RA, PJIA, PSA: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. AGVHD: 12 MONTHS.
Other Criteria	INITIAL: RA: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PJIA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ IR. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, TREMFYA, XELJANZ, RINVOQ, SKYRIZI. RENEWAL: RA, PJIA, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ABATACEPT SQ

Products Affected

- ORENCIA
- ORENCIA CLICKJECT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: RA: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PJIA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ IR. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, COSENTYX, ENBREL, TREMFYA, XELJANZ, RINVOQ, SKYRIZI. RENEWAL: RA, PJIA, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

ABEMACICLIB

Products Affected

• VERZENIO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ABIRATERONE

Products Affected

• abiraterone

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC HIGH-RISK CASTRATION-SENSITIVE PROSTATE CANCER (MCSPC), METASTATIC CASTRATION- RESISTANT PROSTATE CANCER (MCRPC): 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ABIRATERONE SUBMICRONIZED

Products Affected

• YONSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ACALABRUTINIB

Products Affected

• CALQUENCE (ACALABRUTINIB MAL)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT BRUKINSA, WHERE INDICATIONS ALIGN.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ADAGRASIB

Products Affected

• KRAZATI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ADALIMUMAB

Products Affected

- HUMIRA PEN
- HUMIRA PEN CROHNS-UC-HS START
- HUMIRA PEN PSOR-UVEITS-ADOL HS
- HUMIRA SUBCUTANEOUS SYRINGE KIT 40 MG/0.8 ML
- HUMIRA(CF)

- HUMIRA(CF) PEDI CROHNS STARTER
- HUMIRA(CF) PEN
- HUMIRA(CF) PEN CROHNS-UC-HS
- HUMIRA(CF) PEN PEDIATRIC UC
- HUMIRA(CF) PEN PSOR-UV-ADOL HS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, OR GENITAL AREA.
Age Restrictions	
Prescriber Restrictions	INITIAL: RA, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA), ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST. UVEITIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OPTHALMOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE- MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PJIA, PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD. PSO: TRIAL OF OR CONTRAINDICATION TO ONE DMARD. PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. CD, UC: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPURINE, METHOTREXATE, OR MESALAMINE. UVEITIS: NO ISOLATED ANTERIOR UVEITIS. RENEWAL: RA, PJIA, PSA, AS, PSO, HIDRADENITIS SUPPURATIVA, UVEITIS: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

AFATINIB

Products Affected

• GILOTRIF

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS.
Other Criteria	METASTATIC NSCLC WITH EGFR MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

AGALSIDASE BETA

Products Affected

• FABRAZYME

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	FABRY DISEASE: INITIAL: 1) SYMPTOMATIC OR EVIDENCE OF INJURY FROM GL-3 TO THE KIDNEY, HEART, OR CENTRAL NERVOUS SYSTEM RECOGNIZED BY LABORATORY, HISTOLOGICAL, OR IMAGING FINDINGS, AND 2) ONE OF THE FOLLOWING: (A) FEMALES: GALACTOSIDASE ALPHA (GLA) GENE MUTATION VIA GENETIC TESTING, OR (B) MALES: ENZYME ASSAY INDICATING ALPHA GALACTOSIDASE A DEFICIENCY OR GLA GENE MUTATION VIA GENETIC TESTING.
Age Restrictions	
Prescriber Restrictions	FABRY DISEASE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH NEPHROLOGIST, CARDIOLOGIST, OR SPECIALIST IN GENETICS OR INHERITED METABOLIC DISORDERS.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	FABRY DISEASE: INITIAL: NO CONCURRENT USE WITH ANOTHER FABRY DISEASE THERAPY. RENEWAL: 1) DEMONSTRATED IMPROVEMENT OR STABILIZATION, AND 2) NO CONCURRENT USE WITH ANOTHER FABRY DISEASE THERAPY.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

ALECTINIB

Products Affected

• ALECENSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ALPELISIB-PIQRAY

Products Affected

 PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1), 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

AMBRISENTAN

Products Affected

• ambrisentan

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS.
Age Restrictions	
Prescriber Restrictions	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	PAH: INITIAL: DOES NOT HAVE IDIOPATHIC PULMONARY FIBROSIS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

AMIVANTAMAB-VMJW

Products Affected

• RYBREVANT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ANAKINRA

Products Affected

• KINERET

PA Criteria	Criteria Details
Exclusion Criteria	CORONAVIRUS DISEASE 2019 (COVID-19) IN HOSPITALIZED ADULTS.
Criteria	HOSPITALIZED ADULTS.
Required Medical Information	
Age Restrictions	
Prescriber	INITIAL: RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY
Restrictions	OR IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage	RA: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. ALL
Duration	OTHERS: 12 MONTHS.
Other Criteria	RA: INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, RINVOQ, ENBREL, XELJANZ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

APALUTAMIDE

Products Affected

• ERLEADA ORAL TABLET 240 MG, 60 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: NON-METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (NMCRPC): HIGH RISK PROSTATE CANCER (I.E., RAPIDLY INCREASING PROSTATE SPECIFIC ANTIGEN [PSA] LEVELS). NMCRPC, METASTATIC CASTRATION-SENSITIVE PROSTATE CANCER (MCSPC): 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. RENEWAL: NMCRPC, MCSPC: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GNRH ANALOG.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

APOMORPHINE

Products Affected

• apomorphine

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PARKINSONS DISEASE (PD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	PD: INITIAL: PHYSICIAN HAS OPTIMIZED DRUG THERAPY FOR PD. RENEWAL: IMPROVEMENT WITH MOTOR FLUCTUATIONS DURING OFF EPISODES WHILE ON THERAPY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

APOMORPHINE - SL

Products Affected

 KYNMOBI SUBLINGUAL FILM 10 MG, 10-15-20-25-30 MG, 15 MG, 20 MG, 25 MG, 30 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	PARKINSONS DISEASE (PD): INITIAL: 18 YEARS OF AGE OR OLDER.
Prescriber Restrictions	PD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	PD: INITIAL: PHYSICIAN HAS OPTIMIZED DRUG THERAPY FOR PARKINSONS DISEASE. RENEWAL: IMPROVEMENT WITH MOTOR FLUCTUATIONS DURING OFF EPISODES WITH THE USE OF THERAPY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

APREMILAST

Products Affected

- OTEZLA
- OTEZLA STARTER

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: MILD PLAQUE PSORIASIS (PSO): ONE OF THE FOLLOWING: 1) PSORIASIS COVERING 2 PERCENT OF BODY SURFACE AREA (BSA), 2) STATIC PHYSICIAN GLOBAL ASSESSMENT (SPGA) SCORE OF 2, OR 3) PSORIASIS AREA AND SEVERITY INDEX (PASI) SCORE OF 2 TO 9. MODERATE TO SEVERE PSO: 1) PSORIASIS COVERING 3 PERCENT OR MORE OF BSA, OR 2) PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, OR GENITAL AREA.
Age Restrictions	
Prescriber Restrictions	INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. BEHCETS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, TREMFYA, XELJANZ, RINVOQ, SKYRIZI. MILD PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL SYSTEMIC AGENT (E.G., METHOTREXATE, ACITRETIN, CYCLOSPORINE) AND ONE CONVENTIONAL TOPICAL AGENT (E.G., PUVA, UVB, TOPICAL CORTICOSTEROIDS). MODERATE TO SEVERE PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, SKYRIZI, TREMFYA. BEHCETS DISEASE: 1) HAS ORAL ULCERS OR A HISTORY OF RECURRENT ORAL ULCERS BASED ON CLINICAL SYMPTOMS, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE OR MORE CONSERVATIVE TREATMENTS (E.G., COLCHICINE, TOPICAL CORTICOSTEROID, ORAL CORTICOSTEROID). RENEWAL: PSA, PSO, BEHCETS DISEASE: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ASCIMINIB

Products Affected

SCEMBLIX ORAL TABLET 20 MG, 40 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PHILADELPHIA CHROMOSOME-POSITIVE CHRONIC MYELOID LEUKEMIA (Ph+ CML): MUTATIONAL ANALYSIS PRIOR TO INITIATION AND SCEMBLIX IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ASFOTASE ALFA

Products Affected

• STRENSIQ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	HYPOPHOSPHATASIA (HPP): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST, GENETICIST, OR METABOLIC SPECIALIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: PERINATAL/INFANTILE-ONSET HPP: 1) 6
	MONTHS OF AGE OR YOUNGER AT ONSET OF HPP, AND 2)
	POSITIVE FOR A TISSUE NON-SPECIFIC ALKALINE
	PHOSPHATASE (TNSALP) (ALPL) GENE MUTATION AS
	CONFIRMED BY GENETIC TESTING OR TWO OF THE
	FOLLOWING: (A) SERUM ALKALINE PHOSPHATASE (ALP)
	LEVEL BELOW THAT OF NORMAL RANGE FOR PATIENT
	AGE, (B) ELEVATED SERUM PYRIDOXAL-5'-PHOSPHATE
	(PLP) LEVELS AND NO VITAMIN B6 SUPPLEMENTATION IN
	THE PREVIOUS WEEK, (C) URINE
	PHOSPHOETHANOLAMINE (PEA) LEVEL ABOVE THAT OF
	NORMAL RANGE FOR PATIENT AGE, (D) RADIOGRAPHIC
	EVIDENCE OF HPP, (E) AT LEAST TWO OF THE
	FOLLOWING: (I) RACHITIC CHEST DEFORMITY, (II)
	CRANIOSYNOSTOSIS, (III) DELAY IN SKELETAL GROWTH
	RESULTING IN DELAY OF MOTOR DEVELOPMENT, (IV)
	HISTORY OF VITAMIN B6 DEPENDENT SEIZURES, (V)
	NEPHROCALCINOSIS OR HISTORY OF ELEVATED SERUM
	CALCIUM, (VI) HISTORY OR PRESENCE OF NON-
	TRAUMATIC POSTNATAL FRACTURE AND DELAYED
	FRACTURE HEALING. JUVENILE-ONSET HPP: 1) 18 YEARS
	OF AGE OR YOUNGER AT ONSET OF HPP, AND 2) POSITIVE
	FOR A TNSALP ALPL GENE MUTATION AS CONFIRMED
	BY GENETIC TESTING OR TWO OF THE FOLLOWING: (A)
	SERUM ALP LEVEL BELOW THAT OF NORMAL RANGE
	FOR PATIENT AGE, (B) ELEVATED SERUM PLP LEVELS
	AND NO VITAMIN B6 SUPPLEMENTATION IN THE
	PREVIOUS WEEK, (C) URINE PEA LEVEL ABOVE THAT OF
	NORMAL RANGE FOR PATIENT AGE, (D) RADIOGRAPHIC
	EVIDENCE OF HPP, (E) AT LEAST TWO OF THE
	FOLLOWING: (I) RACHITIC DEFORMITIES, (II)
	PREMATURE LOSS OF PRIMARY TEETH PRIOR TO 5
	YEARS OF AGE, (III) DELAY IN SKELETAL GROWTH
	RESULTING IN DELAY OF MOTOR DEVELOPMENT, (IV)

PA Criteria	Criteria Details
	HISTORY OR PRESENCE OF NON-TRAUMATIC FRACTURES OR DELAYED FRACTURE HEALING. ALL INDICATIONS: 1) NOT CURRENTLY RECEIVING TREATMENT WITH A BISPHOSPHONATE, 2) CALCIUM OR PHOSPHATE LEVELS ARE NOT BELOW THE NORMAL RANGE, 3) NOT HAVE A TREATABLE FORM OF RICKETS. RENEWAL: ALL INDICATIONS: 1) IMPROVEMENT IN THE SKELETAL CHARACTERISTICS OF HPP, AND 2) NOT CURRENTLY RECEIVING TREATMENT WITH A BISPHOSPHONATE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ATEZOLIZUMAB

Products Affected

• TECENTRIQ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ATOGEPANT

Products Affected

• QULIPTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	MIGRAINE PREVENTION: INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREVENTIVE MIGRAINE TREATMENTS: DIVALPROEX SODIUM, TOPIRAMATE, PROPRANOLOL, TIMOLOL, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. RENEWAL: 1) REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

AVAPRITINIB

Products Affected

• AYVAKIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

AVATROMBOPAG

Products Affected

- DOPTELET (10 TAB PACK)
- DOPTELET (15 TAB PACK)
- DOPTELET (30 TAB PACK)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: CHRONIC LIVER DISEASE (CLD): PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST, GASTROENTEROLOGIST, HEPATOLOGIST, IMMUNOLOGIST, ENDOCRINOLOGIST, OR A SURGEON. CHRONIC IMMUNE THROMBOCYTOPENIA (ITP): PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR IMMUNOLOGIST.
Coverage Duration	CLD: 1 MONTH. CHRONIC ITP: INITIAL: 2 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: CLD: 1) PLANNED PROCEDURE 10 TO 13 DAYS AFTER INITIATION OF DOPTELET, AND 2) NOT RECEIVING OTHER THROMBOPOIETIN RECEPTOR AGONISTS (E.G., ROMIPLOSTIM, ELTROMBOPAG, ETC.). CHRONIC ITP: TRIAL OF OR CONTRAINDICATION TO CORTICOSTEROIDS OR IMMUNOGLOBULINS OR INSUFFICIENT RESPONSE TO SPLENECTOMY. RENEWAL: CHRONIC ITP: PATIENT HAD A CLINICAL RESPONSE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

AXITINIB

Products Affected

• INLYTA ORAL TABLET 1 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

AZACITIDINE

Products Affected

• ONUREG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

AZTREONAM INHALED

Products Affected

• CAYSTON

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	7 YEARS OF AGE OR OLDER
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BARICITINIB

Products Affected

• OLUMIANT

PA Criteria	Criteria Details
Exclusion	CORONAVIRUS DISEASE 2019 (COVID-19) IN
Criteria	HOSPITALIZED ADULTS.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. SEVERE ALOPECIA AREATA: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST.
Coverage Duration	RA, SEVERE ALOPECIA AREATA: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO ANY TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ. SEVERE ALOPECIA AREATA: 1) AT LEAST 50 PERCENT SCALP HAIR LOSS AS MEASURED BY THE SEVERITY OF ALOPECIA TOOL (SALT) FOR MORE THAN 6 MONTHS, AND 2) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGICS FOR SEVERE ALOPECIA AREATA OR OTHER JAK INHIBITORS FOR ANY INDICATION. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION. SEVERE ALOPECIA AREATA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGICS FOR SEVERE ALOPECIA AREATA OR OTHER JAK INHIBITORS FOR ANY INDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

BECAPLERMIN

Products Affected

• REGRANEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	DIABETIC NEUROPATHIC ULCERS: PRESCRIBED BY OR IN CONSULTATION WITH A VASCULAR SURGEON, PODIATRIST, ENDOCRINOLOGIST, PHYSICIAN PRACTICING IN A SPECIALTY WOUND CLINIC OR INFECTIOUS DISEASE SPECIALIST.
Coverage Duration	3 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BEDAQUILINE

Products Affected

• SIRTURO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	24 WEEKS
Other Criteria	PULMONARY MULTI-DRUG RESISTANT TUBERCULOSIS (MDR-TB): SIRTURO USED IN COMBINATION WITH AT LEAST 3 OTHER ANTIBIOTICS FOR THE TREATMENT OF PULMONARY MDR-TB.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BELIMUMAB

- BENLYSTA INTRAVENOUS
- BENLYSTA SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: SYSTEMIC LUPUS ERYTHEMATOSUS (SLE): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. LUPUS NEPHRITIS (LN): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR NEPHROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: SLE: CURRENTLY TAKING CORTICOSTEROIDS, ANTIMALARIALS, NSAIDS, OR IMMUNOSUPPRESSIVE AGENTS. RENEWAL: SLE: PATIENT HAD CLINICAL IMPROVEMENT. LN: IMPROVEMENT IN RENAL RESPONSE FROM BASELINE LABORATORY VALUES (I.E., EGFR OR PROTEINURIA) AND/OR CLINICAL PARAMETERS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BELUMOSUDIL

Products Affected

• REZUROCK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BELZUTIFAN

Products Affected

• WELIREG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BENDAMUSTINE

- bendamustine intravenous recon soln
- BENDAMUSTINE INTRAVENOUS
- SOLUTION
- BENDEKA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BENRALIZUMAB

- FASENRA
- FASENRA PEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	ASTHMA: INITIAL: BLOOD EOSINOPHIL LEVEL GREATER THAN OR EQUAL TO 150 CELLS/MCL WITHIN THE PAST 12 MONTHS.
Age Restrictions	
Prescriber Restrictions	ASTHMA: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE.
Coverage Duration	INITIAL: 4 MONTHS, RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	ASTHMA: INITIAL: 1) CONCURRENT THERAPY WITH A
	MEDIUM, HIGH-DOSE, OR MAXIMALLY TOLERATED
	DOSE OF AN INHALED CORTICOSTEROID (ICS) AND ONE
	OTHER MAINTENANCE MEDICATION, AND 2) ONE
	ASTHMA EXACERBATION REQUIRING SYSTEMIC
	CORTICOSTEROID BURST LASTING 3 OR MORE DAYS
	WITHIN THE PAST 12 MONTHS, OR AT LEAST ONE
	SERIOUS EXACERBATION REQUIRING HOSPITALIZATION
	OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR
	SYMPTOM CONTROL DESPITE CURRENT THERAPY AS
	EVIDENCED BY AT LEAST THREE OF THE FOLLOWING
	WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA
	SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT
	WAKING DUE TO ASTHMA, SABA RELIEVER FOR
	SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY
	LIMITATION DUE TO ASTHMA, AND 3) NO CONCURRENT
	USE WITH XOLAIR, DUPIXENT OR OTHER ANTI-IL5
	BIOLOGICS WHEN USED FOR ASTHMA. RENEWAL: 1) NO
	CONCURRENT USE WITH XOLAIR, DUPIXENT OR OTHER
	ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA, 2)
	CONTINUED USE OF ICS AND ONE OTHER MAINTENANCE
	MEDICATION, AND 3) CLINICAL RESPONSE AS
	EVIDENCED BY: (A) REDUCTION IN ASTHMA
	EXACERBATIONS FROM BASELINE, (B) DECREASED
	UTILIZATION OF RESCUE MEDICATIONS, (C) INCREASE
	IN PERCENT PREDICTED FEV1 FROM PRETREATMENT
	BASELINE, OR (D) REDUCTION IN SEVERITY OR
	FREQUENCY OF ASTHMA-RELATED SYMPTOMS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B	No
Prerequisite	

BETAINE

Products Affected

• betaine

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BEVACIZUMAB-ADCD

Products Affected

• VEGZELMA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BEVACIZUMAB-AWWB

Products Affected

• MVASI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BEVACIZUMAB-BVZR

Products Affected

• ZIRABEV

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BEXAROTENE

Products Affected

• bexarotene

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BINIMETINIB

Products Affected

• MEKTOVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BORTEZOMIB

- bortezomib injection VELCADE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BOSENTAN

- bosentan
- TRACLEER ORAL TABLET FOR SUSPENSION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS.
Age Restrictions	
Prescriber Restrictions	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	PAH: INITIAL: 1) DOES NOT HAVE ELEVATED LIVER ENZYMES (ALT, AST) MORE THAN 3 TIMES UPPER LIMIT OF NORMAL (ULN) OR INCREASE IN BILIRUBIN BY 2 OR MORE TIMES ULN, AND 2) NOT CONCURRENTLY TAKING CYCLOSPORINE A OR GLYBURIDE. RENEWAL: NOT CONCURRENTLY TAKING CYCLOSPORINE A OR GLYBURIDE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BOSUTINIB

- BOSULIF ORAL CAPSULE 100 MG, 50 MG
- BOSULIF ORAL TABLET 100 MG, 400 MG, 500 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PREVIOUSLY TREATED (Ph+ CML): MUTATIONAL ANALYSIS PRIOR TO INITIATION AND BOSULIF IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BRIGATINIB

- ALUNBRIG ORAL TABLET 180 MG, 30 MG, 90 MG
- ALUNBRIG ORAL TABLETS, DOSE PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

C1 ESTERASE INHIBITOR-CINRYZE

Products Affected

• CINRYZE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HEREDITARY ANGIOEDEMA (HAE): INITIAL: DIAGNOSIS CONFIRMED BY COMPLEMENT TESTING.
Age Restrictions	
Prescriber Restrictions	HAE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST, IMMUNOLOGIST, OR ALLERGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	HAE: INITIAL: NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS. RENEWAL: 1) IMPROVEMENT IN HAE ATTACKS (I.E., REDUCTIONS IN ATTACK FREQUENCY OR ATTACK SEVERITY), AND 2) NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

C1 ESTERASE INHIBITOR-HAEGARDA

Products Affected

• HAEGARDA SUBCUTANEOUS RECON SOLN 2,000 UNIT, 3,000 UNIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HEREDITARY ANGIOEDEMA (HAE): INITIAL: DIAGNOSIS CONFIRMED BY COMPLEMENT TESTING.
Age Restrictions	
Prescriber Restrictions	HAE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST, IMMUNOLOGIST, OR ALLERGIST.
Coverage Duration	INITIAL AND RENEWAL: 12 MONTHS.
Other Criteria	HAE: INITIAL: NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS. RENEWAL: 1) IMPROVEMENT COMPARED TO BASELINE IN HAE ATTACKS (I.E., REDUCTIONS IN ATTACK FREQUENCY OR ATTACK SEVERITY), AND 2) NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CABOZANTINIB CAPSULE

Products Affected

 COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X1), 140 MG/DAY(80 MG X1-20 MG X3), 60 MG/DAY (20 MG X 3/DAY)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CABOZANTINIB TABLET

Products Affected

• CABOMETYX ORAL TABLET 20 MG, 40 MG, 60 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CANAKINUMAB

Products Affected

• ILARIS (PF)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: CRYOPYRIN-ASSOCIATED PERIODIC SYNDROMES (CAPS): 1) ONE OF THE FOLLOWING: (A) GENETIC TEST FOR GAIN-OF-FUNCTION MUTATIONS IN THE NLRP3 GENE, OR (B) HAS INFLAMMATORY MARKERS (I.E., ELEVATED CRP, ESR, SERUM AMYLOID A PROTEIN (SAA) OR S100 PROTEINS), AND 2) TWO OF THE FOLLOWING: URTICARIAL-LIKE RASH (NEUTROPHILIC DERMATITIS), COLD-TRIGGERED EPISODES, SENSORINEURAL HEARING LOSS, MUSCULOSKELETAL SYMPTOMS, CHRONIC ASEPTIC MENINGITIS, SKELETAL ABNORMALITIES.
Age Restrictions	
Prescriber Restrictions	INITIAL: SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS (SJIA), ADULT-ONSET STILLS DISEASE (AOSD): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, OR IMMUNOLOGIST. GOUT: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. RENEWAL: GOUT: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	INITIAL: AOSD/SJIA: 6 MO, CAPS: LIFETIME, ALL OTHER DIAGNOSES: 12 MO. RENEWAL: AOSD/SJIA/GOUT: 12 MO

PA Criteria	Criteria Details
Other Criteria	INITIAL: CAPS: NO CONCURRENT USE WITH OTHER IL-1 INHIBITORS. AOSD, SJIA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE- MODIFYING ANTIRHEUMATIC DRUG). GOUT: NO CONCURRENT USE WITH OTHER IL-1 INHIBITORS. RENEWAL: AOSD, SJIA: CONTINUES TO BENEFIT FROM THE MEDICATION. GOUT: 1) NO CONCURRENT USE WITH OTHER IL-1 INHIBITORS, AND 2) IMPROVEMENT IN GOUT FLARES.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CANNABIDIOL

Products Affected

• EPIDIOLEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	DRAVET SYNDROME (DS), LENNOX-GASTAUT SYNDROME (LGS), TUBEROUS SCLEROSIS COMPLEX (TSC): PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL: 12 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: LENNOX-GASTAUT SYNDROME (LGS): TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING ANTIEPILEPTIC MEDICATIONS: RUFINAMIDE, FELBAMATE, CLOBAZAM, TOPIRAMATE, LAMOTRIGINE, CLONAZEPAM. RENEWAL: DS, LGS, TSC: CONFIRMATION OF DIAGNOSIS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CAPIVASERTIB

Products Affected

• TRUQAP

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CAPLACIZUMAB YHDP

Products Affected

• CABLIVI INJECTION KIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	ACQUIRED THROMBOTIC THROMBOCYTOPENIA PURPURA (ATTP): PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST
Coverage Duration	12 MONTHS
Other Criteria	ATTP: CABLIVI WAS PREVIOUSLY INITIATED AS PART OF THE FDA APPROVED TREATMENT REGIMEN IN COMBINATION WITH PLASMA EXCHANGE AND IMMUNOSUPPRESSIVE THERAPY WITHIN AN INPATIENT SETTING. THE PATIENT HAS NOT EXPERIENCED MORE THAN TWO RECURRENCES OF ATTP WHILE ON CABLIVI THERAPY (I.E., NEW DROP IN PLATELET COUNT REQUIRING REPEAT PLASMA EXCHANGE DURING 30 DAYS POST-PLASMA EXCHANGE THERAPY [PEX] AND UP TO 28 DAYS OF EXTENDED THERAPY).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CAPMATINIB

Products Affected

• TABRECTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CARGLUMIC ACID

Products Affected

• carglumic acid

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: ACUTE OR CHRONIC HYPERAMMONEMIA (HA) DUE TO N ACETYLGLUTAMATE SYNTHASE (NAGS) DEFICIENCY: NAGS GENE MUTATION IS CONFIRMED BY BIOCHEMICAL OR GENETIC TESTING. ACUTE HA DUE TO PROPIONIC ACIDEMIA (PA): 1) CONFIRMED BY ELEVATED METHYLCITRIC ACID AND NORMAL METHYLMALONIC ACID, OR 2) GENETIC TESTING CONFIRMS MUTATION IN THE PCCA OR PCCB GENE. ACUTE HA DUE TO METHYLMALONIC ACIDEMIA (MMA): 1) CONFIRMED BY ELEVATED METHYLMALONIC ACID, METHYLCITRIC ACID, OR 2) GENETIC TESTING CONFIRMS MUTATION IN THE METHYLMALONIC ACID, METHYLCITRIC ACID, OR 2) GENETIC TESTING CONFIRMS MUTATION IN THE MMUT, MMA, MMAB OR MMADHC GENES.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	ACUTE HA DUE TO NAGS/PA/MMA: 7 DAYS. CHRONIC HA DUE TO NAGS: INITIAL: 6 MOS, RENEWAL: 12 MOS.
Other Criteria	RENEWAL: CHRONIC HA DUE TO NAGS: PATIENT HAS SHOWN CLINICAL IMPROVEMENT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CERITINIB

Products Affected

• ZYKADIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CETUXIMAB

Products Affected

• ERBITUX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CLADRIBINE

- MAVENCLAD (10 TABLET PACK)
- MAVENCLAD (4 TABLET PACK)
- MAVENCLAD (5 TABLET PACK)
- MAVENCLAD (6 TABLET PACK)
- MAVENCLAD (7 TABLET PACK)
- MAVENCLAD (8 TABLET PACK)
- MAVENCLAD (9 TABLET PACK)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL/RENEWAL: 48 WEEKS.
Other Criteria	RELAPSING FORM OF MULTIPLE SCLEROSIS (MS): INITIAL: HAS NOT RECEIVED A TOTAL OF TWO YEARS OF MAVENCLAD TREATMENT (I.E., TWO YEARLY TREATMENT COURSES OF TWO CYCLES IN EACH). RENEWAL: 1) HAS DEMONSTRATED CLINICAL BENEFIT COMPARED TO PRE-TREATMENT BASELINE, 2) DOES NOT HAVE LYMPHOPENIA, AND 3) HAS NOT RECEIVED A TOTAL OF TWO YEARS OF MAVENCLAD TREATMENT (I.E., TWO YEARLY TREATMENT COURSES OF TWO CYCLES IN EACH).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CLOBAZAM-SYMPAZAN

Products Affected

• SYMPAZAN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	LENNOX-GASTAUT SYNDROME (LGS): THERAPY IS PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	LGS: 1) UNABLE TO TAKE TABLETS OR SUSPENSIONS, AND 2) TRIAL OF OR CONTRAINDICATION TO A FORMULARY VERSION OF CLOBAZAM.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

COBIMETINIB

Products Affected

• COTELLIC

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

COLCHICINE

Products Affected

• colchicine oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	PROPHYLAXIS OF GOUT FLARES: 16 YEARS AND OLDER
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PROPHYLAXIS OF GOUT FLARES: TRIAL OF OR CONTRAINDICATION TO COLCHICINE CAPSULES (MITIGARE) IF AGE 18 YEARS OR OLDER.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CORTICOTROPIN

Products Affected

- ACTHAR
- CORTROPHIN GEL

PA Criteria	Criteria Details
Exclusion Criteria	INITIAL: NOT APPROVED FOR DIAGNOSTIC PURPOSES.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS AND MULTIPLE SCLEROSIS (MS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, ALLERGIST/IMMUNOLOGIST, OPHTHALMOLOGIST, PULMONOLOGIST OR NEPHROLOGIST.
Coverage Duration	INFANTILE SPASMS AND MS: 28 DAYS. OTHER FDA APPROVED INDICATIONS: INITIAL AND RENEWAL: 28 DAYS
Other Criteria	INITIAL: ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS: TRIAL OF OR CONTRAINDICATION TO INTRAVENOUS (IV) CORTICOSTEROIDS. ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS AND MS: TRIAL OF OR CONTRAINDICATION TO A STANDARD OF CARE THERAPY. RENEWAL: ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS AND MS: 1) DEMONSTRATED CLINICAL BENEFIT WHILE ON THERAPY AS INDICATED BY SYMPTOM RESOLUTION AND/OR NORMALIZATION OF LABORATORY TESTS, AND 2) CONTINUES TO POSSESS CONTRAINDICATION TO IV CORTICOSTEROIDS. PART B BEFORE PART D STEP THERAPY, APPLIES ONLY TO BENEFICIARIES IN AN MA- PD PLAN.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

CRIZANLIZUMAB-TMCA

Products Affected

• ADAKVEO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	SICKLE CELL DISEASE (SCD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST
Coverage Duration	INITIAL: 12 MONTHS. RENEWAL: LIFETIME
Other Criteria	SCD: INITIAL: AGES 18 YEARS OR OLDER: 1) AT LEAST 2 SICKLE CELL CRISES IN THE PAST YEAR, 2) SICKLE-CELL ASSOCIATED SYMPTOMS WHICH ARE INTERFERING WITH ACTIVITIES OF DAILY LIVING, OR 3) HISTORY OF OR HAS RECURRENT ACUTE CHEST SYNDROME. AGES 16 TO 17 YEARS: APPROVED WITHOUT ADDITIONAL CRITERIA. RENEWAL: MAINTAINED OR EXPERIENCED A REDUCTION IN ACUTE COMPLICATIONS OF SCD. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CRIZOTINIB

Products Affected

• XALKORI ORAL CAPSULE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CRIZOTINIB PELLETS

Products Affected

• XALKORI ORAL PELLET 150 MG, 20 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	NON-SMALL CELL LUNG CANCER (NSCLC), ANAPLASTIC LARGE CELL LYMPHOMA (ALCL), INFLAMMATORY MYOFIBROBLASTIC TUMOR (IMT): UNABLE TO SWALLOW CAPSULES.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CYSTEAMINE HYDROCHLORIDE

Products Affected

- CYSTADROPS
- CYSTARAN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DABRAFENIB

Products Affected

• TAFINLAR ORAL CAPSULE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DABRAFENIB SUSPENSION

Products Affected

 TAFINLAR ORAL TABLET FOR SUSPENSION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	UNABLE TO SWALLOW TAFINILAR CAPSULES.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DACOMITINIB

Products Affected

• VIZIMPRO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC): NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DALFAMPRIDINE

Products Affected

• dalfampridine

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	MULTIPLE SCLEROSIS (MS): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL: 3 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	MS: INITIAL: HAS SYMPTOMS OF A WALKING DISABILITY SUCH AS MILD TO MODERATE BILATERAL LOWER EXTREMITY WEAKNESS OR UNILATERAL WEAKNESS PLUS LOWER EXTREMITY OR TRUNCAL ATAXIA. RENEWAL: IMPROVEMENT IN WALKING ABILITY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DARATUMUMAB

Products Affected

• DARZALEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DARATUMUMAB-HYALURONIDASE-FIHJ

Products Affected

• DARZALEX FASPRO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DAROLUTAMIDE

Products Affected

• NUBEQA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS
Other Criteria	INITIAL: NON-METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (NMCRPC): HIGH RISK PROSTATE CANCER (I.E., RAPIDLY INCREASING PROSTATE SPECIFIC ANTIGEN [PSA] LEVELS). NMCRPC, METASTATIC HORMONE-SENSITIVE PROSTATE CANCER (MHSPC): 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. RENEWAL: NMCRPC, MHSPC: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GNRH ANALOG.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DASATINIB

Products Affected

• SPRYCEL ORAL TABLET 100 MG, 140 MG, 20 MG, 50 MG, 70 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PREVIOUSLY-TREATED PHILADELPHIA CHROMOSOME- POSITIVE CHRONIC MYELOID LEUKEMIA (Ph+ CML): MUTATIONAL ANALYSIS PRIOR TO INITIATION AND SPRYCEL IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR- ABL1 MUTATION PROFILE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DECITABINE/CEDAZURIDINE

Products Affected

• INQOVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DEFERASIROX

Products Affected

• deferasirox

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: CHRONIC IRON OVERLOAD DUE TO BLOOD TRANSFUSIONS: SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 1000 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS). NON-TRANSFUSION DEPENDENT THALASSEMIA (NTDT): 1) SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 300 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS) AND 2) LIVER IRON CONCENTRATION (LIC) OF 5 MG FE/G DRY WEIGHT OR GREATER. RENEWAL: CHRONIC IRON OVERLOAD DUE TO BLOOD TRANSFUSIONS: SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 500 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS). NTDT: 1) SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 300 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS) OR 2) LIC OF 3 MG FE/G DRY WEIGHT OR GREATER.
Age Restrictions	
Prescriber Restrictions	INITIAL: CHRONIC IRON OVERLOAD: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR HEMATOLOGIST/ONCOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL FOR ALL INDICATIONS: FORMULARY VERSION OF DEFERASIROX SPRINKLE: TRIAL OF OR CONTRAINDICATION TO GENERIC DEFERASIROX TABLET OR TABLET FOR ORAL SUSPENSION.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

DEFERIPRONE

Products Affected

- *deferiprone*FERRIPROX (2 TIMES A DAY)
 FERRIPROX ORAL SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	TRANSFUSIONAL IRON OVERLOAD: RENEWAL: SERUM FERRITIN LEVELS CONSISTENTLY ABOVE 500 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS).
Age Restrictions	
Prescriber Restrictions	TRANSFUSIONAL IRON OVERLOAD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR HEMATOLOGIST/ONCOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: TRANSFUSIONAL IRON OVERLOAD DUE TO THALASSEMIA SYNDROMES: 1) TRIAL OF, CONTRAINDICATION, INTOLERABLE TOXICITIES, OR CLINICALLY SIGNIFICANT ADVERSE EFFECTS TO A FORMULARY VERSION OF DEFERASIROX OR DEFEROXAMINE, OR 2) CURRENT CHELATION THERAPY (I.E., FORMULARY VERSION OF DEFERASIROX OR DEFEROXAMINE) IS INADEQUATE. TRANSFUSIONAL IRON OVERLOAD DUE TO SICKLE CELL DISEASE OR OTHER ANEMIAS: TRIAL OF OR CONTRAINDICATION TO A FORMULARY VERSION OF DEFERASIROX OR DEFEROXAMINE.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

DEFEROXAMINE

Products Affected

• deferoxamine

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	CHRONIC IRON OVERLOAD: INITIAL: SERUM FERRITIN LEVELS CONSISTENTLY ABOVE 1000 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS). RENEWAL: SERUM FERRITIN LEVELS CONSISTENTLY ABOVE 500 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS).
Age Restrictions	CHRONIC IRON OVERLOAD: INITIAL: 3 YEARS OR OLDER
Prescriber Restrictions	CHRONIC IRON OVERLOAD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR HEMATOLOGIST/ONCOLOGIST
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. THIS DRUG MAY BE EITHER BUNDLED WITH AND COVERED UNDER END STAGE RENAL DISEASE DIALYSIS RELATED SERVICES OR COVERED UNDER MEDICARE D DEPENDING UPON THE CIRCUMSTANCES.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DENOSUMAB-XGEVA

Products Affected

• XGEVA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DEUTETRABENAZINE

Products Affected

• AUSTEDO ORAL TABLET 12 MG, 6 MG, 9 MG 24 MG, 6 MG

• AUSTEDO XR ORAL TABLET EXTENDED RELEASE 24 HR 12 MG, • AUSTEDO XR TITRATION KT(WK1-4)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	HUNTINGTON DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR MOVEMENT DISORDER SPECIALIST. TARDIVE DYSKINESIA: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR MOVEMENT DISORDER SPECIALIST.
Coverage Duration	12 MONTHS
Other Criteria	TARDIVE DYSKINESIA: HISTORY OF USING AGENTS THAT CAUSE TARDIVE DYSKINESIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DICLOFENAC TOPICAL GEL

Products Affected

• diclofenac sodium topical gel 3 %

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DICLOFENAC TOPICAL SOLUTION

Products Affected

• *diclofenac sodium topical solution in metered-dose pump*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	OSTEOARTHRITIS OF THE KNEE: TRIAL OF OR CONTRAINDICATION TO A FORMULARY VERSION OF DICLOFENAC SODIUM 1% TOPICAL GEL AND A FORMULARY VERSION OF DICLOFENAC SODIUM 1.5% TOPICAL DROPS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DIMETHYL FUMARATE

Products Affected

 dimethyl fumarate oral capsule, delayed release(drlec) 120 mg, 120 mg (14)-240 mg (46), 240 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DIROXIMEL FUMARATE

Products Affected

• VUMERITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DOSTARLIMAB-GXLY

Products Affected

• JEMPERLI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DRONABINOL CAPSULE

Products Affected

• dronabinol

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	NAUSEA AND VOMITING ASSOCIATED WITH CANCER CHEMOTHERAPY: TRIAL OF OR CONTRAINDICATION TO ONE ANTIEMETIC THERAPY. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D FOR THE INDICATION OF NAUSEA AND VOMITING ASSOCIATED WITH CANCER CHEMOTHERAPY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DROXIDOPA

Products Affected

• droxidopa

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	NEUROGENIC ORTHOSTATIC HYPOTENSION (NOH): INITIAL: 1) BASELINE BLOOD PRESSURE READINGS WHILE THE PATIENT IS SITTING AND ALSO WITHIN 3 MINUTES OF STANDING FROM A SUPINE POSITION. 2) A DECREASE OF AT LEAST 20 MMHG IN SYSTOLIC BLOOD PRESSURE OR 10 MMHG DIASTOLIC BLOOD PRESSURE WITHIN THREE MINUTES AFTER STANDING FROM A SITTING POSITION.
Age Restrictions	
Prescriber Restrictions	NOH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR CARDIOLOGIST.
Coverage Duration	INITIAL: 3 MONTHS RENEWAL: 12 MONTHS
Other Criteria	NOH: RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DUPILUMAB

Products Affected

- DUPIXENT PEN
- DUPIXENT SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: EOSINOPHILIC ASTHMA: BLOOD EOSINOPHIL LEVEL OF 150 TO 1500 CELLS/MCL WITHIN THE PAST 12 MONTHS. EOSINOPHILIC ESOPHAGITIS (EOE): DIAGNOSIS CONFIRMED BY ESOPHAGOGASTRODUODENOSCOPY (EGD) WITH BIOPSY.
Age Restrictions	
Prescriber Restrictions	INITIAL: AD, PN: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST, ALLERGIST OR IMMUNOLOGIST. ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE. CRSWNP: PRESCRIBED BY OR IN CONSULTATION WITH AN OTOLARYNGOLOGIST, ALLERGIST OR IMMUNOLOGIST. EOE: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST, ALLERGIST, OR IMMUNOLOGIST.
Coverage Duration	INITIAL: AD, CRSWNP, EOE, PN: 6 MOS, ASTHMA: 4 MOS. RENEWAL: ALL INDICATIONS: 12 MOS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: AD: 1) AD COVERING AT LEAST 10 PERCENT OF BODY SURFACE AREA OR AD AFFECTING THE FACE,
	HEAD, NECK, HANDS, FEET, GROIN, OR INTERTRIGINOUS
	AREAS, 2) INTRACTABLE PRURITUS OR
	CRACKING/OOZING/BLEEDING OF AFFECTED SKIN, 3)
	TRIAL OF OR CONTRAINDICATION TO ONE TOPICAL
	(CORTICOSTEROID, CALCINEURIN INHIBITOR, PDE4
	INHIBITOR, OR JAK INHIBITOR), AND 4) NO
	CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGICS
	OR JAK INHIBITORS FOR AD. ASTHMA: 1) CONCURRENT
	THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY-
	TOLERATED DOSE OF AN INHALED CORTICOSTEROID
	(ICS) AND ONE OTHER MAINTENANCE MEDICATION, 2)
	ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC
	CORTICOSTEROID BURST LASTING 3 OR MORE DAYS
	WITHIN THE PAST 12 MONTHS, OR ONE SERIOUS
	EXACERBATION REQUIRING HOSPITALIZATION OR ER
	VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM
	CONTROL DESPITE CURRENT THERAPY AS EVIDENCED
	BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE
	THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO
	ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN
	TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO
	ASTHMA, AND 3) NO CONCURRENT USE WITH XOLAIR OR
	OTHER ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA.
	CRSWNP: 1) EVIDENCE OF NASAL POLYPS BY DIRECT
	EXAMINATION, ENDOSCOPY OR SINUS CT SCAN, 2)
	INADEQUATELY CONTROLLED DISEASE AS
	DETERMINED BY USE OF SYSTEMIC STEROIDS IN THE
	PAST 2 YEARS OR ENDOSCOPIC SINUS SURGERY, AND 3) A
	56 DAY TRIAL OF ONE TOPICAL NASAL
	CORTICOSTEROID. PN: 1) CHRONIC PRURITIS (ITCH MORE
	THAN 6 WEEKS), MULTIPLE PRURIGINOUS LESIONS, AND

PA Criteria	Criteria Details
	HISTORY OR SIGN OF A PROLONGED SCRATCHING BEHAVIOR, 2) TRIAL OF OR CONTRAINDICATION TO ONE TOPICAL (CORTICOSTEROID OR CALCIPOTRIOL). RENEWAL: AD: 1) IMPROVEMENT WHILE ON THERAPY, AND 2) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGICS OR JAK INHIBITORS FOR AD. CRSWNP, EOE: IMPROVEMENT WHILE ON THERAPY. ASTHMA: 1) NO CONCURRENT USE WITH XOLAIR, OR OTHER ANTI-IL5 BIOLOGICS FOR ASTHMA, 2) CONTINUED USE OF ICS AND ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) INCREASE IN PERCENT PREDICTED FEV1 FROM PRETREATMENT BASELINE, OR (D) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS. PN: IMPROVEMENT OR REDUCTION OF PRURITIS OR PRURIGINOUS LESIONS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DUVELISIB

Products Affected

• COPIKTRA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

EDARAVONE

Products Affected

• RADICAVA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	AMYOTROPHIC LATERAL SCLEROSIS (ALS): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR ALS SPECIALIST AT AN ALS SPECIALTY CENTER OR CARE CLINIC.
Coverage Duration	ALS: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	ALS: INITIAL: 1) DURATION OF DISEASE (FROM ONSET OF SYMPTOMS) IS LESS THAN OR EQUAL TO 2 YEARS, 2) NORMAL RESPIRATORY FUNCTION, 3) HAS MILD TO MODERATE ALS WITH A SCORE OF 2 OR HIGHER IN ALL OF THE FOLLOWING 12 ITEMS OF THE AMYOTROPHIC LATERAL SCLEROSIS FUNCTIONAL RATING SCALE REVISED (ALSFRS-R): SPEECH, SALIVATION, SWALLOWING, HANDWRITING, CUTTING FOOD, DRESSING AND HYGIENE, TURNING IN BED, WALKING, CLIMBING STAIRS, DYSPNEA, ORTHOPNEA, RESPIRATORY INSUFFICIENCY, AND 4) TRIAL OF RILUZOLE TABLET OR CURRENTLY TAKING RILUZOLE TABLET. RENEWAL: 1) DOES NOT REQUIRE INVASIVE VENTILATION, AND 2) HAS IMPROVED BASELINE FUNCTIONAL ABILITY OR HAS MAINTAINED A SCORE OF 2 OR HIGHER IN ALL 12 ITEMS OF THE ALSFRS-R.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

EFLAPEGRASTIM-XNST

Products Affected

• ROLVEDON

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NON MYELOID MALIGNANCY: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	NON MYELOID MALIGNANCY: TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: NYVEPRIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

EFLORNITHINE

Products Affected

• IWILFIN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	24 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ELACESTRANT

Products Affected

 ORSERDU ORAL TABLET 345 MG, 86 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ELAGOLIX

Products Affected

• ORILISSA ORAL TABLET 150 MG, 200 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS.
Age Restrictions	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: 18 YEARS OF AGE OR OLDER.
Prescriber Restrictions	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS
Other Criteria	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, AND 2) TRIAL OF OR CONTRAINDICATION TO AN NSAID AND A PROGESTIN-CONTAINING PREPARATION. RENEWAL: 1) IMPROVEMENT IN PAIN ASSOCIATED WITH ENDOMETRIOSIS WHILE ON THERAPY, AND 2) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

ELAPEGADEMASE-LVLR

Products Affected

• REVCOVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	ADENOSINE DEAMINASE SEVERE COMBINED IMMUNE DEFICIENCY (ADA-SCID): INITIAL: ADA-SCID AS MANIFESTED BY: 1) CONFIRMATORY GENETIC TEST, OR 2) SUGGESTIVE LABORATORY FINDINGS (E.G., ELEVATED DEOXYADENOSINE NUCLEOTIDE [DAXP] LEVELS, LYMPHOPENIA) AND HALLMARK SIGNS/SYMPTOMS (E.G., RECURRENT INFECTIONS, FAILURE TO THRIVE, PERSISTENT DIARRHEA).
Age Restrictions	
Prescriber Restrictions	ADA-SCID: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH IMMUNOLOGIST, HEMATOLOGIST/ONCOLOGIST, OR PHYSICIAN SPECIALIZING IN INHERITED METABOLIC DISORDERS.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	ADA-SCID: RENEWAL: 1) IMPROVEMENT OR MAINTENANCE OF IMMUNE FUNCTION FROM BASELINE, AND 2) HAS NOT RECEIVED SUCCESSFUL HCT OR GENE THERAPY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ELEXACAFTOR-TEZACAFTOR-IVACAFTOR

Products Affected

- TRIKAFTA ORAL GRANULES IN PACKET, SEQUENTIAL
- TRIKAFTA ORAL TABLETS, SEQUENTIAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	CYSTIC FIBROSIS (CF): INITIAL: CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CYSTIC FIBROSIS.
Age Restrictions	
Prescriber Restrictions	CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CYSTIC FIBROSIS EXPERT.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: LIFETIME.
Other Criteria	CF: RENEWAL: 1) MAINTAINED, IMPROVED, OR DEMONSTRATED LESS THAN EXPECTED DECLINE IN FEV1 OR BODY MASS INDEX (BMI), OR 2) REDUCTION IN NUMBER OF PULMONARY EXACERBATIONS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ELIGLUSTAT

Products Affected

• CERDELGA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ELOSULFASE ALFA

Products Affected

• VIMIZIM

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ELRANATAMAB-BCMM

Products Affected

- ELREXFIO 44 MG/1.1 ML VIAL OUTER, SUV, P/F
- ELREXFIO SUBCUTANEOUS SOLUTION 40 MG/ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	RELAPSED OR REFRACTORY MULTIPLE MYELOMA: RENEWAL: 1) HAS RECEIVED AT LEAST 24 WEEKS OF TREATMENT WITH ELREXFIO, AND 2) HAS RESPONDED TO TREATMENT (PARTIAL RESPONSE OR BETTER), AND HAS MAINTAINED THIS RESPONSE FOR AT LEAST 2 MONTHS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ELTROMBOPAG

Products Affected

- PROMACTA ORAL POWDER IN PACKET 12.5 MG, 25 MG
- PROMACTA ORAL TABLET 12.5 MG, 25 MG, 50 MG, 75 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: ITP: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR IMMUNOLOGIST.
Coverage Duration	ITP: INITIAL: 2 MO, RENEWAL: 12 MO. HEPATITIS C, SEVERE APLASTIC ANEMIA: 12 MO.
Other Criteria	INITIAL: PERSISTENT OR CHRONIC IMMUNE THROMBOCYTOPENIA PURPURA (ITP): TRIAL OF OR CONTRAINDICATION TO CORTICOSTEROIDS, IMMUNOGLOBULINS, OR AN INSUFFICIENT RESPONSE TO SPLENECTOMY. ALL INDICATIONS: APPROVAL FOR PROMACTA ORAL SUSPENSION PACKETS REQUIRES A TRIAL OF PROMACTA TABLETS OR PATIENT IS UNABLE TO TAKE TABLET FORMULATION. RENEWAL: ITP: PATIENT HAS SHOWN A CLINICAL RESPONSE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

EMAPALUMAB-LZSG

Products Affected

• GAMIFANT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS (HLH): INITIAL: 1) A GENETIC TEST IDENTIFYING HLH- ASSOCIATED GENE MUTATION (E.G., PRF1, UNC13D), OR 2) HAS AT LEAST FIVE OF THE FOLLOWING EIGHT DIAGNOSTIC CRITERIA FOR HLH: (A) FEVER, (B) SPLENOMEGALY, (C) CYTOPENIAS (AFFECTING AT LEAST 2 OF 3 CELL LINEAGES), (D) HYPERTRIGLYCERIDEMIA OR HYPOFIBRINOGENEMIA, (E) HEMOPHAGOCYTOSIS IN BONE MARROW OR SPLEEN OR LYMPH NODES AND NO EVIDENCE OF MALIGNANCY, (F) LOW OR ABSENT NATURAL KILLER-CELL ACTIVITY, (G) FERRITIN LEVEL OF 500 MCG/L OR GREATER, (H) SOLUBLE CD25 LEVEL OF 2,400 U/ML OR GREATER.
Age Restrictions	
Prescriber Restrictions	HLH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN IMMUNOLOGIST, HEMATOLOGIST, OR ONCOLOGIST.
Coverage Duration	INITIAL AND RENEWAL: 8 WEEKS.

PA Criteria	Criteria Details
Other Criteria	HLH: INITIAL: 1) CONCURRENT THERAPY WITH DEXAMETHASONE, AND 2) ONE OF THE FOLLOWING: (A) HAS REFRACTORY, RECURRENT, OR PROGRESSIVE DISEASE, OR (B) HAD A TRIAL OF OR INTOLERANCE TO CONVENTIONAL HLH THERAPY (I.E., CHEMOTHERAPY, STEROIDS, IMMUNOTHERAPY). RENEWAL: 1) HAS NOT RECEIVED SUCCESSFUL HEMATOPOIETIC STEM CELL TRANSPLANTATION, AND 2) DEMONSTRATED IMPROVED IMMUNE SYSTEM RESPONSE FROM BASELINE (E.G., RESOLUTION OF FEVER, DECREASED SPLENOMEGALY, IMPROVEMENT IN CNS SYMPTOMS, IMPROVED CBC, INCREASED FIBRINOGEN LEVELS, REDUCED D-DIMER, REDUCED FERRITIN, REDUCED SOLUBLE CD25 LEVELS.)
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ENASIDENIB

Products Affected

• IDHIFA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ENCORAFENIB

Products Affected

• BRAFTOVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ENTRECTINIB

Products Affected

ROZLYTREK ORAL CAPSULE 100
 MG, 200 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ENTRECTINIB PELLETS

Products Affected

• ROZLYTREK ORAL PELLETS IN PACKET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC), SOLID TUMORS: 1) TRIAL OF OR CONTRAINDICATION TO ROZLYTREK CAPSULES MADE INTO AN ORAL SUSPENSION, AND 2) DIFFICULTY OR UNABLE TO SWALLOW CAPSULES.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ENZALUTAMIDE

Products Affected

- XTANDI ORAL CAPSULE
- XTANDI ORAL TABLET 40 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	INITIAL: NON-METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (NMCRPC): HIGH RISK PROSTATE CANCER (I.E. RAPIDLY INCREASING PROSTATE SPECIFIC ANTIGEN [PSA] LEVELS). NON-METASTATIC CASTRATION- SENSITIVE PROSTATE CANCER (NMCSPC): HIGH RISK FOR METASTASIS (I.E. PSA DOUBLED OVER 9 MONTHS OR LESS). METASTATIC CRPC (MCRPC), NMCRPC, METASTATIC CSPC (MCSPC), NMCSPC : 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. RENEWAL: MCRPC, NMCRPC, MCSPC: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GNRH ANALOG.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

EPCORITAMAB-BYSP

Products Affected

• EPKINLY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

EPOETIN ALFA-EPBX

Products Affected

 RETACRIT INJECTION SOLUTION 10,000 UNIT/ML, 2,000 UNIT/ML, 20,000 UNIT/2 ML, 20,000 UNIT/ML, 3,000 UNIT/ML, 4,000 UNIT/ML, 40,000 UNIT/ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: CHRONIC KIDNEY DISEASE (CKD), ANEMIA RELATED TO ZIDOVUDINE, OR CANCER CHEMOTHERAPY: HEMOGLOBIN LEVEL OF LESS THAN 10G/DL. ELECTIVE, NON-CARDIAC, NON-VASCULAR SURGERY: HEMOGLOBIN LEVEL LESS THAN 13G/DL. RENEWAL: 1) CKD IN ADULTS NOT ON DIALYSIS: (A) HEMOGLOBIN LEVEL IS LESS THAN 10G/DL, OR (B) HEMOGLOBIN LEVEL HAS REACHED 10G/DL AND THE DOSE IS BEING REDUCED/INTERRUPTED TO DECREASE THE NEED FOR BLOOD TRANSFUSIONS. CKD IN PEDIATRIC PATIENTS: (A) HEMOGLOBIN LEVEL IS LESS THAN 10G/DL, OR (B) HEMOGLOBIN LEVEL HAS APPROACHED OR EXCEEDS 12G/DL AND THE DOSE IS BEING REDUCED/INTERRUPTED TO DECREASE THE NEED FOR BLOOD TRANSFUSIONS. ANEMIA RELATED TO ZIDOVUDINE: HEMOGLOBIN LEVEL BETWEEN 10G/DL AND 12G/DL. CANCER CHEMOTHERAPY: 1) HEMOGLOBIN LEVEL OF LESS THAN 10 G/DL, OR 2) THE HEMOGLOBIN LEVEL DOES NOT EXCEED A LEVEL NEEDED TO AVOID RBC TRANSFUSION.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	ANEMIA FROM CHEMO/CKD WITHOUT DIALYSIS/ZIDOVUDINE: INITIAL/RENEWAL: 12 MONTHS. SURGERY: 1 MONTH.

PA Criteria	Criteria Details
Other Criteria	RENEWAL: CKD: NOT RECEIVING DIALYSIS TREATMENT. THIS DRUG MAY BE EITHER BUNDLED WITH AND COVERED UNDER END STAGE RENAL DISEASE DIALYSIS RELATED SERVICES OR COVERED UNDER MEDICARE D DEPENDING UPON THE CIRCUMSTANCES.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

EPOPROSTENOL IV

Products Affected

• epoprostenol

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS.
Age Restrictions	
Prescriber Restrictions	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
Coverage Duration	INITIAL AND RENEWAL: 12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ERDAFITINIB

Products Affected

 BALVERSA ORAL TABLET 3 MG, 4 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ERLOTINIB

Products Affected

• erlotinib oral tablet 100 mg, 150 mg, 25 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC NSCLC WITH EGFR MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ESKETAMINE

Products Affected

• SPRAVATO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: TREATMENT-RESISTANT DEPRESSION (TRD), MAJOR DEPRESSIVE DISORDER (MDD): PRESCRIBED BY OR IN CONSULTATION WITH A PSYCHIATRIST.
Coverage Duration	INITIAL: TRD: 3 MONTHS. MDD: 4 WEEKS. RENEWAL: TRD, MDD: 12 MONTHS.
Other Criteria	INITIAL: TRD: 1) NON-PSYCHOTIC, UNIPOLAR DEPRESSION, 2) NO ACTIVE SUBSTANCE ABUSE, AND 3) ADEQUATE TRIAL (AT LEAST 4 WEEKS) OF AT LEAST TWO ANTIDEPRESSANT AGENTS FROM DIFFERENT CLASSES THAT ARE INDICATED FOR DEPRESSION. MDD: 1) NON- PSYCHOTIC, UNIPOLAR DEPRESSION, AND 2) NO ACTIVE SUBSTANCE ABUSE. RENEWAL: TRD, MDD: DEMONSTRATED CLINICAL BENEFIT (IMPROVEMENT IN DEPRESSION) COMPARED TO BASELINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ETANERCEPT

Products Affected

- ENBREL
- ENBREL MINI
- ENBREL SURECLICK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE OR GENITAL AREA.
Age Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS): 18 YEARS OR OLDER. PSORIATIC ARTHRITIS (PSA): 2 YEARS OR OLDER.
Prescriber Restrictions	INITIAL: RA, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA), AS: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSA: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE- MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PJIA, PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD. PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. RENEWAL: RA, PJIA, PSA, AS, PSO: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ETEPLIRSEN

Products Affected

• EXONDYS-51

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	DUCHENNE MUSCULAR DYSTROPHY (DMD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	DMD: INITIAL: PATIENT IS AMBULATORY AND CURRENTLY RECEIVING TREATMENT WITH OR HAS A CONTRAINDICATION TO CORTICOSTEROIDS. RENEWAL: MAINTAINED OR DEMONSTRATED A LESS THAN EXPECTED DECLINE IN AMBULATORY ABILITY IN MUSCLE FUNCTION ASSESSMENTS OR OTHER MUSCLE FUNCTION (I.E., PULMONARY OR CARDIAC FUNCTION). THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

EVEROLIMUS-AFINITOR

Products Affected

• everolimus (antineoplastic) oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

EVEROLIMUS-AFINITOR DISPERZ

Products Affected

• *everolimus (antineoplastic) oral tablet for suspension*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FECAL MICROBIOTA CAPSULE

Products Affected

• VOWST

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	30 DAYS
Other Criteria	CLOSTRIDIOIDES DIFFICILE INFECTION (CDI): 1) HAS NOT PREVIOUSLY RECEIVED VOWST: COMPLETION OF ANTIBIOTIC TREATMENT FOR RECURRENT CDI (AT LEAST 3 CDI EPISODES), OR 2) PREVIOUSLY RECEIVED VOWST: (A) TREATMENT FAILURE (DEFINED AS THE PRESENCE OF CDI DIARRHEA WITHIN 8 WEEKS OF FIRST DOSE OF VOWST AND A POSITIVE STOOL TEST FOR C. DIFFICILE), AND (B) HAS NOT RECEIVED MORE THAN ONE TREATMENT COURSE OF VOWST WHICH WAS AT LEAST 12 DAYS AND NOT MORE THAN 8 WEEKS PRIOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FEDRATINIB

Products Affected

• INREBIC

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	MYELOFIBROSIS: INITIAL: TRIAL OF OR CONTRAINDICATION TO JAKAFI (RUXOLITINIB). RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FENFLURAMINE

Products Affected

• FINTEPLA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: DRAVET SYNDROME, LENNOX-GASTAUT SYNDROME (LGS): PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	DRAVET SYNDROME: INITIAL/RENEWAL: 12 MONTHS. LGS: 12 MONTHS.
Other Criteria	INITIAL: LGS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING ANTIEPILEPTIC MEDICATIONS: RUFINAMIDE, FELBAMATE, CLOBAZAM, TOPIRAMATE, LAMOTRIGINE, CLONAZEPAM. RENEWAL: DRAVET SYNDROME: PATIENT HAS SHOWN CONTINUED CLINICAL BENEFIT (E.G. REDUCTION OF SEIZURES, REDUCED LENGTH OF SEIZURES, SEIZURE CONTROL MAINTAINED).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FENTANYL CITRATE

Products Affected

• fentanyl citrate buccal lozenge on a handle

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	CANCER RELATED PAIN: 1) CURRENTLY ON A MAINTENANCE DOSE OF CONTROLLED-RELEASE OPIOID PAIN MEDICATION, AND 2) TRIAL OF OR CONTRAINDICATION TO AT LEAST ONE IMMEDIATE- RELEASE ORAL OPIOID PAIN AGENT OR PATIENT HAS DIFFICULTY SWALLOWING TABLETS/CAPSULES. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FILGRASTIM-AAFI

Products Affected

• NIVESTYM

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FILGRASTIM-AYOW

Products Affected

• RELEUKO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT NIVESTYM, WHERE INDICATIONS ALIGN.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FILGRASTIM-SNDZ

Products Affected

• ZARXIO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: NIVESTYM.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FINASTERIDE/TADALAFIL

Products Affected

• ENTADFI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	BENIGN PROSTATIC HYPERPLASIA (BPH): 18 YEARS OF AGE OR OLDER
Prescriber Restrictions	
Coverage Duration	26 WEEKS
Other Criteria	BPH: 1) TRIAL OF OR CONTRAINDICATION TO ONE 5- ALPHA-REDUCTASE INHIBITOR, AND 2) TRIAL OF OR CONTRAINDICATION TO TADALAFIL 2.5MG OR TADALAFIL 5MG.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FINERENONE

Products Affected

• KERENDIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FINGOLIMOD

- *fingolimod*GILENYA ORAL CAPSULE 0.25 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FINGOLIMOD LAURYL SULFATE

Products Affected

TASCENSO ODT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	MULTIPLE SCLEROSIS (MS): (1) UNABLE TO SWALLOW FINGOLIMOD CAPSULES, AND (2) TRIAL OF OR CONTRAINDICATION TO FINGOLIMOD CAPSULES.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FOSTAMATINIB

Products Affected

• TAVALISSE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	CHRONIC IMMUNE THROMBOCYTOPENIA (ITP): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR IMMUNOLOGIST.
Coverage Duration	INITIAL: 3 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	ITP: RENEWAL: PATIENT HAS SHOWN A CLINICAL RESPONSE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FREMANEZUMAB-VFRM

- AJOVY AUTOINJECTOR
- AJOVY SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	MIGRAINE PREVENTION: INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREVENTIVE MIGRAINE TREATMENTS: DIVALPROEX SODIUM, TOPIRAMATE, PROPRANOLOL, TIMOLOL, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. RENEWAL: 1) REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FRUQUINTINIB

Products Affected

 FRUZAQLA ORAL CAPSULE 1 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FUTIBATINIB

Products Affected

• LYTGOBI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INTRAHEPATIC CHOLANGIOCARCINOMA (ICCA): COMPLETE A COMPREHENSIVE OPHTHALMOLOGICAL EXAMINATION, INCLUDING OPTICAL COHERENCE TOMOGRAPHY (OCT), PRIOR TO THE INITIATION OF THERAPY AND AT THE RECOMMENDED SCHEDULED INTERVALS.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

GALCANEZUMAB-GNLM

Products Affected

• EMGALITY PEN

 EMGALITY SYRINGE SUBCUTANEOUS SYRINGE 120 MG/ML, 300 MG/3 ML (100 MG/ML X

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: MIGRAINE PREVENTION: 6 MOS. EPISODIC CLUSTER HEADACHE: 3 MOS. RENEWAL (ALL): 12 MOS.
Other Criteria	INITIAL: MIGRAINE PREVENTION: 1) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREVENTIVE MIGRAINE TREATMENTS: DIVALPROEX SODIUM, TOPIRAMATE, PROPRANOLOL, TIMOLOL. RENEWAL: MIGRAINE PREVENTION: 1) REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. EPISODIC CLUSTER HEADACHE: IMPROVEMENT IN EPISODIC CLUSTER HEADACHE FREQUENCY AS COMPARED TO BASELINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

3)

GANAXOLONE

Products Affected

• ZTALMY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

GEFITINIB

Products Affected

• gefitinib

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC NSCLC WITH EGFR MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

GILTERITINIB

Products Affected

• XOSPATA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

GIVOSIRAN

Products Affected

• GIVLAARI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	ACUTE HEPATIC PORPHYRIA (AHP): INITIAL: GENETIC CONFIRMATION OF MUTATION OR ELEVATED URINARY OR PLASMA PBG (PORPHOBILINOGEN) OR ALA (AMINOLEVULINIC ACID).
Age Restrictions	
Prescriber Restrictions	ACUTE HEPATIC PORPHYRIA (AHP): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A GENETICIST, HEPATOLOGIST, HEMATOLOGIST, GASTROENTEROLOGIST, NEUROLOGIST, DERMATOLOGIST, OR A HEALTHCARE PROVIDER EXPERIENCED IN MANAGING AHP.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS
Other Criteria	AHP: INITIAL: EXPERIENCED TWO OR MORE AHP ATTACKS IN THE PAST 12 MONTHS. RENEWAL: 1) ACHIEVED OR MAINTAINED CLINICAL BENEFIT COMPARED TO BASELINE, AND 2) HAS NOT RECEIVED A LIVER TRANSPLANT. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

GLASDEGIB

Products Affected

 DAURISMO ORAL TABLET 100 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

GLATIRAMER

Products Affected

- COPAXONE SUBCUTANEOUS SYRINGE 20 MG/ML, 40 MG/ML
- glatiramer subcutaneous syringe 20 mg/ml, 40 mg/ml

• glatopa subcutaneous syringe 20 mg/ml, 40 mg/ml

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

GLECAPREVIR/PIBRENTASVIR

Products Affected

• MAVYRET ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HCV RNA LEVEL WITHIN PAST 6 MONTHS
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
Other Criteria	1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, 2) TRIAL OF A PREFERRED FORMULARY ALTERNATIVE INCLUDING HARVONI OR EPCLUSA WHEN THESE AGENTS ARE CONSIDERED ACCEPTABLE FOR TREATMENT OF THE SPECIFIC GENOTYPE PER AASLD/IDSA GUIDANCE, 3) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING MEDICATIONS: CARBAMAZEPINE, RIFAMPIN, ETHINYL ESTRADIOL-CONTAINING MEDICATION, ATAZANAVIR, DARUNAVIR, LOPINAVIR, RITONAVIR, EFAVIRENZ, ATORVASTATIN, LOVASTATIN, SIMVASTATIN, ROSUVASTATIN AT DOSES GREATER THAN 10MG, CYCLOSPORINE AT DOSES GREATER THAN 100MG PER DAY, EPCLUSA, HARVONI, VOSEVI, OR ZEPATIER, 4) PATIENT MUST NOT HAVE PRIOR FAILURE OF A DAA REGIMEN WITH NS5A INHIBITOR AND HCV PROTEASE INHIBITOR, AND 5) DOES NOT HAVE MODERATE OR SEVERE HEPATIC IMPAIRMENT (CHILD PUGH B OR C).
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

GLP1-DULAGLUTIDE

Products Affected

• TRULICITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

GLP1-SEMAGLUTIDE

Products Affected

 OZEMPIC SUBCUTANEOUS PEN INJECTOR 0.25 MG OR 0.5 MG (2 MG/3 ML), 0.25 MG OR 0.5 MG(2 MG/1.5 ML), 1 MG/DOSE (4 MG/3 ML), 2 MG/DOSE (8 MG/3 ML)

• RYBELSUS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

GLP1-TIRZEPATIDE

Products Affected

• MOUNJARO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

GLYCEROL PHENYLBUTYRATE

Products Affected

• RAVICTI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	UREA CYCLE DISORDER (UCD): INITIAL: DIAGNOSIS IS CONFIRMED BY ENZYMATIC, BIOCHEMICAL OR GENETIC TESTING
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS
Other Criteria	UCD: INITIAL: TRIAL OF OR CONTRAINDICATION TO SODIUM PHENYLBUTYRATE. RENEWAL: PATIENT HAS CLINICAL BENEFIT FROM BASELINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

GOSERELIN

Products Affected

• ZOLADEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	ENDOMETRIOSIS: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS.
Age Restrictions	
Prescriber Restrictions	ENDOMETRIOSIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST.
Coverage Duration	STAGE B2-C PROSTATIC CARCINOMA: 4 MOS. ENDOMETRIOSIS: 6 MOS PER LIFETIME. ALL OTHERS: 12 MONTHS.
Other Criteria	ENDOMETRIOSIS: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, 2) TRIAL OF OR CONTRAINDICATION TO NSAID AND PROGESTIN- CONTAINING PREPARATION, AND 3) HAS NOT RECEIVED A TOTAL OF 6 MONTHS OF TREATMENT PER LIFETIME. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

GUSELKUMAB

Products Affected

• TREMFYA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE OR GENITAL AREA.
Age Restrictions	
Prescriber Restrictions	INITIAL: PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG). PSO: TRIAL OF OR CONTRAINDICATION ONE CONVENTIONAL THERAPY SUCH AS A PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. RENEWAL: PSO, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH CONCENTRATION ORAL OPIOID SOLUTIONS

- morphine concentrate oral solution
- oxycodone oral concentrate

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	OPIOID TOLERANT: 12 MONTHS. HOSPICE, PALLIATIVE CARE OR END OF LIFE CARE: LIFETIME.
Other Criteria	1) OPIOID TOLERANT (I.E. PREVIOUS USE OF 60 MG ORAL MORPHINE PER DAY, 25 MCG TRANSDERMAL FENTANYL PER HOUR, 30 MG ORAL OXYCODONE PER DAY, 8 MG ORAL HYDROMORPHONE PER DAY, 25 MG ORAL OXYMORPHONE PER DAY, 60 MG ORAL HYDROCODONE PER DAY, OR AN EQUIANALGESIC DOSE OF ANOTHER OPIOID) AND HAS TROUBLE SWALLOWING OPIOID TABLETS, CAPSULES, OR LARGE VOLUMES OF LIQUID, OR 2) ENROLLED IN HOSPICE OR PALLIATIVE CARE OR END OF LIFE CARE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY - CARBINOXAMINE

- carbinoxamine maleate oral liquid
- carbinoxamine maleate oral tablet 4 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY - CYPROHEPTADINE

Products Affected

• cyproheptadine

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY -**BUTALBITAL-CONTAINING AGENTS**

- ascomp with codeine
- butalbital-acetaminop-caf-cod
- butalbital-acetaminophen oral tablet 50-325 codeine-butalbital-asa-caff mg
- *butalbital-acetaminophen-caff oral capsule* 50-325-40 mg
- butalbital-acetaminophen-caff oral tablet
- butalbital-aspirin-caffeine
- tencon
- *zebutal*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS ARE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY -CLEMASTINE

Products Affected

• clemastine oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY -CONJUGATED ESTROGEN

Products Affected

• PREMARIN ORAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	VULVAR/VAGINAL ATROPHY, OSTEOPOROSIS AND VASOMOTOR SYMPTOMS OF MENOPAUSE: PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HYPOESTROGENISM TREATMENT, PALLIATIVE TREATMENT, AND HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY -DIPYRIDAMOLE

Products Affected

• dipyridamole oral

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY -DISOPYRAMIDE

Products Affected

• disopyramide phosphate oral capsule

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY -ESTRADIOL

Products Affected

• dotti

• lyllana

- estradiol oral
- estradiol transdermal patch semiweekly
- estradiol transdermal patch weekly

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	VULVAR/VAGINAL ATROPHY, OSTEOPOROSIS AND VASOMOTOR SYMPTOMS OF MENOPAUSE: PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HYPOESTROGENISM TREATMENT, PALLIATIVE TREATMENT, AND HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY -ESTRADIOL-NORETHINDRONE

- amabelz
- estradiol-norethindrone acet oral tablet 0.5-0.1 mg
- mimvey

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	VULVAR/VAGINAL ATROPHY, OSTEOPOROSIS, AND VASOMOTOR SYMPTOMS OF MENOPAUSE: PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HYPOESTROGENISM TREATMENT AND HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY -ESTROGEN-BAZEDOXIFENE

Products Affected

• DUAVEE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY -ESTROGEN-MEDROXYPROGESTERONE

- PREMPHASE
- PREMPRO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY -GLYBURIDE FORMULATIONS

- glyburide
- glyburide micronized
- glyburide-metformin

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	TRIAL OF GLIMEPIRIDE, GLIPIZIDE, OR PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS ARE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY -KETOROLAC

- ketorolac injection solution 15 mg/ml, 30 mg/ml, 30 mg/ml (1 ml)
- ketorolac injection syringe 15 mg/ml, 30 mg/ml
- ketorolac intramuscular solution
- *ketorolac intramuscular syringe*
- *ketorolac oral*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	30 DAYS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY -NORETHINDRONE-ESTRADIOL

- fyavolv
- jinteli
- norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY -PHENOBARBITAL

Products Affected

• phenobarbital

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	FOR TREATMENT OF EPILEPSY/SEIZURES IN PATIENTS WHO ARE NEWLY PRESCRIBED PHENOBARBITAL: PATIENT HAS NOT RESPONDED TO AT LEAST ONE ANTICONVULSANT OR PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY -PROMETHAZINE

- promethazine injection solution
- promethazine oral
- promethazine rectal
- promethegan

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRURITUS/URTICARIA/SEASONAL/PERENNIAL ALLERGY: 1) TRIAL OF OR CONTRAINDICATION TO A NON- SEDATING ANTIHISTAMINE SUCH AS LEVOCETIRIZINE, OR 2) PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. NAUSEA AND VOMITING: PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH-RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS REQUIRE PHYSICIAN ATTESTATION THAT REQUESTED MEDICATION IS USED TO TREAT A DIAGNOSIS UNRELATED TO THE TERMINAL ILLNESS OR RELATED CONDITION, AND ARE APPROVED WITHOUT TRIAL OF FORMULARY ALTERNATIVES NOR REQUIRING PRESCRIBER ACKNOWLEDGEMENT.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY -SCOPOLAMINE

Products Affected

• scopolamine base

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS REQUIRE PHYSICIAN ATTESTATION THAT REQUESTED MEDICATION IS USED TO TREAT A DIAGNOSIS UNRELATED TO THE TERMINAL ILLNESS OR RELATED CONDITION, AND ARE APPROVED WITHOUT REQUIRING PRESCRIBER ACKNOWLEDGEMENT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY -SKELETAL MUSCLE RELAXANTS

- chlorzoxazone oral tablet 250 mg, 500 mg, 750 mg
- cyclobenzaprine oral tablet 10 mg, 5 mg
- methocarbamol oral tablet 500 mg, 750 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED A HIGH RISK MEDICATION FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY-DIPHENHYDRAMINE ELIXIR

Products Affected

• diphenhydramine hcl oral elixir

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRURITUS OR URTICARIA: TRIAL OF OR CONTRAINDICATION TO A NON-SEDATING ANTIHISTAMINE SUCH AS LEVOCETIRIZINE. INSOMNIA: TRIAL OF SILENOR AND BELSOMRA. MOTION SICKNESS AND ANTIPARKINSONISM: PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS AND ANAPHYLACTIC REACTIONS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY-DIPHENOXYLATE-ATROPINE

Products Affected

• diphenoxylate-atropine

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY-INDOMETHACIN

- indomethacin oral capsule 25 mg, 50 mg
- indomethacin oral capsule, extended release

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY-MEGESTROL

- megestrol oral suspension 400 mg/10 ml (40 mg/ml)
- megestrol oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY-PAROXETINE

- paroxetine hcl oral suspension
- paroxetine hcl oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HISTRELIN-SUPPRELIN LA

Products Affected

• SUPPRELIN LA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	CENTRAL PRECOCIOUS PUBERTY (CPP): INITIAL: FEMALES: ELEVATED LEVELS OF FOLLICLE- STIMULATING HORMONE (FSH) GREATER THAN 4.0 MIU/ML AND LUTEINIZING HORMONE (LH) LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS. MALES: ELEVATED LEVELS OF FSH GREATER THAN 5.0 MIU/ML AND LH LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS.
Age Restrictions	
Prescriber Restrictions	CPP: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	CPP: INITIAL: FEMALES: 1) YOUNGER THAN 8 YEARS OF AGE AT ONSET OF CPP, AND 2) AT TANNER STAGE 2 OR ABOVE FOR BREAST DEVELOPMENT AND PUBIC HAIR GROWTH. MALES: 1) YOUNGER THAN 9 YEARS OF AGE AT ONSET OF CPP, AND 2) AT TANNER STAGE 2 OR ABOVE FOR GENITAL DEVELOPMENT AND PUBIC HAIR GROWTH. RENEWAL: 1) TANNER STAGING AT INITIAL DIAGNOSIS HAS STABILIZED OR REGRESSED DURING THREE SEPARATE MEDICAL VISITS IN THE PREVIOUS YEAR, AND 2) HAS NOT REACHED ACTUAL AGE WHICH CORRESPONDS TO CURRENT PUBERTAL AGE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

IBRUTINIB

Products Affected

• IMBRUVICA ORAL CAPSULE 140 MG, 70 MG 280 MG, 420 MG

- IMBRUVICA ORAL SUSPENSION
- IMBRUVICA ORAL TABLET 140 MG,

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: BRUKINSA, WHERE INDICATIONS ALIGN.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

IBUPROFEN-FAMOTIDINE

Products Affected

• ibuprofen-famotidine

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	TRIAL OF ONE OF THE FOLLOWING GENERIC, FEDERAL LEGEND HISTAMINE H2-RECEPTOR ANTAGONISTS: FAMOTIDINE, CIMETIDINE, OR NIZATIDINE, AND TRIAL OF GENERIC, FEDERAL LEGEND IBUPROFEN.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ICATIBANT

- icatibant
- sajazir

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HEREDITARY ANGIOEDEMA (HAE): DIAGNOSIS CONFIRMED BY COMPLEMENT TESTING.
Age Restrictions	
Prescriber Restrictions	HAE: PRESCRIBED BY OR IN CONSULTATION WITH AN ALLERGIST, IMMUNOLOGIST, OR HEMATOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	HAE: NO CONCURRENT USE WITH OTHER MEDICATIONS FOR TREATMENT OF ACUTE HAE ATTACKS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

IDELALISIB

Products Affected

• ZYDELIG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

IMATINIB

Products Affected

• imatinib oral tablet 100 mg, 400 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	ADJUVANT GASTROINTESTINAL STROMAL TUMOR TREATMENT: 36 MONTHS. ALL OTHER DIAGNOSES: 12 MONTHS.
Other Criteria	PHILADELPHIA CHROMOSOME POSITIVE CHRONIC MYELOID LEUKEMIA: PATIENT HAS NOT RECEIVED A PREVIOUS TREATMENT WITH ANOTHER TYROSINE KINASE INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

INFIGRATINIB

Products Affected

• TRUSELTIQ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	CHOLANGIOCARCINOMA: COMPREHENSIVE OPHTHALMOLOGICAL EXAMINATION, INCLUDING OPTICAL COHERENCE TOMOGRAPHY (OCT), WILL BE COMPLETED PRIOR TO INITIATION OF THERAPY AND AT THE RECOMMENDED SCHEDULED INTERVALS.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

INFLIXIMAB

Products Affected

• infliximab

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, OR FACE.
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: RA: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, XELJANZ, TREMFYA, RINVOQ, SKYRIZI. PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, SKYRIZI, TREMFYA. AS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, COSENTYX, ENBREL, XELJANZ, RINVOQ. CD: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS; HUMIRA, COSENTYX, ENBREL, XELJANZ, RINVOQ. CD: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, RINVOQ, SKYRIZI. UC: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, RINVOQ, SKYRIZI. UC: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, XELJANZ, RINVOQ. RENEWAL: RA, AS, PSO, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

INFLIXIMAB-ABDA

Products Affected

• RENFLEXIS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, OR FACE.
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: RA: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, XELJANZ, TREMFYA, RINVOQ, SKYRIZI. PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, SKYRIZI, TREMFYA. AS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, COSENTYX, ENBREL, XELJANZ, RINVOQ. CD: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, COSENTYX, ENBREL, XELJANZ, RINVOQ. CD: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, SKYRIZI, RINVOQ. UC: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, SKYRIZI, RINVOQ. UC: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, XELJANZ, RINVOQ. RENEWAL: RA, AS, PSO, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

INFLIXIMAB-AXXQ

Products Affected

• AVSOLA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, OR FACE.
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: RA: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, XELJANZ, TREMFYA, RINVOQ, SKYRIZI. PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, SKYRIZI, TREMFYA. AS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, COSENTYX, ENBREL, XELJANZ, RINVOQ. CD: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, COSENTYX, ENBREL, XELJANZ, RINVOQ. CD: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, SKYRIZI, RINVOQ. UC: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, SKYRIZI, RINVOQ. UC: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, XELJANZ, RINVOQ. RENEWAL: RA, AS, PSO, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

INFLIXIMAB-DYYB

Products Affected

• INFLECTRA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, OR FACE.
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: RA: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, XELJANZ, TREMFYA, RINVOQ, SKYRIZI. PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, SKYRIZI, TREMFYA. AS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, COSENTYX, ENBREL, XELJANZ, RINVOQ. CD: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS; HUMIRA, COSENTYX, ENBREL, XELJANZ, RINVOQ. CD: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, SKYRIZI, RINVOQ. UC: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, SKYRIZI, RINVOQ. UC: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, XELJANZ, RINVOQ. RENEWAL: RA, AS, PSO, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

INTERFERON FOR MS-AVONEX

- AVONEX INTRAMUSCULAR PEN INJECTOR KIT
- AVONEX INTRAMUSCULAR SYRINGE KIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

INTERFERON FOR MS-BETASERON

Products Affected

• BETASERON SUBCUTANEOUS KIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

INTERFERON FOR MS-PLEGRIDY

Products Affected

 PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML, 63 MCG/0.5 ML- 94 MCG/0.5 ML
 PLEGRIDY SUBCUTANEOUS SYRINGE 125 MCG/0.5 ML, 63 MCG/0.5 ML- 94 MCG/0.5 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

INTERFERON GAMMA-1B

Products Affected

• ACTIMMUNE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: CHRONIC GRANULOMATOUS DISEASE (CGD): PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST, INFECTIOUS DISEASE SPECIALIST, OR IMMUNOLOGIST. SEVERE MALIGNANT OSTEOPETROSIS (SMO): PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	RENEWAL: CGD, SMO: 1) DEMONSTRATED CLINICAL BENEFIT COMPARED TO BASELINE, AND 2) HAS NOT RECEIVED HEMATOPOIETIC CELL TRANSPLANTATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

IPILIMUMAB

Products Affected

• YERVOY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: UNRESECT/MET MEL: 4MO, RCC/CRC/HCC: 3MO, ALL OTHERS: 12MO. INITIAL/RENEWAL: CUTAN MEL: 6MO
Other Criteria	RENEWAL: ADJUVANT CUTANEOUS MELANOMA: NO EVIDENCE OF DISEASE RECURRENCE (DEFINED AS THE APPEARANCE OF ONE OR MORE NEW MELANOMA LESIONS: LOCAL, REGIONAL OR DISTANT METASTASIS). THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ITRACONAZOLE SOLUTION

Products Affected

• itraconazole oral solution

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	ESOPHAGEAL CANDIDIASIS AND OROPHARYNGEAL CANDIDIASIS: TRIAL OF OR CONTRAINDICATION TO FLUCONAZOLE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

IVACAFTOR

Products Affected

• KALYDECO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	CYSTIC FIBROSIS (CF): INITIAL: CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CYSTIC FIBROSIS
Age Restrictions	
Prescriber Restrictions	CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CYSTIC FIBROSIS EXPERT
Coverage Duration	INITIAL: 12 MONTHS. RENEWAL: LIFETIME
Other Criteria	CF: INITIAL: NOT HOMOZYGOUS FOR F508DEL MUTATION IN CFTR GENE. RENEWAL: 1) MAINTAINED, IMPROVED, OR DEMONSTRATED LESS THAN EXPECTED DECLINE IN FEV1 OR BODY MASS INDEX (BMI), OR 2) REDUCTION IN NUMBER OF PULMONARY EXACERBATIONS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

IVOSIDENIB

Products Affected

• TIBSOVO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

IXAZOMIB

Products Affected

• NINLARO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

IXEKIZUMAB

- TALTZ AUTOINJECTOR
- TALTZ SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, OR GENITAL AREA. NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN (CRP) LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILLITIS ON MAGNETIC RESONANCE IMAGING (MRI).
Age Restrictions	
Prescriber Restrictions	INITIAL: PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. ANKYLOSING SPONDYLITIS (AS), NR-AXSPA: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, COSENTYX, STELARA, ENBREL, SKYRIZI, TREMFYA. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, COSENTYX, STELARA, ENBREL, XELJANZ, TREMFYA, RINVOQ, SKYRIZI. AS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA, COSENTYX, XELJANZ, RINVOQ. NR-AXSPA: 1) TRIAL OF OR CONTRAINDICATION TO BOTH OF THE PREFERRED AGENTS: COSENTYX, RINVOQ, OR 2) TRIAL OF COSENTYX AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. RENEWAL: PSO, PSA, AS, NR-AXSPA: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LANADELUMAB-FLYO

Products Affected

 TAKHZYRO SUBCUTANEOUS SOLUTION (150 MG/ML)

• TAKHZYRO SUBCUTANEOUS SYRINGE 150 MG/ML, 300 MG/2 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HEREDITARY ANGIOEDEMA (HAE): INITIAL: DIAGNOSIS CONFIRMED BY COMPLEMENT TESTING.
Age Restrictions	
Prescriber Restrictions	HAE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ALLERGIST, IMMUNOLOGIST, OR HEMATOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	HAE: INITIAL: NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS. RENEWAL: 1) IMPROVEMENT COMPARED TO BASELINE IN HAE ATTACKS (I.E., REDUCTIONS IN ATTACK FREQUENCY OR ATTACK SEVERITY), AND 2) NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LANREOTIDE

Products Affected

• lanreotide

 SOMATULINE DEPOT SUBCUTANEOUS SYRINGE 120 MG/0.5 ML, 60 MG/0.2 ML, 90 MG/0.3

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	ACROMEGALY: INITIAL: THERAPY IS PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	ACROMEGALY: INITIAL: 3 MOS, RENEWAL: 12 MOS.GEP- NETS, CARCINOID SYNDROME: 12 MOS.
Other Criteria	ACROMEGALY: INITIAL: TRIAL OF OR CONTRAINDICATION TO ONE GENERIC OCTREOTIDE INJECTION. RENEWAL: 1) REDUCTION, NORMALIZATION, OR MAINTENANCE OF IGF-1 LEVELS BASED ON AGE AND GENDER, AND 2) IMPROVEMENT OR SUSTAINED REMISSION OF CLINICAL SYMPTOMS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ML

LAPATINIB

Products Affected

• lapatinib

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LAROTRECTINIB

- VITRAKVI ORAL CAPSULE 100 MG, 25 MG
- VITRAKVI ORAL SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	APPROVAL FOR VITRAKVI ORAL SOLUTION: TRIAL OF VITRAKVI CAPSULES OR PATIENT IS UNABLE TO TAKE CAPSULE FORMULATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LEDIPASVIR-SOFOSBUVIR

- HARVONI ORAL PELLETS IN PACKET 33.75-150 MG, 45-200 MG
- HARVONI ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HCV RNA LEVEL WITHIN PAST 6 MONTHS.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
Other Criteria	1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, AND 2) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING: CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, ROSUVASTATIN, TIPRANAVIR/RITONAVIR, SOFOSBUVIR (AS A SINGLE AGENT), EPCLUSA, ZEPATIER, MAVYRET, OR VOSEVI. REQUESTS FOR HARVONI 45MG-200MG PELLETS: PATIENT IS UNABLE TO SWALLOW TABLETS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LENALIDOMIDE

Products Affected

• lenalidomide

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LENVATINIB

Products Affected

• LENVIMA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LETERMOVIR

- PREVYMIS INTRAVENOUS SOLUTION 240 MG/12 ML, 480 MG/24 ML
- PREVYMIS ORAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	HSCT: NOT AT RISK FOR LATE CMV: 4 MOS, AT RISK FOR LATE CMV: 7 MOS. KIDNEY TRANSPLANT: 7 MOS.
Other Criteria	HEMATOPOIETIC STEM CELL TRANSPLANT (HSCT): 1) THERAPY WILL BE INITIATED BETWEEN DAY 0 AND DAY 28 POST TRANSPLANT, AND 2) WILL NOT RECEIVE THE MEDICATION BEYOND 100 DAYS POST TRANSPLANT IF NOT AT RISK FOR LATE CYTOMEGALOVIRUS (CMV) INFECTION AND DISEASE, OR BEYOND 200 DAYS POST TRANSPLANT IF AT RISK FOR LATE CMV INFECTION AND DISEASE. KIDNEY TRANSPLANT: 1) THERAPY WILL BE INITIATED BETWEEN DAY 0 AND DAY 7 POST TRANSPLANT, AND 2) WILL NOT RECEIVE THE MEDICATION BEYOND 200 DAYS POST TRANSPLANT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LEUPROLIDE

Products Affected

• leuprolide subcutaneous kit

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	PROSTATE CANCER: 12 MONTHS.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LEUPROLIDE DEPOT

Products Affected

• *leuprolide (3 month)*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LEUPROLIDE-ELIGARD

- ELIGARD
- ELIGARD (3 MONTH)
- ELIGARD (4 MONTH)
- ELIGARD (6 MONTH)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LEUPROLIDE-LUPRON DEPOT

- LUPRON DEPOT
- LUPRON DEPOT (3 MONTH)
- LUPRON DEPOT (4 MONTH)
- LUPRON DEPOT (6 MONTH)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: ENDOMETRIOSIS: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS.
Age Restrictions	
Prescriber Restrictions	INITIAL: ENDOMETRIOSIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST.
Coverage Duration	PROSTATE CA: 12 MOS. UTERINE FIBROIDS: 3 MOS. ENDOMETRIOSIS: INITIAL/RENEWAL: 6 MOS.
Other Criteria	INITIAL: ENDOMETRIOSIS: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, 2) TRIAL OF OR CONTRAINDICATION TO NSAID AND PROGESTIN- CONTAINING PREPARATION, AND 3) HAS NOT RECEIVED A TOTAL OF 12 MONTHS OF TREATMENT PER LIFETIME. RENEWAL: ENDOMETRIOSIS: 1) IMPROVEMENT OF PAIN RELATED TO ENDOMETRIOSIS WHILE ON THERAPY, 2) RECEIVING CONCOMITANT ADD-BACK THERAPY (I.E., COMBINATION ESTROGEN-PROGESTIN OR PROGESTIN- ONLY CONTRACEPTIVE PREPARATION), 3) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, AND 4) HAS NOT RECEIVED A TOTAL OF 12 MONTHS OF TREATMENT PER LIFETIME. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

LEUPROLIDE-LUPRON DEPOT-PED

- LUPRON DEPOT-PED
- LUPRON DEPOT-PED (3 MONTH)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	CENTRAL PRECOCIOUS PUBERTY (CPP): INITIAL: FEMALES: ELEVATED LEVELS OF FOLLICLE- STIMULATING HORMONE (FSH) GREATER THAN 4.0 MIU/ML AND LUTEINIZING HORMONE (LH) LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS. MALES: ELEVATED LEVELS OF FSH GREATER THAN 5.0 MIU/ML AND LH LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS.
Age Restrictions	
Prescriber Restrictions	CPP: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	CPP: INITIAL: FEMALES: 1) YOUNGER THAN 8 YEARS OF AGE AT ONSET OF CPP, AND 2) AT TANNER STAGE 2 OR ABOVE FOR BREAST DEVELOPMENT AND PUBIC HAIR GROWTH. MALES: 1) YOUNGER THAN 9 YEARS OF AGE AT ONSET OF CPP, AND 2) AT TANNER STAGE 2 OR ABOVE FOR GENITAL DEVELOPMENT AND PUBIC HAIR GROWTH. RENEWAL: 1) TANNER STAGING AT INITIAL DIAGNOSIS HAS STABILIZED OR REGRESSED DURING THREE SEPARATE MEDICAL VISITS IN THE PREVIOUS YEAR, AND 2) HAS NOT REACHED ACTUAL AGE WHICH CORRESPONDS TO CURRENT PUBERTAL AGE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

LEVODOPA

Products Affected

• INBRIJA INHALATION CAPSULE, W/INHALATION DEVICE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PARKINSONS DISEASE (PD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	PD: INITIAL: 1) NOT CURRENTLY TAKING MORE THAN 1600MG OF LEVODOPA PER DAY, AND 2) PHYSICIAN HAS OPTIMIZED DRUG THERAPY FOR PARKINSONS DISEASE. RENEWAL: IMPROVEMENT WITH MOTOR FLUCTUATIONS DURING OFF EPISODES WITH THE USE OF INBRIJA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

L-GLUTAMINE

Products Affected

• ENDARI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	SICKLE CELL DISEASE(SCD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST
Coverage Duration	INITIAL: 12 MONTHS. RENEWAL: LIFETIME.
Other Criteria	SCD: INITIAL: AGES 18 YEARS OR OLDER: 1) AT LEAST 2 SICKLE CELL CRISES IN THE PAST YEAR, 2) SICKLE-CELL ASSOCIATED SYMPTOMS WHICH ARE INTERFERING WITH ACTIVITIES OF DAILY LIVING, OR 3) HISTORY OF OR HAS RECURRENT ACUTE CHEST SYNDROME. AGES 5 TO 17 YEARS: APPROVED WITHOUT ADDITIONAL CRITERIA. RENEWAL: MAINTAINED OR EXPERIENCED A REDUCTION IN ACUTE COMPLICATIONS OF SCD.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LIDOCAINE OINTMENT

Products Affected

• lidocaine topical ointment

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION. THIS DRUG MAY BE EITHER BUNDLED WITH AND COVERED UNDER END STAGE RENAL DISEASE DIALYSIS RELATED SERVICES OR COVERED UNDER MEDICARE D DEPENDING UPON THE CIRCUMSTANCES.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

LIDOCAINE PATCH

- lidocaine topical adhesive patch, medicated 5 %
- ZTLIDO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

LIDOCAINE PRILOCAINE

Products Affected

• lidocaine-prilocaine topical cream

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG MAY BE EITHER BUNDLED WITH AND COVERED UNDER END STAGE RENAL DISEASE DIALYSIS RELATED SERVICES OR COVERED UNDER MEDICARE D DEPENDING UPON THE CIRCUMSTANCES. INFORMATION MAY NEED TO BE SUBMITTED DESCRIBING THE USE AND SETTING OF THE DRUG TO MAKE THE DETERMINATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

LIDOCAINE SOLUTION

Products Affected

 lidocaine hcl mucous membrane solution 4 % (40 mg/ml)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

LOMITAPIDE

Products Affected

• JUXTAPID ORAL CAPSULE 10 MG, 20 MG, 30 MG, 40 MG, 5 MG, 60 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST, ENDOCRINOLOGIST OR LIPIDOLOGIST.
Coverage Duration	12 MONTHS

PA Criteria	Criteria Details
Other Criteria	1) DIAGNOSIS DETERMINED BY A) DEFINITE SIMON BROOME DIAGNOSTIC CRITERIA, OR B) DUTCH LIPID NETWORK CRITERIA SCORE OF 8 OR GREATER, OR C) CLINICAL DIAGNOSIS BASED ON A HISTORY OF AN UNTREATED LDL-C CONCENTRATION GREATER THAN 500 MG/DL TOGETHER WITH EITHER XANTHOMA BEFORE 10 YEARS OF AGE, OR EVIDENCE OF HEFH IN BOTH PARENTS. 2) LDL-C LEVEL GREATER THAN OR EQUAL TO 70MG/DL WHILE ON MAXIMAL DRUG TREATMENT. 3) TRIAL OF EVOLOCUMAB UNLESS THE PATIENT HAS NON- FUNCTIONING LDL RECEPTORS. 4) MEETS ONE OF THE FOLLOWING: A) TAKING A HIGH-INTENSITY STATIN (I.E., ATOR VASTATIN 40-80MG DAILY, ROSUVASTATIN 20-40MG DAILY) FOR A DURATION OF AT LEAST 8 WEEKS, B) TAKING A MAXIMALLY TOLERATED DOSE OF ANY STATIN FOR A DURATION OF AT LEAST 8 WEEKS GIVEN THAT THE PATIENT CANNOT TOLERATE A HIGH- INTENSITY STATIN, C) ABSOLUTE CONTRAINDICATION TO STATIN THERAPY (E.G., ACTIVE DECOMPENSATED LIVER DISEASE, NURSING FEMALE, PREGNANCY OR PLANS TO BECOME PREGNANT, HYPERSENSITIVITY REACTIONS), D) STATIN INTOLERANCE, OR E) TRIAL OF ROSUVASTATIN, ATORVASTATIN, OR STATIN THERAPY AT ANY DOSE AND HAS EXPERIENCED SKELETAL- MUSCLE RELATED SYMPTOMS (E.G., MYOPATHY).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LONCASTUXIMAB TESIRINE-LPYL

Products Affected

• ZYNLONTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LORLATINIB

Products Affected

 LORBRENA ORAL TABLET 100 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LUMACAFTOR-IVACAFTOR

- ORKAMBI ORAL GRANULES IN PACKET
- ORKAMBI ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: CYSTIC FIBROSIS (CF): CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CF.
Age Restrictions	
Prescriber Restrictions	CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CF EXPERT.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: LIFETIME.
Other Criteria	CF: RENEWAL: MAINTAINED, IMPROVED, OR DEMONSTRATED LESS THAN EXPECTED DECLINE IN FEV1 OR BODY MASS INDEX (BMI), OR REDUCTION IN NUMBER OF PULMONARY EXACERBATIONS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LUMASIRAN

Products Affected

• OXLUMO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

MACITENTAN

Products Affected

• OPSUMIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS.
Age Restrictions	
Prescriber Restrictions	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

MARGETUXIMAB-CMKB

Products Affected

• MARGENZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

MECHLORETHAMINE

Products Affected

• VALCHLOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

MEPOLIZUMAB

- NUCALA SUBCUTANEOUS AUTO-INJECTOR
- NUCALA SUBCUTANEOUS RECON SOLN
- NUCALA SUBCUTANEOUS SYRINGE 100 MG/ML, 40 MG/0.4 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: ASTHMA: BLOOD EOSINOPHIL LEVEL GREATER THAN OR EQUAL TO 150 CELLS/MCL WITHIN THE PAST 12 MONTHS.
Age Restrictions	
Prescriber Restrictions	INITIAL: ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN PULMONARY OR ALLERGY MEDICINE. NASAL POLYPS: PRESCRIBED BY OR IN CONSULTATION WITH AN OTOLARYNGOLOGIST, ALLERGIST OR IMMUNOLOGIST.
Coverage Duration	INITIAL: ASTHMA: 4 MO. NASAL POLYPS: 6 MO. OTHERS: 12 MO. RENEWAL: NASAL POLYPS, ASTHMA: 12 MO.

PA Criteria	Criteria Details
Other Criteria	INITIAL: ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY TOLERATED DOSE OF AN INHALED CORTICOSTEROID AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 2) ONE OF THE FOLLOWING: (A) AT LEAST ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR (B) POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA, AND 3) NOT CONCURRENTLY RECEIVING XOLAIR, DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA. NASAL POLYPS: PREVIOUS 56 DAY TRIAL OF ONE TOPICAL NASAL CORTICOSTEROID. RENEWAL: ASTHMA: 1) NOT CONCURRENTLY RECEIVING XOLAIR, DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA. OLOTINUED USE OF ICS AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY ONE OF THE FOLLOWING: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS, OR (D) INCREASE IN PERCENT PREDICTED FEVI FROM
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

METHYLNALTREXONE INJECTABLE

Products Affected

- RELISTOR SUBCUTANEOUS
 SOLUTION
- RELISTOR SUBCUTANEOUS SYRINGE 12 MG/0.6 ML, 8 MG/0.4 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	ADVANCED ILLNESS: 6 MONTHS. CHRONIC NON-CANCER PAIN: 12 MONTHS.
Other Criteria	CHRONIC NON-CANCER PAIN: 1) HAS BEEN TAKING OPIOIDS FOR AT LEAST 4 WEEKS, AND 2) TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENTS: NALOXEGOL (MOVANTIK) AND LUBIPROSTONE (AMITIZA)
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

METHYLNALTREXONE ORAL

Products Affected

RELISTOR ORAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	OPIOID INDUCED CONSTIPATION WITH CHRONIC NON- CANCER PAIN: 1) HAS BEEN TAKING OPIOIDS FOR AT LEAST 4 WEEKS, AND 2) TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENTS: NALOXEGOL (MOVANTIK) AND LUBIPROSTONE (AMITIZA)
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

MIDOSTAURIN

Products Affected

• RYDAPT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	ACUTE MYELOID LEUKEMIA: 6 MONTHS. ADVANCED SYSTEMIC MASTOCYTOSIS: 12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

MIFEPRISTONE

Products Affected

- KORLYM
- mifepristone oral tablet 300 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	CUSHINGS SYNDROME (CS): INITIAL: DIAGNOSIS CONFIRMED BY ONE OF THE FOLLOWING: 1) 24-HR URINE FREE CORTISOL (2 OR MORE TESTS TO CONFIRM), 2) OVERNIGHT 1MG DEXAMETHASONE TEST, OR 3) LATE NIGHT SALIVARY CORTISOL (2 OR MORE TESTS TO CONFIRM).
Age Restrictions	
Prescriber Restrictions	CS: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	INITIAL AND RENEWAL: 12 MONTHS.
Other Criteria	CS: INITIAL: HYPERCORTISOLISM IS NOT A RESULT OF CHRONIC GLUCOCORTICOIDS. RENEWAL: 1) CONTINUES TO HAVE IMPROVEMENT OF GLUCOSE TOLERANCE OR STABLE GLUCOSE TOLERANCE (E.G., REDUCED A1C, IMPROVED FASTING GLUCOSE, ETC.), 2) CONTINUES TO HAVE TOLERABILITY TO KORLYM, AND 3) CONTINUES TO NOT BE A CANDIDATE FOR SURGICAL TREATMENT OR HAS FAILED SURGERY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

MIGALASTAT

Products Affected

• GALAFOLD

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	FABRY DISEASE: INITIAL SYMPTOMATIC OR EVIDENCE OF INJURY FROM GL-3 TO THE KIDNEY, HEART, OR CENTRAL NERVOUS SYSTEM RECOGNIZED BY LABORATORY, HISTOLOGICAL, OR IMAGING FINDINGS.
Age Restrictions	
Prescriber Restrictions	FABRY DISEASE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEPHROLOGIST, CARDIOLOGIST, OR SPECIALIST IN GENETICS OR INHERITED METABOLIC DISORDERS.
Coverage Duration	INITIAL: 6 MOS. RENEWAL: 12 MOS.
Other Criteria	FABRY DISEASE: INITIAL: NOT CONCURRENTLY USING ENZYME REPLACEMENT THERAPY (I.E. FABRAZYME). RENEWAL: 1) IMPROVEMENT OR STABILIZATION, AND 2) NOT CONCURRENTLY USING ENZYME REPLACEMENT THERAPY (I.E. FABRAZYME).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

MIGLUSTAT

Products Affected

- miglustat
- yargesa

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

MILTEFOSINE

Products Affected

• IMPAVIDO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

MOBOCERTINIB

Products Affected

• EXKIVITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

MOMELOTINIB

Products Affected

• OJJAARA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

MOSUNETUZUMAB-AXGB

Products Affected

• LUNSUMIO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	RELAPSED OR REFRACTORY FOLLICULAR LYMPHOMA: INITIAL: 6 MONTHS. RENEWAL: 7 MONTHS.
Other Criteria	RELAPSED OR REFRACTORY FOLLICULAR LYMPHOMA: RENEWAL: 1) HAS ACHIEVED A PARTIAL RESPONSE TO TREATMENT, AND 2) HAS NOT PREVIOUSLY RECEIVED MORE THAN 17 CYCLES OF TREATMENT. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

NAFARELIN

Products Affected

• SYNAREL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: ENDOMETRIOSIS: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS. CENTRAL PRECOCIOUS PUBERTY (CPP): FEMALES: ELEVATED LEVELS OF FOLLICLE- STIMULATING HORMONE (FSH) GREATER THAN 4.0 MIU/ML AND LUTEINIZING HORMONE (LH) LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS. MALES: ELEVATED LEVELS OF FSH GREATER THAN 5.0 MIU/ML AND LH LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS.
Age Restrictions	
Prescriber Restrictions	INITIAL: ENDOMETRIOSIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST. CPP: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	ENDOMETRIOSIS: 6 MONTHS. CPP: INITIAL/RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: ENDOMETRIOSIS: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, 2) TRIAL OF OR CONTRAINDICATION TO NSAID AND PROGESTIN- CONTAINING PREPARATION, AND 3) HAS NOT RECEIVED A TOTAL OF 6 MONTHS OF TREATMENT PER LIFETIME. CPP: FEMALES: 1) YOUNGER THAN 8 YEARS OF AGE AT ONSET OF CPP, AND 2) AT TANNER STAGE 2 OR ABOVE FOR BREAST DEVELOPMENT AND PUBIC HAIR GROWTH. MALES: 1) YOUNGER THAN 9 YEARS OF AGE AT ONSET OF CPP, AND 2) AT TANNER STAGE 2 OR ABOVE FOR GENITAL DEVELOPMENT AND PUBIC HAIR GROWTH. RENEWAL: CPP: 1) TANNER STAGE 2 OR ABOVE FOR GENITAL DEVELOPMENT AND PUBIC HAIR GROWTH. RENEWAL: CPP: 1) TANNER STAGING AT INITIAL DIAGNOSIS HAS STABILIZED OR REGRESSED DURING THREE SEPARATE MEDICAL VISITS IN THE PREVIOUS YEAR, AND 2) HAS NOT REACHED ACTUAL AGE WHICH CORRESPONDS TO CURRENT PUBERTAL AGE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

NARCOLEPSY AGENTS

Products Affected

- armodafinil
- modafinil oral tablet 100 mg, 200 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

NATALIZUMAB

Products Affected

• TYSABRI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: CROHNS DISEASE (CD): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	MULTIPLE SCLEROSIS (MS): 12 MOS. CD: INITIAL: 6 MOS, RENEWAL: 12 MOS.
Other Criteria	INITIAL: MS: TRIAL OF TWO AGENTS INDICATED FOR THE TREATMENT OF MS. CD: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, SKYRIZI, RINVOQ. RENEWAL: CD: 1) RECEIVED AT LEAST 12 MONTHS OF THERAPY WITH TYSABRI AND HAS NOT REQUIRED MORE THAN 3 MONTHS OF CORTICOSTEROID USE WITHIN THE PAST 12 MONTHS TO CONTROL CROHNS DISEASE WHILE ON TYSABRI, OR 2) HAS ONLY RECEIVED 6 MONTHS OF THERAPY WITH TYSABRI AND HAS TAPERED OFF CORTICOSTEROIDS DURING THE FIRST 24 WEEKS OF TYSABRI THERAPY. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

NAXITAMAB-GQGK

Products Affected

• DANYELZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

NEDOSIRAN

Products Affected

• RIVFLOZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

NERATINIB

Products Affected

• NERLYNX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	EARLY-STAGE (STAGE I-III) BREAST CANCER: MEDICATION IS BEING REQUESTED WITHIN 2 YEARS OF COMPLETING THE LAST TRASTUZUMAB DOSE. ALL OTHER FDA APPROVED INDICATIONS ARE COVERED WITHOUT ADDITIONAL CRITERIA, EXCEPT THOSE CRITERIA IN THE FDA APPROVED LABEL.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

NILOTINIB

Products Affected

• TASIGNA ORAL CAPSULE 150 MG, 200 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PREVIOUSLY TREATED CML: MUTATIONAL ANALYSIS PRIOR TO INITIATION AND TASIGNA IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

NINTEDANIB

Products Affected

• OFEV

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: IDIOPATHIC PULMONARY FIBROSIS (IPF): 1) A USUAL INTERSTITIAL PNEUMONIA (UIP) PATTERN AS EVIDENCED BY HIGH-RESOLUTION COMPUTED TOMOGRAPHY (HRCT) ALONE OR VIA A COMBINATION OF SURGICAL LUNG BIOPSY AND HRCT, AND 2) BASELINE FORCED VITAL CAPACITY (FVC) AT LEAST 50% OF PREDICTED VALUE. SYSTEMIC SCLEROSIS- ASSOCIATED INTERSTITIAL LUNG DISEASE (SSC-ILD): 1) AT LEAST 10% FIBROSIS ON A CHEST HRCT, AND 2) BASELINE FVC AT LEAST 40% OF PREDICTED VALUE. CHRONIC FIBROSING INTERSTITIAL LUNG DISEASE WITH A PROGRESSIVE PHENOTYPE (PF-ILD): 1) AT LEAST 10% FIBROSIS ON A CHEST HRCT, AND 2) BASELINE FVC AT LEAST 45% OF PREDICTED VALUE.
Age Restrictions	
Prescriber Restrictions	INITIAL: IPF: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST. SSC-ILD, PF-ILD: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR RHEUMATOLOGIST.
Coverage Duration	INITIAL: SSC-ILD: 6 MOS. IPF, PF-ILD: 12 MOS. RENEWAL (ALL INDICATIONS): 12 MOS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: IPF: 1) DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., CONNECTIVE TISSUE DISEASE, DRUG TOXICITY, ASBESTOS OR BERYLLIUM EXPOSURE, HYPERSENSITIVITY PNEUMONITIS), AND 2) TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: ESBRIET (PIRFENIDONE). SSC-ILD: 1) DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., HEART FAILURE/FLUID OVERLOAD, DRUG- INDUCED LUNG TOXICITY, RECURRENT ASPIRATION), AND 2) TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: ACTEMRA SUBQ. PF-ILD: LUNG FUNCTION AND RESPIRATORY SYMPTOMS OR CHEST IMAGING HAVE WORSENED/PROGRESSED DESPITE TREATMENT WITH MEDICATIONS USED IN CLINICAL PRACTICE FOR ILD (NOT ATTRIBUTABLE TO COMORBIDITIES SUCH AS INFECTION, HEART FAILURE). RENEWAL: IPF, SSC-ILD, PF-ILD: CLINICAL MEANINGFUL IMPROVEMENT OR MAINTENANCE IN ANNUAL RATE OF DECLINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

NIRAPARIB

Products Affected

- ZEJULA ORAL CAPSULE
- ZEJULA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	RECURRENT EPITHELIAL OVARIAN, FALLOPIAN TUBE, OR PRIMARY PERITONEAL CANCER: 1) ZEJULA WILL BE USED AS MONOTHERAPY, AND 2) ZEJULA IS STARTED NO LATER THAN 8 WEEKS AFTER THE MOST RECENT PLATINUM-CONTAINING REGIMEN.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

NIRAPARIB/ABIRATERONE

Products Affected

• AKEEGA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (MCRPC): 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

NIROGACESTAT

Products Affected

• OGSIVEO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

NIRSEVIMAB-ALIP

Products Affected

• BEYFORTUS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	RESPIRATORY SYNCYTIAL VIRUS (RSV) PREVENTION: 1) HAS NOT COMPLETED A COURSE OF SYNAGIS OR HAS NOT RECEIVED A DOSE OF SYNAGIS IN THE PAST 30 DAYS, AND WILL NOT RECEIVE FURTHER DOSES OF SYNAGIS WITHIN THE SAME RSV SEASON, AND 2) HAS NOT RECEIVED MORE THAN 2 DOSES OF BEYFORTUS PER LIFETIME, UNLESS HAS UNDERGONE OR WILL UNDERGO A CARDIAC SURGERY WITH CARDIOPULMONARY BYPASS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

NITISINONE

Products Affected

- nitisinone
- ORFADIN ORAL SUSPENSION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HEREDITARY TYROSINEMIA TYPE 1 (HT-1): INITIAL: DIAGNOSIS CONFIRMED BY ELEVATED URINARY OR PLASMA SUCCINYLACETONE LEVELS OR A MUTATION IN THE FUMARYLACETOACETATE HYDROLASE GENE. RENEWAL: URINARY OR PLASMA SUCCINYLACETONE LEVELS HAVE DECREASED FROM BASELINE WHILE ON TREATMENT WITH NITISINONE.
Age Restrictions	
Prescriber Restrictions	HT-1: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PRESCRIBER SPECIALIZING IN INHERITED METABOLIC DISEASES.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	HT-1: INITIAL: ORFADIN SUSPENSION: TRIAL OF OR CONTRAINDICATION TO PREFERRED NITISINONE TABLETS OR CAPSULES.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

NIVOLUMAB

Products Affected

• OPDIVO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	UNRESECTABLE OR METASTATIC MELANOMA: NO CONCURRENT USE WITH TARGETED THERAPY (I.E., BRAF INHIBITORS, MEK INHIBITORS, AND NTRK INHIBITORS).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

NIVOLUMAB-RELATLIMAB-RMBW

Products Affected

• OPDUALAG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

OBETICHOLIC ACID

Products Affected

• OCALIVA

PA Criteria	Criteria Details
Exclusion	PRIMARY BILIARY CHOLANGITIS (PBC):
Criteria	INITIAL/RENEWAL: COMPLETE BILIARY OBSTRUCTION.
Required Medical	PBC: INITIAL: DIAGNOSIS CONFIRMED BY TWO OF THE
Information	FOLLOWING: 1) ALKALINE PHOSPHATASE LEVEL OF AT
	LEAST 1.5 TIMES THE UPPER LIMIT OF NORMAL, 2)
	PRESENCE OF ANTIMITOCHONDRIAL ANTIBODIES AT A
	TITER OF 1:40 OR HIGHER, OR 3) HISTOLOGIC EVIDENCE
	OF NON-SUPPURATIVE DESTRUCTIVE CHOLANGITIS AND DESTRUCTION OF INTERLOBULAR BILE DUCTS.
	DESTRUCTION OF INTERLOBULAR BILE DUCIS.
Age Restrictions	
Prescriber	PBC: INITIAL: PRESCRIBED BY OR IN CONSULTATION
Restrictions	WITH A GASTROENTEROLOGIST OR HEPATOLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS
Other Criteria	PBC: INITIAL: USED IN COMBINATION WITH
	URSODEOXYCHOLIC ACID IN A PATIENT WITH AN
	INADEQUATE RESPONSE TO URSODEOXYCHOLIC ACID,
	OR AS MONOTHERAPY IN A PATIENT WHO IS UNABLE TO
	TOLERATE URSODEOXYCHOLIC ACID. RENEWAL:
	CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B	No
Prerequisite	

OCRELIZUMAB

Products Affected

• OCREVUS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	RELAPSING FORM OF MULTIPLE SCLEROSIS (MS): TRIAL OF TWO AGENTS INDICATED FOR THE TREATMENT OF RELAPSING FORMS OF MS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

OFATUMUMAB-SQ

Products Affected

• KESIMPTA PEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

OLANZAPINE/SAMIDORPHAN

Products Affected

• LYBALVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	SCHIZOPHRENIA/BIPOLAR I: PRESCRIBED BY OR IN CONSULTATION WITH A PSYCHIATRIST
Coverage Duration	12 MONTHS
Other Criteria	SCHIZOPHRENIA: (1) PATIENT IS AT HIGH RISK OF WEIGHT GAIN AND (2) TRIAL OF OR CONTRAINDICATION TO LATUDA OR ONE OF THE FOLLOWING ORAL ANTIPSYCHOTICS: RISPERIDONE, CLOZAPINE TABLET, OLANZAPINE, IMMEDIATE RELEASE QUETIAPINE FUMARATE, ZIPRASIDONE, ARIPIPRAZOLE. BIPOLAR I: (1) PATIENT IS AT HIGH RISK OF WEIGHT GAIN AND (2) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING ORAL ANTIPSYCHOTICS: RISPERIDONE, OLANZAPINE, IMMEDIATE RELEASE QUETIAPINE FUMARATE, ZIPRASIDONE, ARIPIPRAZOLE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

OLAPARIB

Products Affected

• LYNPARZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	RECURRENT EPITHELIAL OVARIAN, FALLOPIAN TUBE OR PRIMARY PERITONEAL CANCER: MEDICATION WILL BE USED AS MONOTHERAPY. METASTATIC CASTRATION- RESISTANT PROSTATE CANCER: 1) PREVIOUSLY RECEIVED A BILATERAL ORCHIECTOMY, OR 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. ALL OTHER FDA APPROVED INDICATIONS ARE COVERED WITHOUT ADDITIONAL CRITERIA, EXCEPT THOSE CRITERIA IN THE FDA APPROVED LABEL.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

OLUTASIDENIB

Products Affected

• REZLIDHIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

OMACETAXINE

Products Affected

• SYNRIBO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

OMALIZUMAB

Products Affected

• XOLAIR SUBCUTANEOUS AUTO-INJECTOR 300 MG/2 ML

• XOLAIR SUBCUTANEOUS AUTO- • XOLAIR SUBCUTANEOUS SYRINGE

 XOLAIR SUBCUTANEOUS RECON SOLN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: ASTHMA: POSITIVE SKIN PRICK OR BLOOD TEST (E.G., ELISA, FEIA) TO A PERENNIAL AEROALLERGEN AND A BASELINE IGE SERUM LEVEL GREATER THAN OR EQUAL TO 30 IU/ML.
Age Restrictions	
Prescriber Restrictions	INITIAL AND RENEWAL: CHRONIC SPONTANEOUS URTICARIA (CSU): PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE, DERMATOLOGY OR IMMUNOLOGY. INITIAL: NASAL POLYPS: PRESCRIBED BY OR IN CONSULTATION WITH AN OTOLARYNGOLOGIST, ALLERGIST OR IMMUNOLOGIST. ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE.
Coverage Duration	INITIAL: ASTHMA: 4 MO. CSU, NASAL POLYPS: 6 MO. RENEWAL: ASTHMA, NASAL POLYPS: 12 MO. CSU: 6 MO.

PA Criteria	Criteria Details
Other Criteria	INITIAL: CSU: TRIAL OF OR CONTRAINDICATION TO A MAXIMALLY TOLERATED DOSE OF AN H1 ANTI- HISTAMINE AND STILL EXPERIENCES HIVES ON MOST DAYS OF THE WEEK. NASAL POLYPS: 1) PREVIOUS 56 DAY TRIAL OF ONE TOPICAL NASAL CORTICOSTEROID, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE PREFERRED AGENT: NUCALA, DUPIXENT. ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY TOLERATED DOSE OF AN INHALED CORTICOSTEROID AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, 2) ONE OF THE FOLLOWING: (A) PATIENT EXPERIENCED AT LEAST ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR (B) PATIENT HAS POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA, 3) NOT CONCURRENTLY RECEIVING DUPIXENT OR ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA. RENEWAL: CSU: DIAGNOSIS OF CSU. NASAL POLYPS: CLINICAL BENEFIT COMPARED TO BASELINE. ASTHMA: 1) NOT CONCURRENTLY RECEIVING DUPIXENT OR ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA, 2) CONTINUED USE OF ICS AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY ONE OF THE FOLLOWING: (A) REDUCTION IN ASTHMA
	EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS, OR (D) INCREASE IN PERCENT PREDICTED FEV1 FROM PRETREATMENT BASELINE.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

OPICAPONE

Products Affected

• ONGENTYS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	PARKINSONS DISEASE: 18 YEARS OF AGE OR OLDER
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

OSIMERTINIB

Products Affected

• TAGRISSO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	EGFR EXON 19 DELETIONS OR EXON 21 L858R MUTATIONS NON-SMALL CELL LUNG CANCER (NSCLC) AND METASTATIC NSCLC WITH EGFR T790M MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

OXANDROLONE

Products Affected

• oxandrolone

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	PROTEIN CATABOLISM, BONE PAIN: 1) MONITORED FOR PELIOSIS HEPATIS, LIVER CELL TUMORS, AND BLOOD LIPID CHANGES, 2) DOES NOT HAVE KNOWN OR SUSPECTED: CARCINOMA OF THE PROSTATE OR BREAST IN MALE PATIENTS, CARCINOMA OF THE BREAST IN FEMALES WITH HYPERCALCEMIA, NEPHROSIS (THE NEPHROTIC PHASE OF NEPHRITIS), OR HYPERCALCEMIA, AND 3) DOES NOT HAVE SEVERE HEPATIC DYSFUNCTION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PACRITINIB

Products Affected

• VONJO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	MYELOFIBROSIS: RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PALBOCICLIB

Products Affected

• IBRANCE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ADVANCED OR METASTATIC BREAST CANCER: TRIAL OF OR CONTRAINDICATION TO ONE OF THE PREFERRED AGENTS, WHERE INDICATIONS ALIGN: KISQALI, VERZENIO.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PALIVIZUMAB

Products Affected

• SYNAGIS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: GESTATIONAL AGE
Age Restrictions	INITIAL AND RENEWAL: LESS THAN 24 MONTHS OF AGE.
Prescriber Restrictions	
Coverage Duration	CRITERIA CONSISTENT WITH THE CDC RSV CENSUS REGIONAL TREND. SEE OTHER CRITERIA.
Other Criteria	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT RECOMMENDATIONS FROM THE AMERICAN ACADEMY OF PEDIATRICS FOR PALIVIZUMAB PROPHYLAXIS FOR RESPIRATORY SYNCYTIAL VIRUS (RSV) INFECTIONS AND THE CENTERS FOR DISEASE CONTROL AND PREVENTION RSV CENSUS REGIONAL TREND. APPROVAL WILL BE FOR UP TO A MAXIMUM OF 5 MONTHS DEPENDENT UPON REMAINING LENGTH OF RESPIRATORY SYNCYTIAL VIRUS SEASON. RENEWAL: ADDITIONAL 1 MONTH OF TREATMENT FOR CARDIOPULMONARY BYPASS SURGERY DURING RSV PROPHYLAXIS SEASON.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PANOBINOSTAT

Products Affected

• FARYDAK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS
Other Criteria	MULTIPLE MYELOMA: RENEWAL: TOLERATED THE FIRST 8 CYCLES OF THERAPY WITHOUT UNRESOLVED SEVERE OR MEDICALLY SIGNIFICANT TOXICITY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PARATHYROID HORMONE

Products Affected

• NATPARA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	HYPOCALCEMIA SECONDARY TO HYPOPARATHYROIDISM: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	HYPOCALCEMIA SECONDARY TO HYPOPARATHYROIDISM: 1) TRIAL OF OR CONTRAINDICATION TO CALCITRIOL, 2) HYPOPARATHYROIDISM IS NOT DUE TO A CALCIUM SENSING RECEPTOR (CSR) MUTATION, AND 3) HYPOPARATHYROIDISM IS NOT CONSIDERED ACUTE POST-SURGICAL HYPOPARATHYROIDISM.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PASIREOTIDE DIASPARTATE

Products Affected

• SIGNIFOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	CUSHINGS DISEASE (CD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	CD: RENEWAL: 1) CONTINUED IMPROVEMENT OF CUSHINGS DISEASE, AND 2) MAINTAINED TOLERABILITY TO SIGNIFOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PAZOPANIB

Products Affected

• pazopanib

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ADVANCED SOFT TISSUE SARCOMA (STS): NOT USED FOR ADIPOCYTIC STS OR GASTROINTESTINAL STROMAL TUMORS (GIST)
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PEGFILGRASTIM - APGF

Products Affected

• NYVEPRIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NON MYELOID MALIGNANCY: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PEGFILGRASTIM - CBQV

Products Affected

- UDENYCA
- UDENYCA AUTOINJECTOR
- UDENYCA ONBODY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NON MYELOID MALIGNANCY, ACUTE RADIATION EXPOSURE: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	NON MYELOID MALIGNANCY: TRIAL OF OR CONTRAINDICATION TO NYVEPRIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PEGFILGRASTIM - JMDB

Products Affected

• FULPHILA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NON MYELOID MALIGNANCY: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	NON MYELOID MALIGNANCY: TRIAL OF OR CONTRAINDICATION TO NYVEPRIA
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PEGFILGRASTIM-BMEZ

Products Affected

• ZIEXTENZO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NON MYELOID MALIGNANCY: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	NON MYELOID MALIGNANCY: TRIAL OF OR CONTRAINDICATION TO NYVEPRIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PEGFILGRASTIM-FPGK

Products Affected

• STIMUFEND

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NON MYELOID MALIGNANCY, ACUTE RADIATION EXPOSURE: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	NON MYELOID MALIGNANCY: TRIAL OF OR CONTRAINDICATION TO NYVEPRIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PEGFILGRASTIM-NEULASTA ONPRO

Products Affected

• NEULASTA ONPRO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NON MYELOID MALIGNANCY, ACUTE RADIATION EXPOSURE: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PEGFILGRASTIM-PBBK

Products Affected

• FYLNETRA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NON MYELOID MALIGNANCY: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	NON MYELOID MALIGNANCY: TRIAL OF OR CONTRAINDICATION TO PREFERRED AGENT: NYVEPRIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PEGINTERFERON ALFA-2A

Products Affected

• PEGASYS

PA Criteria	Criteria Details
Exclusion Criteria	CHRONIC HEPATITIS C VIRUS INFECTION. CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	HEPATITIS B: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST, INFECTIOUS DISEASE SPECIALIST, OR PHYSICIAN SPECIALIZING IN THE TREATMENT OF HEPATITIS (E.G., HEPATOLOGIST).
Coverage Duration	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
Other Criteria	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PEGUNIGALSIDASE ALFA-IWXJ

Products Affected

• ELFABRIO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	FABRY DISEASE: INITIAL: 1) SYMPTOMATIC OR EVIDENCE OF INJURY TO THE KIDNEY, HEART, OR CENTRAL NERVOUS SYSTEM RECOGNIZED BY LABORATORY, HISTOLOGICAL, OR IMAGING FINDINGS, AND 2) ONE OF THE FOLLOWING: (A) FEMALES: GALACTOSIDASE ALPHA (GLA) GENE MUTATION VIA GENETIC TESTING, OR (B) MALES: ENZYME ASSAY INDICATING ALPHA GALACTOSIDASE A DEFICIENCY OR GLA GENE MUTATION VIA GENETIC TESTING.
Age Restrictions	
Prescriber Restrictions	FABRY DISEASE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH NEPHROLOGIST, CARDIOLOGIST, OR SPECIALIST IN GENETICS OR INHERITED METABOLIC DISORDERS.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	FABRY DISEASE: INITIAL: NO CONCURRENT USE WITH ANOTHER FABRY DISEASE THERAPY. RENEWAL: 1) DEMONSTRATED IMPROVEMENT OR STABILIZATION, AND 2) NO CONCURRENT USE WITH ANOTHER FABRY DISEASE THERAPY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PEGVALIASE-PQPZ

Products Affected

• PALYNZIQ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	PHENYLKETONURIA (PKU): INITIAL: NOT ON CONCURRENT TREATMENT WITH KUVAN. RENEWAL: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NOT ON CONCURRENT TREATMENT WITH KUVAN.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PEGVISOMANT

Products Affected

• SOMAVERT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PEMBROLIZUMAB

Products Affected

• KEYTRUDA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	UNRESECTABLE OR METASTATIC MELANOMA: NO CONCURRENT USE WITH TARGETED THERAPY (I.E., BRAF INHIBITORS, MEK INHIBITORS, AND NTRK INHIBITORS).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PEMIGATINIB

Products Affected

• PEMAZYRE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	CHOLANGIOCARCINOMA, MYELOID/LYMPHOID NEOPLASMS: COMPREHENSIVE OPHTHALMOLOGICAL EXAMINATION, INCLUDING OPTICAL COHERENCE TOMOGRAPHY (OCT), WILL BE COMPLETED PRIOR TO INITIATION OF THERAPY AND AT THE RECOMMENDED SCHEDULED INTERVALS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PENICILLAMINE TABLET

Products Affected

• penicillamine oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: CYSTINURIA: HAS NEPHROLITHIASIS AND ONE OF THE FOLLOWING: 1) STONE ANALYSIS SHOWING PRESENCE OF CYSTINE, 2) PRESENCE OF PATHOGNOMONIC HEXAGONAL CYSTINE CRYSTALS ON URINALYSIS, OR 3) FAMILY HISTORY OF CYSTINURIA AND POSITIVE CYANIDE-NITROPRUSSIDE SCREENING.
Age Restrictions	
Prescriber Restrictions	INITIAL: WILSONS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A HEPATOLOGIST OR GASTROENTEROLOGIST. CYSTINURIA: PRESCRIBED BY OR IN CONSULTATION WITH A NEPHROLOGIST. RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	INITIAL: 12 MONTHS, RENEWAL: LIFETIME.

PA Criteria	Criteria Details
Other Criteria	INITIAL: WILSONS DISEASE: 1) LEIPZIG SCORE OF 4 OR GREATER. RA: 1) NO HISTORY OR OTHER EVIDENCE OF RENAL INSUFFICIENCY, AND 2) TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. RENEWAL: RA: 1) NO HISTORY OR OTHER EVIDENCE OF RENAL INSUFFICIENCY, AND 2) EXPERIENCED OR MAINTAINED IMPROVEMENT IN TENDER JOINT COUNT OR SWOLLEN JOINT COUNT COMPARED TO BASELINE. WILSONS DISEASE, CYSTINURIA: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PEXIDARTINIB

Products Affected

• TURALIO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PIMAVANSERIN

Products Affected

• NUPLAZID

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	PSYCHOSIS IN PARKINSONS DISEASE (PD): INITIAL: 18 YEARS OR OLDER
Prescriber Restrictions	PSYCHOSIS IN PD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, GERIATRICIAN, OR A BEHAVIORAL HEALTH SPECIALIST (E.G., PSYCHIATRIST).
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	PSYCHOSIS IN PD: RENEWAL: IMPROVEMENT IN PSYCHOSIS SYMPTOMS FROM BASELINE AND DEMONSTRATES A CONTINUED NEED FOR TREATMENT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PIRFENIDONE

Products Affected

- pirfenidone oral capsule
 pirfenidone oral tablet 267 mg, 534 mg, 801 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	IDIOPATHIC PULMONARY FIBROSIS (IPF): INITIAL: 1) A USUAL INTERSTITIAL PNEUMONIA (UIP) PATTERN AS EVIDENCED BY HIGH-RESOLUTION COMPUTED TOMOGRAPHY (HRCT) ALONE OR VIA A COMBINATION OF SURGICAL LUNG BIOPSY AND HRCT, AND 2) PREDICTED FORCED VITAL CAPACITY (FVC) OF AT LEAST 50% AT BASELINE.
Age Restrictions	IPF: INITIAL: 18 YEARS OR OLDER.
Prescriber Restrictions	IPF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	IPF: INITIAL: 1) DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., CONNECTIVE TISSUE DISEASE, DRUG TOXICITY, ASBESTOS OR BERYLLIUM EXPOSURE, HYPERSENSITIVITY PNEUMONITIS, SYSTEMIC SCLEROSIS, RHEUMATOID ARTHRITIS, RADIATION, SARCOIDOSIS, BRONCHIOLITIS OBLITERANS ORGANIZING PNEUMONIA, HUMAN IMMUNODEFICIENCY VIRUS (HIV) INFECTION, VIRAL HEPATITIS, OR CANCER). RENEWAL: CLINICAL MEANINGFUL IMPROVEMENT OR MAINTENANCE IN ANNUAL RATE OF DECLINE.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

PIRTOBRUTINIB

Products Affected

 JAYPIRCA ORAL TABLET 100 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

POMALIDOMIDE

Products Affected

• POMALYST

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PONATINIB

Products Affected

• ICLUSIG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	CML: MUTATIONAL ANALYSIS PRIOR TO INITIATION AND ICLUSIG IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

POSACONAZOLE

Products Affected

• posaconazole oral

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	OROPHARYNGEAL CANDIDIASIS (OPC): 3 MONTHS. PROPHYLAXIS: 6 MONTHS. TREATMENT: 12 WEEKS.
Other Criteria	POSACONAZOLE SUSPENSION ONLY: 1) OPC: TRIAL OF OR CONTRAINDICATION TO FLUCONAZOLE OR ITRACONAZOLE. 2) PROPHYLAXIS OF INVASIVE ASPERGILLUS AND CANDIDA INFECTION: INABILITY TO SWALLOW TABLETS. POSACONZOLE TABLETS ONLY: 1) TREATMENT OF INVASIVE ASPERGILLOSIS, 2) PROPHYLAXIS OF INVASIVE ASPERGILLUS AND CANDIDA INFECTION: AT HIGH RISK OF DEVELOPING THESE INFECTIONS DUE TO BEING SEVERELY IMMUNOCOMPROMISED. CONTINUATION OF THERAPY AFTER HOSPITAL DISCHARGE REQUIRES NO EXTRA CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

POSACONAZOLE-POWDERMIX

Products Affected

• NOXAFIL ORAL SUSP, DELAYED RELEASE FOR RECON

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	PROPHYLAXIS OF INVASIVE ASPERGILLUS AND CANDIDA INFECTION: INABILITY TO SWALLOW TABLETS. CONTINUATION OF THERAPY AFTER HOSPITAL DISCHARGE REQUIRES NO EXTRA CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PRALSETINIB

Products Affected

• GAVRETO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PRAMLINTIDE

Products Affected

- SYMLINPEN 120
- SYMLINPEN 60

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	TYPE I OR TYPE II DIABETES: REQUIRING INSULIN OR CONTINUOUS INSULIN INFUSION (INSULIN PUMP) FOR GLYCEMIC CONTROL
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PYRIMETHAMINE

Products Affected

• pyrimethamine

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	TOXOPLASMOSIS: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN INFECTIOUS DISEASE SPECIALIST.
Coverage Duration	TOXOPLASMOSIS: INITIAL: 8 WEEKS, RENEWAL: 6 MOS.
Other Criteria	TOXOPLASMOSIS: RENEWAL: ONE OF THE FOLLOWING: (1) PERSISTENT CLINICAL DISEASE (HEADACHE, NEUROLOGICAL SYMPTOMS, OR FEVER) AND PERSISTENT RADIOGRAPHIC DISEASE (ONE OR MORE MASS LESIONS ON BRAIN IMAGING), OR (2) CD4 COUNT LESS THAN 200 CELLS/MM3 AND CURRENTLY TAKING AN ANTI-RETROVIRAL THERAPY IF HIV POSITIVE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

QUININE

Products Affected

• quinine sulfate

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

QUIZARTINIB

Products Affected

• VANFLYTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RAMUCIRUMAB

Products Affected

• CYRAMZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

REGORAFENIB

Products Affected

• STIVARGA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RELUGOLIX

Products Affected

• ORGOVYX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

REPOTRECTINIB

Products Affected

• AUGTYRO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RESLIZUMAB

Products Affected

• CINQAIR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	ASTHMA: INITIAL: BLOOD EOSINOPHIL LEVEL GREATER THAN OR EQUAL TO 150 CELLS/MCL WITHIN THE PAST 12 MONTHS.
Age Restrictions	
Prescriber Restrictions	ASTHMA: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE.
Coverage Duration	ASTHMA: INITIAL: 4 MONTHS. RENEWAL: 12 MONTHS

PA Criteria	Criteria Details
Other Criteria	ASTHMA: INITIAL: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND ONE OTHER MAINTENANCE MEDICATION, AND 2) ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS, OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA, 3) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: FASENRA, NUCALA, DUPIXENT, AND 4) NO CONCURRENT USE WITH XOLAIR, DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA. RENEWAL: 1) NO CONCURRENT USE WITH XOLAIR, DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA, 2) CONTINUED USE OF ICS AND ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA- RELATED SYMPTOMS, OR (D) INCREASE IN PERCENT PREDICTED FEVI FROM PRETREATMENT BASELINE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
UII Ladel Uses	
Part B Prerequisite	No

RETIFANLIMAB-DLWR

Products Affected

• ZYNYZ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RIBOCICLIB

Products Affected

 KISQALI ORAL TABLET 200 MG/DAY (200 MG X 1), 400 MG/DAY (200 MG X 2), 600 MG/DAY (200 MG X 3)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RIBOCICLIB-LETROZOLE

Products Affected

 KISQALI FEMARA CO-PACK ORAL TABLET 200 MG/DAY(200 MG X 1)-2.5 MG, 400 MG/DAY(200 MG X 2)-2.5 MG, 600 MG/DAY(200 MG X 3)-2.5 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RIFAXIMIN

Products Affected

• XIFAXAN ORAL TABLET 200 MG, 550 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	TRAVELERS DIARRHEA/HE: 12 MOS. IBS-D: 8 WKS.
Other Criteria	RIFAXIMIN 550 MG TABLETS: HE: TRIAL OF OR CONTRAINDICATION TO LACTULOSE OR CONCURRENT LACTULOSE THERAPY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RIMEGEPANT

Products Affected

• NURTEC ODT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: ACUTE MIGRAINE TREATMENT: TRIAL OF OR CONTRAINDICATION TO ONE TRIPTAN (E.G., SUMATRIPTAN, RIZATRIPTAN). EPISODIC MIGRAINE PREVENTION: 1) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREVENTIVE MIGRAINE TREATMENTS: DIVALPROEX SODIUM, TOPIRAMATE, PROPRANOLOL, TIMOLOL. RENEWAL: ACUTE MIGRAINE TREATMENT: 1) IMPROVEMENT FROM BASELINE IN A VALIDATED ACUTE TREATMENT PATIENT-REPORTED OUTCOME QUESTIONNAIRE, OR 2) THERAPY WORKS CONSISTENTLY IN MAJORITY OF MIGRAINE ATTACKS. EPISODIC MIGRAINE PREVENTION: 1) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION, AND 2) REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

RIOCIGUAT

Products Affected

• ADEMPAS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PULMONARY ARTERIAL HYPERTENSION (PAH): DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS. PERSISTENT/RECURRENT CHRONIC THROMBOEMBOLIC PULMONARY HYPERTENSION (CTEPH) (WHO GROUP 4): WHO FUNCTIONAL CLASS II-IV SYMPTOMS.
Age Restrictions	
Prescriber Restrictions	INITIAL: PAH, CTEPH: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: PAH: NOT CONCURRENTLY TAKING NITRATES, NITRIC OXIDE DONORS, PHOSPHODIESTERASE INHIBITORS, OR NON-SPECIFIC PDE INHIBITORS. CTEPH: 1) NOT CONCURRENTLY TAKING NITRATES, NITRIC OXIDE DONORS, PHOSPHODIESTERASE INHIBITORS, OR NON-SPECIFIC PDE INHIBITORS, AND 2) NOT A CANDIDATE FOR SURGERY OR HAS INOPERABLE CTEPH OR HAS PERSISTENT OR RECURRENT DISEASE AFTER SURGICAL TREATMENT. RENEWAL: PAH, CTEPH: NOT CONCURRENTLY TAKING NITRATES, NITRIC OXIDE DONORS, PHOSPHODIESTERASE INHIBITORS, OR NON- SPECIFIC PDE INHIBITORS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RIPRETINIB

Products Affected

• QINLOCK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RISANKIZUMAB-RZAA

Products Affected

• SKYRIZI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PLAQUE PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE OR GENITAL AREA.
Age Restrictions	
Prescriber Restrictions	INITIAL: PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. CROHNS DISEASE (CD): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG). CD: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPURINE, METHOTREXATE, OR MESALAMINE. RENEWAL: PSO, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

RISDIPLAM

Products Affected

• EVRYSDI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	SPINAL MUSCULAR ATROPHY (SMA): INITIAL: GENE MUTATION ANALYSIS INDICATING MUTATIONS OR DELETIONS OF BOTH ALLELES OF THE SURVIVAL MOTOR NEURON 1 (SMN1) GENE. FOR PRESYMPTOMATIC PATIENTS: UP TO THREE COPIES OF SURVIVAL MOTOR NEURON 2 (SMN2) BASED ON NEWBORN SCREENING.
Age Restrictions	
Prescriber Restrictions	SMA: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROMUSCULAR SPECIALIST OR SMA SPECIALIST AT A SMA SPECIALTY CENTER.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS
Other Criteria	SMA: INITIAL: FOR SYMPTOMATIC PATIENTS: (1) BASELINE MOTOR FUNCTION ASSESSMENT BY A NEUROMUSCULAR SPECIALIST OR SMA SPECIALIST, AND (2) IF PATIENT RECEIVED GENE THERAPY, THE PATIENT HAD LESS THAN EXPECTED CLINICAL BENEFIT. RENEWAL: IMPROVED, MAINTAINED, OR DEMONSTRATED LESS THAN EXPECTED DECLINE IN: (1) MOTOR FUNCTION ASSESSMENTS COMPARED TO BASELINE, OR (2) OTHER MUSCLE FUNCTION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RITUXIMAB AND HYALURONIDASE HUMAN-SQ

Products Affected

• RITUXAN HYCELA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	FOLLICULAR LYMPHOMA (FL), DIFFUSE LARGE B-CELL LYMPHOMA (DLBCL), CHRONIC LYMPHOCYTIC LEUKEMIA (CLL): HAS RECEIVED OR WILL RECEIVE AT LEAST ONE FULL DOSE OF A RITUXIMAB PRODUCT BY INTRAVENOUS INFUSION PRIOR TO INITIATION OF RITUXIMAB AND HYALURONIDASE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RITUXIMAB-ABBS

Products Affected

• TRUXIMA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. NHL, CLL: PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST.
Coverage Duration	RA: INITIAL: 6 MO, RENEWAL: 12 MO. NHL, GPA, MPA: 12 MO. CLL: 6 MO.
Other Criteria	RA: INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RITUXIMAB-ARRX

Products Affected

• RIABNI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	RHEUMATOID ARTHRITIS (RA): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. NHL, CLL: PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST.
Coverage Duration	RA: INITIAL: 6 MO, RENEWAL: 12 MO. NHL, WG, MPA: 12 MO. CLL: 6 MO.
Other Criteria	RA: INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RITUXIMAB-PVVR

Products Affected

• RUXIENCE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	RA: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. NHL, CLL: PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST.
Coverage Duration	NHL, GPA, MPA: 12 MONTHS. CLL: 6 MONTHS. RA: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	RA: INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO TWO PREFERRED AGENTS: HUMIRA, ENBREL, RINVOQ, XELJANZ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ROMIPLOSTIM

Products Affected

• NPLATE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	IMMUNE THROMBOCYTOPENIA (ITP): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR IMMUNOLOGIST.
Coverage Duration	ITP: INITIAL: 4 MO, RENEWAL: 12 MO. HEMATOPOIETIC SYNDROME OF ACUTE RADIATION SYNDROME: 12 MO.
Other Criteria	ITP: INITIAL: TRIAL OF OR CONTRAINDICATION TO CORTICOSTEROIDS, IMMUNOGLOBULINS, OR AN INSUFFICIENT RESPONSE TO SPLENECTOMY. RENEWAL: CLINICAL RESPONSE TO THERAPY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ROPEGINTERFERON ALFA-2B-NJFT

Products Affected

• BESREMI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RUCAPARIB

Products Affected

• RUBRACA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: ONE OF THE FOLLOWING: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RUXOLITINIB

Products Affected

• JAKAFI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	MYELOFIBROSIS: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. POLYCYTHEMIA VERA, GVHD: 12 MONTHS.
Other Criteria	MYELOFIBROSIS: RENEWAL: PATIENT CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SAFINAMIDE

Products Affected

• XADAGO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SAPROPTERIN

Products Affected

- *javygtor oral tablet,soluble*
- sapropterin oral tablet, soluble

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 1 MONTH, RENEWAL 12 MONTHS.
Other Criteria	HYPERPHENYLALANINEMIA (HPA): INITIAL: NOT CONCURRENTLY USING PALYNZIQ. RENEWAL: 1) CONTINUES TO BENEFIT FROM TREATMENT, AND 2) NOT CONCURRENTLY USING PALYNZIQ.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SARILUMAB

Products Affected

• KEVZARA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	RHEUMATOID ARTHRITIS (RA): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	RA: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. POLYMYALGIA RHEUMATICA (PMR): 12 MONTHS.
Other Criteria	RA: INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SATRALIZUMAB-MWGE

Products Affected

• ENSPRYNG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NEUROMYELITIS OPTICA SPECTRUM DISORDER (NMOSD): INITIAL: PRESCRIBED BY AN OPHTHALMOLOGIST OR PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	NMOSD: INITIAL: 1) ONE OF THE FOLLOWING CORE CLINICAL CHARACTERISTIC: (A) OPTIC NEURITIS, (B) ACUTE MYELITIS, (C) AREA POSTREMA SYNDROME, (D) ACUTE BRAINSTEM SYNDROME, (E) SYMPTOMATIC NARCOLEPSY OR ACUTE DIENCEPHALIC CLINICAL SYNDROME WITH NMOSD-TYPICAL DIENCEPHALIC MRI LESIONS, OR (F) SYMPTOMATIC CEREBRAL SYNDROME WITH NMOSD-TYPICAL BRAIN LESIONS, AND 2) NO CONCURRENT USE WITH RITUXIMAB, INEBILIZUMAB, OR ECULIZUMAB. RENEWAL: 1) REDUCTION IN RELAPSE FREQUENCY FROM BASELINE, AND 2) NO CONCURRENT USE WITH RITUXIMAB, INEBILIZUMAB, OR ECULIZUMAB.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SEBELIPASE ALFA

Products Affected

• KANUMA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	LYSOSOMAL ACID LIPASE (LAL) DEFICIENCY: INITIAL: DIAGNOSIS CONFIRMED BY 1) PRESENCE OF CLINICAL FEATURES (E.G., HEPATOMEGALY, ELEVATED SERUM TRANSAMINASES, DYSLIPIDEMIA, SPLENOMEGALY), AND 2) ONE OF THE FOLLOWING: (A) BLOOD TEST INDICATING LOW OR ABSENT LEVELS OF LAL ENZYME ACTIVITY, (B) DRIED BLOOD SPOT TEST INDICATING LOW OR ABSENT LAL ENZYME ACTIVITY, OR (C) GENETIC TEST INDICATING THE BI-ALLELIC PRESENCE OF ALTERED LIPA GENE(S).
Age Restrictions	
Prescriber Restrictions	LAL DEFICIENCY: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST, HEPATOLOGIST, GASTROENTEROLOGIST, MEDICAL GENETICIST, LIPIDOLOGIST, OR METABOLIC SPECIALIST.
Coverage Duration	LAL DEFICIENCY: INITIAL: 6 OR 12 MONTHS, SEE OTHER CRITERIA. RENEWAL: 12 MONTHS

PA Criteria	Criteria Details
Other Criteria	LAL DEFICIENCY: INITIAL: RAPIDLY PROGRESSIVE LAL DEFICIENCY PRESENTING WITHIN THE FIRST 6 MONTHS OF LIFE: 12 MONTHS. LAL DEFICIENCY PRESENTING AFTER THE FIRST 6 MONTHS OF LIFE AND NOT CONSIDERED RAPIDLY PROGRESSIVE: 6 MONTHS. RENEWAL: LAL DEFICIENCY PRESENTING AFTER THE FIRST 6 MONTHS OF LIFE AND NOT CONSIDERED RAPIDLY PROGRESSIVE: IMPROVEMENT IN ONE OF THE FOLLOWING CLINICAL PARAMETERS DURING THE PAST 6 MONTHS: (1) A RELATIVE REDUCTION FROM BASELINE IN LDL-C, NON-HDL-C, OR TRIGLYCERIDES, (2) NORMALIZATION OF ASPARTATE AMINOTRANSFERASE (AST) BASED ON AGE- AND GENDER-SPECIFIC NORMAL RANGES, OR (3) DECREASE IN LIVER FAT CONTENT COMPARED TO BASELINE ASSESSED BY ABDOMINAL IMAGING.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SECUKINUMAB

Products Affected

• COSENTYX (2 SYRINGES)

S) • COSENTYX UNOREADY PEN

- COSENTYX PEN (2 PENS)
- COSENTYX SUBCUTANEOUS SYRINGE 75 MG/0.5 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, OR FACE. NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN (CRP) LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI).
Age Restrictions	
Prescriber Restrictions	INITIAL: PSO, HIDRADENITIS SUPPURATIVA (HS): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR A DERMATOLOGIST. ANKYLOSING SPONDYLITIS (AS), NR-AXSPA, ENTHESITIS- RELATED ARTHRITIS (ERA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	INITIAL: HS: 4 MONTHS, ALL OTHER INDICATIONS: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTI- RHEUMATIC DRUG). NR-AXSPA: TRIAL OF OR CONTRAINDICATION TO AN NSAID (NON-STEROIDAL ANTI-INFLAMMATORY DRUG). ERA: TRIAL OF OR CONTRAINDICATION TO ONE NSAID, SULFASALAZINE, OR METHOTREXATE. HS: NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGICS FOR HS OR OTHER IL-17 INHIBITORS FOR ANY INDICATION. RENEWAL: PSO, PSA, AS, NR-AXSPA, ERA: CONTINUES TO BENEFIT FROM THE MEDICATION. HS: 1) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGICS FOR HS OR OTHER IL-17 INHIBITORS FOR ANY INDICATION, AND 2) CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SECUKINUMAB IV

Products Affected

• COSENTYX INTRAVENOUS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN (CRP) LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI).
Age Restrictions	
Prescriber Restrictions	INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR A DERMATOLOGIST. ANKYLOSING SPONDYLITIS (AS), NR- AXSPA: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTI-RHEUMATIC DRUG). NR-AXSPA: TRIAL OF OR CONTRAINDICATION TO AN NSAID (NON-STEROIDAL ANTI-INFLAMMATORY DRUG). RENEWAL: PSA, AS, NR-AXSPA: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SELEXIPAG

Products Affected

- UPTRAVI INTRAVENOUS
- UPTRAVI ORAL TABLET 1,000 MCG, 1,200 MCG, 1,400 MCG, 1,600 MCG, 200 MCG, 400 MCG, 600 MCG, 800 MCG
- UPTRAVI ORAL TABLETS, DOSE PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS.
Age Restrictions	
Prescriber Restrictions	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS
Other Criteria	PAH: INITIAL: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING AGENTS FROM DIFFERENT DRUG CLASSES: 1) FORMULARY VERSION OF AN ORAL ENDOTHELIN RECEPTOR ANTAGONIST, 2) FORMULARY VERSION OF AN ORAL PHOSPHODIESTERASE TYPE-5 INHIBITOR, OR 3) FORMULARY VERSION OF AN ORAL CGMP STIMULATOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SELINEXOR

Products Affected

 XPOVIO ORAL TABLET 100 MG/WEEK (50 MG X 2), 40 MG/WEEK (40 MG X 1), 40MG TWICE WEEK (40 MG X 2), 60 MG/WEEK (60 MG X 1), 60MG TWICE WEEK (120 MG/WEEK), 80 MG/WEEK (40 MG X 2), 80MG TWICE WEEK (160 MG/WEEK)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SELPERCATINIB

Products Affected

• RETEVMO ORAL CAPSULE 40 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SELUMETINIB

Products Affected

 KOSELUGO ORAL CAPSULE 10 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SILDENAFIL IV

Products Affected

• sildenafil (pulm.hypertension) intravenous

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS.
Age Restrictions	
Prescriber Restrictions	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	PAH: INITIAL/RENEWAL: NOT CONCURRENTLY OR INTERMITTENTLY TAKING ORAL ERECTILE DYSFUNCTION AGENTS (E.G. CIALIS, VIAGRA), ANY ORGANIC NITRATES IN ANY FORM, OR GUANYLATE CYCLASE STIMULATORS.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

SILDENAFIL TABLET

Products Affected

• sildenafil (pulm.hypertension) oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: AGES 18 YEARS OR OLDER: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS. AGES 1 TO 17 YEARS: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PAP GREATER THAN 20 MMHG, 2) PCWP OF 15 MMHG OR LESS, AND 3) PVR OF 3 WOOD UNITS OR GREATER.
Age Restrictions	
Prescriber Restrictions	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	PAH: INITIAL/RENEWAL: NOT CONCURRENTLY OR INTERMITTENTLY TAKING ORAL ERECTILE DYSFUNCTION AGENTS (E.G. CIALIS, VIAGRA), ANY ORGANIC NITRATES IN ANY FORM, OR GUANYLATE CYCLASE STIMULATORS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
Indications	All Medically-accepted Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

SIPONIMOD

Products Affected

• MAYZENT ORAL TABLET 0.25 MG, 1 • MAYZENT STARTER(FOR 2MG MG, 2 MG

MAINT)

• MAYZENT STARTER(FOR 1MG MAINT)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	MULTIPLE SCLEROSIS: RENEWAL: 1) DEMONSTRATION OF CLINICAL BENEFIT COMPARED TO PRE-TREATMENT BASELINE AND 2) DOES NOT HAVE LYMPHOPENIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SIROLIMUS PROTEIN-BOUND

Products Affected

• FYARRO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SODIUM OXYBATE-XYREM

Products Affected

• sodium oxybate

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: ALL INDICATIONS: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR SPECIALIST IN SLEEP MEDICINE
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: EXCESSIVE DAYTIME SLEEPINESS (EDS) IN NARCOLEPSY: 1) NOT CURRENTLY TAKING A SEDATIVE HYPNOTIC AGENT, 2) FOR PATIENTS 18 YEARS OR OLDER: TRIAL OF OR CONTRAINDICATION TO A FORMULARY VERSION OF MODAFINIL, ARMODAFINIL, OR SOLRIAMFETOL AND ONE OTHER GENERIC STIMULANT INDICATED FOR EDS IN NARCOLEPSY, AND 3) FOR PATIENTS 7 TO 17 YEARS OF AGE: TRIAL OF OR CONTRAINDICATION TO ONE GENERIC STIMULANT INDICATED FOR EDS IN NARCOLEPSY. CATAPLEXY IN NARCOLEPSY: NOT CURRENTLY TAKING A SEDATIVE HYPNOTIC AGENT. RENEWAL (ALL INDICATIONS): 1) SUSTAINED IMPROVEMENT OF SYMPTOMS COMPARED TO BASELINE, AND 2) NOT CURRENTLY TAKING A SEDATIVE HYPNOTIC AGENT.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

SODIUM PHENYLBUTYRATE TABLETS

Products Affected

• sodium phenylbutyrate oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	UREA CYCLE DISORDER (UCD): INITIAL: UCD IS CONFIRMED VIA ENZYMATIC, BIOCHEMICAL OR GENETIC TESTING.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	UCD: RENEWAL: CLINICAL BENEFIT FROM BASELINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SOFOSBUVIR/VELPATASVIR

Products Affected

- EPCLUSA ORAL PELLETS IN PACKET 150-37.5 MG, 200-50 MG
- EPCLUSA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HCV RNA LEVEL WITHIN PAST 6 MONTHS.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
Other Criteria	1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, 2) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING MEDICATIONS: AMIODARONE, CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, HIV REGIMEN THAT CONTAINS EFAVIRENZ, ROSUVASTATIN AT DOSES ABOVE 10MG, TIPRANAVIR/RITONAVIR, TOPOTECAN, SOVALDI (AS A SINGLE AGENT), HARVONI, ZEPATIER, MAVYRET, OR VOSEVI, AND 3) PATIENTS WITH DECOMPENSATED CIRRHOSIS REQUIRE CONCURRENT RIBAVIRIN UNLESS RIBAVIRIN INELIGIBLE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SOFOSBUVIR/VELPATASVIR/VOXILAPREVIR

Products Affected

• VOSEVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HCV RNA LEVEL WITHIN PAST 6 MONTHS
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
Other Criteria	1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, 2) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING MEDICATIONS: AMIODARONE, CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, CYCLOSPORINE, PITAVASTATIN, PRAVASTATIN (DOSES ABOVE 40MG), ROSUVASTATIN, METHOTREXATE, MITOXANTRONE, IMATINIB, IRINOTECAN, LAPATINIB, SULFASALAZINE, TOPOTECAN, OR HIV REGIMEN THAT CONTAINS EFAVIRENZ, ATAZANAVIR, LOPINAVIR, TIPRANAVIR/RITONAVIR, SOVALDI (AS A SINGLE AGENT), EPCLUSA, HARVONI, ZEPATIER, OR MAVYRET, AND 3) DOES NOT HAVE MODERATE OR SEVERE HEPATIC IMPAIRMENT (CHILD-PUGH B OR C).
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

SOLRIAMFETOL

Products Affected

• SUNOSI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: EXCESSIVE DAYTIME SLEEPINESS (EDS) IN NARCOLEPSY: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR SPECIALIST IN SLEEP MEDICINE.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: EDS IN NARCOLEPSY: TRIED THE FORMULARY VERSION OF MODAFINIL OR ARMODAFINIL, AND ONE OTHER GENERIC STIMULANT INDICATED FOR EDS IN NARCOLEPSY. EDS IN OBSTRUCTIVE SLEEP APNEA (OSA): TRIED THE FORMULARY VERSION OF MODAFINIL OR ARMODAFINIL. RENEWAL: EDS IN NARCOLEPSY OR OSA: SUSTAINED IMPROVEMENT OF SYMPTOMS COMPARED TO BASELINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SOMATROPIN - NORDITROPIN

Products Affected

• NORDITROPIN FLEXPRO

PA Criteria	Criteria Details
Exclusion Criteria	INITIAL/RENEWAL: ATHLETIC ENHANCEMENT, ANTI- AGING PURPOSES.
Required Medical Information	INITIAL: PEDIATRIC GROWTH HORMONE DEFICIENCY (GHD), IDIOPATHIC SHORT STATURE (ISS), SMALL FOR GESTATIONAL AGE (SGA), TURNER SYNDROME (TS), NOONAN SYNDROME: HEIGHT AT LEAST 2 STANDARD DEVIATIONS (SD) BELOW THE MEAN HEIGHT FOR NORMAL CHILDREN OF THE SAME AGE AND GENDER. PRADER WILLI SYNDROME (PWS): CONFIRMED GENETIC DIAGNOSIS.
Age Restrictions	
Prescriber Restrictions	INITIAL/RENEWAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: ADULT GHD: GROWTH HORMONE DEFICIENCY ALONE OR ASSOCIATED WITH MULTIPLE HORMONE DEFICIENCIES (HYPOPITUITARISM), AS A RESULT OF PITUITARY DISEASES, HYPOTHALAMIC DISEASE, SURGERY, RADIATION THERAPY, TRAUMA, OR CONTINUATION OF THERAPY FROM CHILDHOOD ONSET GROWTH HORMONE DEFICIENCY. PEDIATRIC GHD, ISS, SGA, TS, NOONAN SYNDROME: OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF THE WRIST AND HAND. RENEWAL: PEDIATRIC GHD: 1) IMPROVEMENT WHILE ON THERAPY (I.E., INCREASED HEIGHT OR INCREASED GROWTH VELOCITY), AND 2) OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF THE WRIST AND HAND OR PATIENT HAS NOT COMPLETED PREPUBERTALGROWTH. ISS, SGA, TS, NOONAN SYNDROME: 1) IMPROVEMENT WHILE ON THERAPY (I.E., INCREASED HEIGHT OR INCREASED GROWTH VELOCITY), AND 2) OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF THE WRIST AND HAND. PWS: IMPROVEMENT IN BODY COMPOSITION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SOMATROPIN - SEROSTIM

Products Affected

• SEROSTIM SUBCUTANEOUS RECON SOLN 4 MG, 5 MG, 6 MG

PA Criteria	Criteria Details
Exclusion Criteria	INITIAL/RENEWAL: ATHLETIC ENHANCEMENT, ANTI- AGING PURPOSES
Required Medical Information	INITIAL: HIV/WASTING: MEETS ONE OF THE FOLLOWING CRITERIA FOR WEIGHT LOSS: 1) 10% UNINTENTIONAL WEIGHT LOSS OVER 12 MONTHS, 2) 7.5% UNINTENTIONAL WEIGHT LOSS OVER 6 MONTHS, 3) 5% BODY CELL MASS (BCM) LOSS WITHIN 6 MONTHS, 4) BODY CELL MASS (BCM) LESS THAN 35% (MEN) OF TOTAL BODY WEIGHT AND BODY MASS INDEX (BMI) LESS THAN 27 KG PER METER SQUARED, 5) BCM LESS THAN 23% (WOMEN) OF TOTAL BODY WEIGHT AND A BODY MASS INDEX (BMI) LESS THAN 27 KG PER METER SQUARED, OR 6) BMI LESS THAN 18.5 KG PER METER SQUARED.
Age Restrictions	
Prescriber Restrictions	HIV/WASTING: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST, NUTRITIONAL SUPPORT SPECIALIST, OR INFECTIOUS DISEASE SPECIALIST.
Coverage Duration	INITIAL/RENEWAL: 3 MONTHS.
Other Criteria	HIV/WASTING: INITIAL: 1) CURRENTLY ON HIV ANTIRETROVIRAL THERAPY, AND 2) INADEQUATE RESPONSE TO ONE PREVIOUS THERAPY (E.G., MEGACE, APPETITE STIMULANTS, ANABOLIC STEROIDS). RENEWAL: 1) CURRENTLY ON HIV ANTIRETROVIRAL THERAPY, AND 2) CLINICAL BENEFIT IN MUSCLE MASS AND WEIGHT.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

SONIDEGIB

Products Affected

• ODOMZO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	LOCALLY ADVANCED BASAL CELL CARCINOMA (BCC): BASELINE SERUM CREATINE KINASE (CK) AND SERUM CREATININE LEVELS
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SORAFENIB

Products Affected

• sorafenib

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SOTORASIB

Products Affected

• LUMAKRAS ORAL TABLET 120 MG, 320 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

STIRIPENTOL

Products Affected

- DIACOMIT ORAL CAPSULE 250 MG, 500 MG
- DIACOMIT ORAL POWDER IN PACKET 250 MG, 500 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	DRAVET SYNDROME: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SUNITINIB

Products Affected

• *sunitinib malate*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	GASTROINTESTINAL STROMAL TUMORS (GIST): TRIAL OF OR CONTRAINDICATION TO IMATINIB (GLEEVEC).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TADALAFIL - ADCIRCA, ALYQ

Products Affected

- alyq
- tadalafil (pulm. hypertension)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS.
Age Restrictions	
Prescriber Restrictions	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	PAH: INITIAL/RENEWAL: NOT CONCURRENTLY OR INTERMITTENTLY TAKING ORAL ERECTILE DYSFUNCTION AGENTS (E.G. CIALIS, VIAGRA), ANY ORGANIC NITRATES IN ANY FORM, OR GUANYLATE CYCLASE STIMULATORS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

TADALAFIL-CIALIS

Products Affected

• tadalafil oral tablet 2.5 mg, 5 mg

PA Criteria	Criteria Details
Exclusion Criteria	ERECTILE DYSFUNCTION WITHOUT DIAGNOSIS OF BENIGN PROSTATIC HYPERPLASIA (BPH).
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	BPH: 1) TRIAL OF ONE ALPHA BLOCKER (E.G., DOXAZOSIN, TERAZOSIN, TAMSULOSIN, ALFUZOSIN), AND 2) TRIAL OF ONE 5-ALPHA-REDUCTASE INHIBITOR (E.G., FINASTERIDE, DUTASTERIDE). APPLIES TO 2.5MG AND 5MG STRENGTHS ONLY. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TALAZOPARIB

Products Affected

• TALZENNA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ADVANCED OR METASTATIC BREAST CANCER: PATIENT HAS BEEN TREATED WITH CHEMOTHERAPY IN THE NEOADJUVANT, ADJUVANT, OR METASTATIC SETTING. PATIENTS WITH HORMONE RECEPTOR (HR)-POSITIVE BREAST CANCER MUST HAVE ADDITIONAL PRIOR TREATMENT WITH ENDOCRINE THERAPY OR BE CONSIDERED INAPPROPRIATE FOR ENDOCRINE THERAPY. METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TALIMOGENE

Products Affected

• IMLYGIC INJECTION SUSPENSION 10EXP6 (1 MILLION) PFU/ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	UNRESECTABLE MELANOMA: 1) IMLYGIC TO BE INJECTED INTO CUTANEOUS, SUBCUTANEOUS, AND/OR NODAL LESIONS THAT ARE VISIBLE, PALPABLE, OR DETECTABLE BY ULTRASOUND GUIDANCE, 2) NOT CURRENTLY RECEIVING IMMUNOSUPPRESSIVE THERAPY, 3) NO HISTORY OF PRIMARY OR ACQUIRED IMMUNODEFICIENT STATES, LEUKEMIA, LYMPHOMA, OR AIDS, AND 4) NO CONCURRENT USE WITH PEMBROLIZUMAB, NIVOLUMAB, IPILIMUMAB, DABRAFENIB, TRAMETINIB, VEMURAFENIB, INTERLEUKIN-2, INTERFERON, DACARBAZINE, TEMOZOLOMIDE, PACLITAXEL, CARBOPLATIN, IMATINIB, MELPHALAN, IMIQUIMOD, OR RADIATION THERAPY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TALQUETAMAB-TGVS

Products Affected

• TALVEY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TASIMELTEON

Products Affected

- HETLIOZ LQ
- tasimelteon

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	LIFETIME
Other Criteria	NON-24 HOUR SLEEP-WAKE DISORDER: PATIENT IS LIGHT-INSENSITIVE OR HAS TOTAL BLINDNESS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TAZEMETOSTAT

Products Affected

• TAZVERIK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TBO-FILGRASTIM

Products Affected

• GRANIX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NON-MYELOID MALIGNANCIES: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	NON-MYELOID MALIGNANCIES: TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: NIVESTYM.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TEBENTAFUSP-TEBN

Products Affected

• KIMMTRAK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TECLISTAMAB-CQYV

Products Affected

• TECVAYLI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TEDUGLUTIDE

Products Affected

• GATTEX 30-VIAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	SHORT BOWEL SYNDROME (SBS): INITIAL/RENEWAL: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS
Other Criteria	SBS: INITIAL: PATIENT IS DEPENDENT ON INTRAVENOUS PARENTERAL NUTRITION DEFINED AS REQUIRING PARENTERAL NUTRITION AT LEAST THREE TIMES PER WEEK. RENEWAL: ACHIEVED OR MAINTAINED A DECREASED NEED FOR PARENTERAL SUPPORT COMPARED TO BASELINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TELOTRISTAT

Products Affected

• XERMELO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	CARCINOID SYNDROME DIARRHEA: PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST OR GASTROENTEROLOGIST
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TEPOTINIB

Products Affected

• ТЕРМЕТКО

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TEPROTUMUMAB-TRBW

Products Affected

• TEPEZZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TERIFLUNOMIDE

Products Affected

• teriflunomide

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TESAMORELIN

Products Affected

• EGRIFTA SV

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TESTOSTERONE

Products Affected

- testosterone transdermal gel in metereddose pump 12.5 mgl 1.25 gram (1%), 20.25
 mgl1.25 gram (1.62%)
- testosterone transdermal gel in packet 1 %

(25 mg/2.5gram), 1 % (50 mg/5 gram) testosterone transdermal solution in metered pump wlapp

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	MALE HYPOGONADISM: INITIAL: CONFIRMED BY: 1) AT LEAST TWO TOTAL SERUM TESTOSTERONE LEVELS OF LESS THAN 300 NG/DL TAKEN ON SEPARATE OCCASIONS, OR 2) FREE SERUM TESTOSTERONE LEVEL OF LESS THAN 5 NG/DL.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS
Other Criteria	MALE HYPOGONADISM: INITIAL: 1) 40 YEARS OR OLDER: PROSTATE SPECIFIC ANTIGEN HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING. RENEWAL: 1) 40 YEARS OR OLDER: PROSTATE SPECIFIC ANTIGEN HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING, AND 2) IMPROVED SYMPTOMS COMPARED TO BASELINE AND TOLERANCE TO TREATMENT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TESTOSTERONE CYPIONATE

Products Affected

• testosterone cypionate

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	MALE HYPOGONADISM: INITIAL: CONFIRMED BY: 1) AT LEAST TWO TOTAL SERUM TESTOSTERONE LEVELS OF LESS THAN 300 NG/DL TAKEN ON SEPARATE OCCASIONS, OR 2) FREE SERUM TESTOSTERONE LEVEL OF LESS THAN 5 NG/DL.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS
Other Criteria	MALE HYPOGONADISM: INITIAL: 1) 40 YEARS OR OLDER: PROSTATE SPECIFIC ANTIGEN HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING. RENEWAL: 1) 40 YEARS OR OLDER: PROSTATE SPECIFIC ANTIGEN HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING, AND 2) IMPROVED SYMPTOMS COMPARED TO BASELINE AND TOLERANCE TO TREATMENT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TESTOSTERONE ENANTHATE

- *testosterone enanthate*
- XYOSTED

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	MALE HYPOGONADISM: INITIAL: CONFIRMED BY: 1) AT LEAST TWO TOTAL SERUM TESTOSTERONE LEVELS OF LESS THAN 300 NG/DL TAKEN ON SEPARATE OCCASIONS, OR 2) FREE SERUM TESTOSTERONE LEVEL OF LESS THAN 5 NG/DL.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	MALE HYPOGONADISM: INITIAL/RENEWAL: 12 MO. ALL OTHER INDICATIONS: LIFETIME OF MEMBERSHIP IN PLAN.
Other Criteria	MALE HYPOGONADISM: INITIAL: 1) 40 YEARS OR OLDER: PROSTATE SPECIFIC ANTIGEN HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING. RENEWAL: 1) 40 YEARS OR OLDER: PROSTATE SPECIFIC ANTIGEN HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING, AND 2) IMPROVED SYMPTOMS COMPARED TO BASELINE AND TOLERANCE TO TREATMENT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TETRABENAZINE

Products Affected

• *tetrabenazine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	HUNTINGTONS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR MOVEMENT DISORDER SPECIALIST
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TEZACAFTOR/IVACAFTOR

Products Affected

• SYMDEKO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	CYSTIC FIBROSIS (CF): INITIAL: CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CYSTIC FIBROSIS.
Age Restrictions	
Prescriber Restrictions	CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CYSTIC FIBROSIS EXPERT
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: LIFETIME
Other Criteria	CF: RENEWAL: MAINTAINED, IMPROVED, OR DEMONSTRATED LESS THAN EXPECTED DECLINE IN FEV1 OR BODY MASS INDEX (BMI), OR REDUCTION IN NUMBER OF PULMONARY EXACERBATIONS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

THALIDOMIDE

Products Affected

• THALOMID

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TILDRAKIZUMAB-ASMN

Products Affected

• ILUMYA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	PLAQUE PSORIASIS (PSO): INITIAL: PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, OR FACE.
Age Restrictions	
Prescriber Restrictions	PSO: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	PSO: INITIAL: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, COSENTYX, STELARA, ENBREL, SKYRIZI, TREMFYA. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TISOTUMAB VEDOTIN-TFTV

Products Affected

• TIVDAK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TIVOZANIB

Products Affected

• FOTIVDA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TOCILIZUMAB IV

Products Affected

• ACTEMRA

PA Criteria	Criteria Details
Exclusion Criteria	CORONAVIRUS DISEASE 2019 (COVID-19) IN HOSPITALIZED ADULTS
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS (SJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, OR IMMUNOLOGIST.
Coverage Duration	INITIAL: RA, PJIA, SJIA, GCA: 6 MONTHS. CRS: 1 MONTH. RENEWAL: RA, PJIA, SJIA, GCA: 12 MONTHS.
Other Criteria	INITIAL: RA: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PJIA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ IR. SJIA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG). RENEWAL: RA, PJIA, SJIA: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

TOCILIZUMAB SQ

- ACTEMRA
- ACTEMRA ACTPEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS (SJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, OR IMMUNOLOGIST. SYSTEMIC SCLEROSIS-ASSOCIATED INTERSTITIAL LUNG DISEASE (SSC-ILD): PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR RHEUMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: RA: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PJIA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ IR. SJIA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG). SSC-ILD: DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., CONNECTIVE TISSUE DISEASE, DRUG TOXICITY, ASBESTOS OR BERYLLIUM EXPOSURE, HYPERSENSITIVITY PNEUMONITIS). RENEWAL: RA, PJIA, SJIA: CONTINUES TO BENEFIT FROM THE MEDICATION. SSC-ILD: CLINICAL MEANINGFUL IMPROVEMENT OR MAINTENANCE IN ANNUAL RATE OF DECLINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TOFACITINIB

- XELJANZ
- XELJANZ XR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS), POLYARTICULAR COURSE JUVENILE IDIOPATHIC ARTHRITIS (PCJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE- MODIFYING ANTIRHEUMATIC DRUG) - IF A PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PSA, PCJIA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD. UC: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPURINE, METHOTREXATE, OR MESALAMINE. RENEWAL: RA, PSA, AS, PCJIA: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

TOLVAPTAN

- JYNARQUE ORAL TABLET
- JYNARQUE ORAL TABLETS, SEQUENTIAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	AUTOSOMAL DOMINANT POLYCYSTIC KIDNEY DISEASE (ADPKD): INITIAL: 1) CONFIRMED POLYCYSTIC KIDNEY DISEASE VIA CT, MRI, OR ULTRASOUND, AND 2) GENETIC TESTING FOR CAUSATIVE MUTATIONS OR FAMILY HISTORY OF CONFIRMED POLYCYSTIC KIDNEY DISEASE IN ONE OR BOTH PARENTS.
Age Restrictions	
Prescriber Restrictions	ADPKD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEPHROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	ADPKD: INITIAL: DOES NOT HAVE ESRD (I.E., RECEIVING DIALYSIS OR HAS UNDERGONE RENAL TRANSPLANT). RENEWAL: HAS NOT PROGRESSED TO ESRD/DIALYSIS OR TRANSPLANT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TOPICAL TRETINOIN

- ALTRENO
- tretinoin

PA Criteria	Criteria Details
Exclusion Criteria	COSMETIC INDICATIONS SUCH AS WRINKLES, PHOTOAGING, MELASMA.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ACNE VULGARIS: BRAND TOPICAL TRETINOIN REQUIRES TRIAL OF OR CONTRAINDICATION TO A GENERIC TOPICAL TRETINOIN PRODUCT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TORIPALIMAB-TPZI

Products Affected

• LOQTORZI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	NASOPHARYNGEAL CARCINOMA (NPC): FIRST LINE TREATMENT: 24 MOS, PREVIOUSLY TREATED: LIFETIME.
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TRAMETINIB

Products Affected

• MEKINIST ORAL TABLET 0.5 MG, 2 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TRAMETINIB SOLUTION

Products Affected

• MEKINIST ORAL RECON SOLN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	UNABLE TO SWALLOW MEKINIST TABLETS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TRASTUZUMAB HYALURONIDASE

Products Affected

• HERCEPTIN HYLECTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ADJUVANT BREAST CANCER, METASTATIC BREAST CANCER: TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREFERRED AGENTS: HERZUMA, OGIVRI, ONTRUZANT, TRAZIMERA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TRASTUZUMAB-ANNS

Products Affected

• KANJINTI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ADJUVANT BREAST CANCER, METASTATIC BREAST CANCER: TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREFERRED AGENTS: HERZUMA, OGIVRI, ONTRUZANT, TRAZIMERA. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TRASTUZUMAB-DKST

Products Affected

• OGIVRI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TRASTUZUMAB-DTTB

Products Affected

• ONTRUZANT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TRASTUZUMAB-PKRB

Products Affected

• HERZUMA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TRASTUZUMAB-QYYP

Products Affected

• TRAZIMERA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TREMELIMUMAB-ACTL

Products Affected

• IMJUDO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	UHCC: 30 DAYS. METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC): 5 MONTHS.
Other Criteria	UNRESECTABLE HEPATOCELLULAR CARCINOMA (UHCC): HAS NOT RECEIVED PRIOR TREATMENT WITH IMJUDO.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TREPROSTINIL INHALED

Products Affected

• TYVASO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PULMONARY ARTERIAL HYPERTENSION (PAH), PULMONARY HYPERTENSION-INTERSTITIAL LUNG DISEASE (PH-ILD): DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS.
Age Restrictions	
Prescriber Restrictions	INITIAL: PAH, PH-ILD: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
Coverage Duration	INITIAL: PAH: 12 MONTHS, PH-ILD: 6 MONTHS. RENEWAL: PAH, PH-ILD: 12 MONTHS.
Other Criteria	INITIAL: PAH: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING AGENTS FROM DIFFERENT DRUG CLASSES: 1) FORMULARY VERSION OF AN ORAL ENDOTHELIN RECEPTOR ANTAGONIST, 2) FORMULARY VERSION OF AN ORAL PHOSPHODIESTERASE TYPE-5 INHIBITOR, 3) FORMULARY VERSION OF AN ORAL CGMP STIMULATOR, 4) FORMULARY VERSION OF AN IV/SQ PROSTACYCLIN. THIS DRUG MAY BE COVERED UNDER MEDICARE PART B OR D DEPENDING UPON THE CIRCUMSTANCES.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

TREPROSTINIL INJECTABLE

Products Affected

• treprostinil sodium

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS.
Age Restrictions	
Prescriber Restrictions	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS
Other Criteria	PAH: INITIAL: 1) CONTINUATION OF THERAPY FROM HOSPITAL DISCHARGE, 2) NEW START AND PHYSICIAN INDICATED PATIENT IS INTERMEDIATE OR HIGH RISK, OR 3) NEW START AND TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING AGENTS FROM DIFFERENT DRUG CLASSES: (A) FORMULARY VERSION OF AN ORAL ENDOTHELIN RECEPTOR ANTAGONIST, (B) FORMULARY VERSION OF AN ORAL PHOSPHODIESTERASE TYPE-5 INHIBITOR, OR (C) FORMULARY VERSION OF AN ORAL CGMP STIMULATOR. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

TRIENTINE CAPSULE

Products Affected

• trientine oral capsule 250 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	WILSONS DISEASE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEPATOLOGIST OR GASTROENTEROLOGIST.
Coverage Duration	INITIAL: 12 MONTHS, RENEWAL: LIFETIME.
Other Criteria	WILSONS DISEASE: INITIAL: 1) LEIPZIG SCORE OF 4 OR GREATER, AND 2) TRIAL OF OR CONTRAINDICATION TO FORMULARY VERSION OF PENICILLAMINE TABLET. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TRIFLURIDINE/TIPIRACIL

Products Affected

• LONSURF ORAL TABLET 15-6.14 MG, 20-8.19 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TRIPTORELIN-TRELSTAR

Products Affected

 TRELSTAR INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS.
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TRIPTORELIN-TRIPTODUR

Products Affected

• TRIPTODUR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	CENTRAL PRECOCIOUS PUBERTY (CPP): INITIAL: FEMALES: ELEVATED LEVELS OF FOLLICLE- STIMULATING HORMONE (FSH) GREATER THAN 4.0 MIU/ML AND LUTEINIZING HORMONE (LH) LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS. MALES: ELEVATED LEVELS OF FSH GREATER THAN 5.0 MIU/ML AND LH LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS.
Age Restrictions	
Prescriber Restrictions	CPP: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	CPP: INITIAL: FEMALES: 1) YOUNGER THAN 8 YEARS OF AGE AT ONSET OF CPP, AND 2) AT TANNER STAGE 2 OR ABOVE FOR BREAST DEVELOPMENT AND PUBIC HAIR GROWTH. MALES: 1) YOUNGER THAN 9 YEARS OF AGE AT ONSET OF CPP, AND 2) AT TANNER STAGE 2 OR ABOVE FOR GENITAL DEVELOPMENT AND PUBIC HAIR GROWTH. RENEWAL: 1) TANNER STAGING AT INITIAL DIAGNOSIS HAS STABILIZED OR REGRESSED DURING THREE SEPARATE MEDICAL VISITS IN THE PREVIOUS YEAR, AND 2) HAS NOT REACHED ACTUAL AGE WHICH CORRESPONDS TO CURRENT PUBERTAL AGE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

TUCATINIB

Products Affected

 TUKYSA ORAL TABLET 150 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

UBROGEPANT

Products Affected

• UBRELVY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	ACUTE MIGRAINE TREATMENT: INITIAL: TRIAL OF OR CONTRAINDICATION TO ONE TRIPTAN (E.G., SUMATRIPTAN, RIZATRIPTAN). RENEWAL: 1) IMPROVEMENT FROM BASELINE IN A VALIDATED ACUTE TREATMENT PATIENT-REPORTED OUTCOME QUESTIONNAIRE, OR 2) THERAPY WORKS CONSISTENTLY IN MAJORITY OF MIGRAINE ATTACKS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

UPADACITINIB

Products Affected

• RINVOQ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR- AXSPA): 1) C-REACTIVE PROTEIN LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI).
Age Restrictions	
Prescriber Restrictions	INITIAL: RA, AS, NR-AXSPA: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSA: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. AD: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST, ALLERGIST, OR IMMUNOLOGIST. UC, CD: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE- MODIFYING ANTIRHEUMATIC DRUG) - IF A PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD. ATOPIC DERMATITIS: 1) ATOPIC DERMATITIS COVERING AT LEAST 10 PERCENT OF BODY SURFACE AREA OR ATOPIC DERMATITIS AFFECTING THE FACE, HEAD, NECK, HANDS, FEET, GROIN, OR INTERTRIGINOUS AREAS, 2) INTRACTABLE PRURITUS OR CRACKING/OOZING/BLEEDING OF AFFECTED SKIN, 3) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING: TOPICAL CORTICOSTEROID, TOPICAL CALCINEURIN INHIBITOR, TOPICAL PDE4 INHIBITOR, OR TOPICAL JAK INHIBITOR, AND 4) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGIC/JAK INHIBITOR FOR THE TREATMENT OF ATOPIC DERMATITIS. UC: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPURINE, METHOTREXATE, OR MESALAMINE. NR-AXSPA: TRIAL OF OR CONTRAINDICATION TO AN NSAID (NON-STEROIDAL ANTI-INFLAMMATORY DRUG). CD: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPURINE, METHOTREXATE, OR MESALAMINE. RENEWAL: RA, PSA, AS, NR-AXSPA: CONTINUES TO BENEFIT FROM THE MEDICATION. ATOPIC DERMATITIS: 1) IMPROVEMENT WHILE ON THERAPY, AND 2) NO CONCURRENT USE WITH OTHER
Indications	SYSTEMIC BIOLOGIC/JAK INHIBITOR FOR THE TREATMENT OF ATOPIC DERMATITIS.All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

USTEKINUMAB

Products Affected

• STELARA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, OR FACE.
Age Restrictions	
Prescriber Restrictions	INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG). PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. CD, UC: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPURINE, METHOTREXATE, OR MESALAMINE. RENEWAL: PSA, PSO: CONTINUES TO BENEFIT FROM THE MEDICATION.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

USTEKINUMAB IV

Products Affected

• STELARA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	2 MONTHS
Other Criteria	CD, UC: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPURINE, METHOTREXATE, OR MESALAMINE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

VALBENAZINE

Products Affected

- INGREZZA
- INGREZZA INITIATION PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	TARDIVE DYSKINESIA (TD): PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR MOVEMENT DISORDER SPECIALIST.
Coverage Duration	12 MONTHS
Other Criteria	TD: 1) PRIOR HISTORY OF USING AGENTS THAT CAUSE TARDIVE DYSKINESIA, AND 2) TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: AUSTEDO.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

VANDETANIB

Products Affected

• CAPRELSA ORAL TABLET 100 MG, 300 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	CURRENTLY STABLE ON CAPRELSA REQUIRES NO EXTRA CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

VEMURAFENIB

Products Affected

• ZELBORAF

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	MELANOMA: ZELBORAF WILL BE USED ALONE OR IN COMBINATION WITH COTELLIC
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

VENETOCLAX

Products Affected

- VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG
- VENCLEXTA STARTING PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

VERICIGUAT

Products Affected

• VERQUVO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL/RENEWAL:12 MONTHS.
Other Criteria	HEART FAILURE (HF): INITIAL: TRIAL OF OR CONTRAINDICATION TO 1) ONE AGENT FROM ANY OF THE FOLLOWING STANDARD OF CARE CLASSES: A) ACE INHIBITOR, ARB, OR ARNI, B) BETA BLOCKER (I.E., BISOPROLOL, CARVEDILOL, METOPROLOL SUCCINATE), OR C) ALDOSTERONE ANTAGONIST (I.E., SPIRONOLACTONE, EPLERENONE), AND 2) ONE PREFERRED SGLT-2 INHIBITOR (E.G., FARXIGA, XIGDUO XR, JARDIANCE). INITIAL/RENEWAL: NOT CONCURRENTLY TAKING LONG-ACTING NITRATES OR NITRIC OXIDE DONORS, RIOCIGUAT, OR PDE-5 INHIBITORS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

VESTRONIDASE ALFA VJBK

Products Affected

• MEPSEVII

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	MUCOPOLYSACCHARIDOSIS VII (MPS VII): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN GENETIC OR METABOLIC DISORDERS.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	MPS VII: INITIAL: 1) HAS NOT UNDERGONE SUCCESSFUL BONE MARROW OR STEM CELL TREATMENT FOR MPS VII, 2) LIMITATION IN MOBILITY, BUT REMAINS SUFFICIENTLY AMBULATORY, AND 3) DIAGNOSIS CONFIRMED BY: (A) URINARY GAG (GLYCOSAMINOGLYCAN) LEVEL IS GREATER THAN THREE TIMES THE UPPER LEVEL OF NORMAL BASED ON THE LABORATORY ASSAY, (B) BETA-GLUCURONIDASE ENZYME ACTIVITY DEFICIENCY OR GENETIC TESTING, AND (C) ONE OF THE FOLLOWING CLINICAL SIGNS OF MPS VII: ENLARGED LIVER AND SPLEEN, JOINT LIMITATIONS, AIRWAY OBSTRUCTIONS OR PULMONARY DYSFUNCTION. RENEWAL: IMPROVED, MAINTAINED, OR DEMONSTRATED A LESS THAN EXPECTED DECLINE IN AMBULATORY ABILITY FROM BASELINE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

VIGABATRIN

Products Affected

- vigabatrin
- vigadrone
- vigpoder

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	REFRACTORY COMPLEX PARTIAL SEIZURES (CPS), INFANTILE SPASMS: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	CPS: TRIAL OF OR CONTRAINDICATION TO TWO ANTIEPILEPTIC AGENTS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

VISMODEGIB

Products Affected

• ERIVEDGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

VORICONAZOLE SUSPENSION

Products Affected

• voriconazole oral suspension for reconstitution

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	CANDIDA INFECTIONS: 3 MOS. ALL OTHER INDICATIONS: 6 MOS.
Other Criteria	CANDIDA INFECTIONS: TRIAL OF OR CONTRAINDICATION TO FLUCONAZOLE. ALL INDICATIONS: INABILITY TO SWALLOW TABLETS OR AN INDICATION FOR ESOPHAGEAL CANDIDIASIS. CONTINUATION OF THERAPY AFTER HOSPITAL DISCHARGE REQUIRES NO EXTRA CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ZANUBRUTINIB

Products Affected

• BRUKINSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ZURANOLONE

Products Affected

• ZURZUVAE ORAL CAPSULE 20 MG, 25 MG, 30 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	14 DAYS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

INDEX

abiraterone	6
ACTEMRA4	15, 417
ACTEMRA ACTPEN	417
ACTHAR	73
ACTIMMUNE	213
ADAKVEO	75
ADEMPAS	
AJOVY AUTOINJECTOR	151
AJOVY SYRINGE	151
AKEEGA	
ALECENSA	
ALTRENO	
ALUNBRIG ORAL TABLET 180 M	
30 MG, 90 MG	55
ALUNBRIG ORAL TABLETS, DO	SE
PACK	
alyq	390
amabelz	178
ambrisentan	17
apomorphine	21
armodafinil	266
ascomp with codeine	172
AUGTYRO	333
AUSTEDO ORAL TABLET 12 MG	i, 6
MG, 9 MG	94
AUSTEDO XR ORAL TABLET	
EXTENDED RELEASE 24 HR 12	
MG, 24 MG, 6 MG	94
AUSTEDO XR TITRATION	
KT(WK1-4)	94
AVONEX INTRAMUSCULAR PE	N
INJECTOR KIT	210
AVONEX INTRAMUSCULAR	
SYRINGE KIT	210
AVSOLA	
AYVAKIT	31
BALVERSA ORAL TABLET 3 MG	i, 4
MG, 5 MG	131
bendamustine intravenous recon soln	
BENDAMUSTINE INTRAVENOU	
SOLUTION	
BENDEKA	
BENLYSTA INTRAVENOUS	40

BENLYSTA SUBCUTANEOUS
BESREMI
<i>betaine</i> 46
BETASERON SUBCUTANEOUS KIT
bexarotene
BEYFORTUS
<i>bortezomib injection</i>
bosentan
BOSULIF ORAL CAPSULE 100 MG,
50 MG 54
BOSULIF ORAL TABLET 100 MG,
400 MG, 500 MG 54
BRAFTOVI122
BRUKINSA
<i>butalbital-acetaminop-caf-cod</i> 172
butalbital-acetaminophen oral tablet 50-
<i>325 mg</i> 172
butalbital-acetaminophen-caff oral
<i>capsule 50-325-40 mg</i> 172
butalbital-acetaminophen-caff oral tablet 172
butalbital-aspirin-caffeine172
CABLIVI INJECTION KIT
CABOMETYX ORAL TABLET 20
MG, 40 MG, 60 MG 59
CALQUENCE (ACALABRUTINIB
MAL)
CAPRELSA ORAL TABLET 100 MG,
300 MG451
carbinoxamine maleate oral liquid170
carbinoxamine maleate oral tablet 4 mg 170
<i>carglumic acid</i>
CAYSTON
CERDELGA115
chlorzoxazone oral tablet 250 mg, 500
<i>mg</i> , <i>750 mg</i>
CINQAIR
CINRYZE
clemastine oral tablet
codeine-butalbital-asa-caff172
<i>colchicine oral tablet</i> 72

COMETRIQ ORAL CAPSULE 100	
MG/DAY(80 MG X1-20 MG X1), 140	
MG/DAY(80 MG X1-20 MG X3), 60	
MG/DAY (20 MG X 3/DAY)	58
COPAXONE SUBCUTANEOUS	
SYRINGE 20 MG/ML, 40 MG/ML	. 160
COPIKTRA	. 105
CORTROPHIN GEL	73
COSENTYX (2 SYRINGES)	.362
COSENTYX INTRAVENOUS	
COSENTYX PEN (2 PENS)	
COSENTYX SUBCUTANEOUS	
SYRINGE 75 MG/0.5 ML	. 362
COSENTYX UNOREADY PEN	. 362
COTELLIC	
cyclobenzaprine oral tablet 10 mg, 5 mg.	. 188
cyproheptadine	
ĊYRAMZA	
CYSTADROPS	78
CYSTARAN	
dalfampridine	
DÁNÝELZA	
DARZALEX	
DARZALEX FASPRO	
DAURISMO ORAL TABLET 100	
MG, 25 MG	.159
deferasirox	
deferiprone	90
deferoxamine	
DIACOMIT ORAL CAPSULE 250	
MG, 500 MG	. 388
DIACOMIT ORAL POWDER IN	
PACKET 250 MG, 500 MG	. 388
diclofenac sodium topical gel 3 %	
diclofenac sodium topical solution in	
metered-dose pump	96
dimethyl fumarate oral capsule, delayed	
release(drlec) 120 mg, 120 mg (14)-240)
mg (46), 240 mg	97
diphenhydramine hcl oral elixir	. 189
diphenoxylate-atropine	
dipyridamole oral	
disopyramide phosphate oral capsule	.176
DOPTELET (10 TAB PACK)	
DOPTELET (15 TAB PACK)	

DOPTELET (30 TAB PACK)	32
dotti	177
dronabinol	. 100
droxidopa	. 101
DUAVEE	. 179
DUPIXENT PEN	102
DUPIXENT SYRINGE	102
EGRIFTA SV	405
ELFABRIO	. 309
ELIGARD	231
ELIGARD (3 MONTH)	. 231
ELIGARD (4 MONTH)	. 231
ELIGARD (6 MONTH)	
ELREXFIO 44 MG/1.1 ML VIAL	
OUTER, SUV, P/F	
ELREXFIO SUBCUTANEOUS	
SOLUTION 40 MG/ML	. 117
EMGALITY PEN	
EMGALITY SYRINGE	
SUBCUTANEOUS SYRINGE 120	
MG/ML, 300 MG/3 ML (100 MG/ML	
X 3)	.154
ENBREL	
ENBREL MINI	134
ENBREL SURECLICK	
ENDARI	. 237
ENSPRYNG	. 359
ENTADFI	. 146
EPCLUSA ORAL PELLETS IN	
PACKET 150-37.5 MG, 200-50 MG	. 377
EPCLUSA ORAL TABLET	. 377
EPIDIOLEX	62
EPKINLY	. 127
epoprostenol	. 130
ERBITUX	68
ERIVEDGE	. 458
ERLEADA ORAL TABLET 240 MG,	
60 MG	20
erlotinib oral tablet 100 mg, 150 mg, 25	
<i>mg</i>	
estradiol oral	
estradiol transdermal patch semiweekly.	177
estradiol transdermal patch weekly	177
estradiol-norethindrone acet oral tablet	
0.5-0.1 mg	178

everolimus (antineoplastic) oral table	
mg, 2.5 mg, 5 mg, 7.5 mg	137
everolimus (antineoplastic) oral table	t for
suspension	138
EVRYSDI	347
EXKIVITY	
EXONDYS-51	
FABRAZYME	
FARYDAK	
FASENRA	
FASENRA PEN	
fentanyl citrate buccal lozenge on a	
handle	142
FERRIPROX (2 TIMES A DAY)	
FERRIPROX ORAL SOLUTION.	
fingolimod	
FINTEPLA	
FOTIVDA	
FRUZAQLA ORAL CAPSULE 1 M	
5 MG	
FULPHILA	303
FYARRO	373
fyavolv	183
FYLNETRA	307
GALAFOLD	258
GAMIFANT	119
GATTEX 30-VIAL	400
GAVRETO	
gefitinib	156
GILENYA ORAL CAPSULE 0.25	MG
GILOTRIF	
GIVLAARI	
glatiramer subcutaneous syringe 20	150
mg/ml, 40 mg/ml	160
glatopa subcutaneous syringe 20 mg/n	
40 mg/ml	
glyburide	
glyburide micronized	
glyburide-metformin	
GRANIX	
HAEGARDA SUBCUTANEOUS	
RECON SOLN 2,000 UNIT, 3,000	
UNIT	57

HARVONI ORAL PELLETS IN	
PACKET 33.75-150 MG, 45-200 MG	. 225
HARVONI ORAL TABLET	225
HERCEPTIN HYLECTA	. 426
HERZUMA	. 430
HETLIOZ LQ	. 395
HUMIRA PEN	
HUMIRA PEN CROHNS-UC-HS	
START	10
HUMIRA PEN PSOR-UVEITS-ADO	L
HS	10
HUMIRA SUBCUTANEOUS	
SYRINGE KIT 40 MG/0.8 ML	10
HUMIRA(CF)	
HUMIRA(CF) PEDI CROHNS	
STARTER	10
HUMIRA(CF) PEN	
HUMIRA(CF) PEN CROHNS-UC-H	S.10
HUMIRA(CF) PEN PEDIATRIC UC	
HUMIRA(CF) PEN PSOR-UV-ADO	
HS	
IBRANCE	295
ibuprofen-famotidine	197
icatibant	. 198
ICLUSIG	322
IDHIFA	121
ILARIS (PF)	60
ILUMYA	
imatinib oral tablet 100 mg, 400 mg	. 200
IMBRUVICA ORAL CAPSULE 140	
MG, 70 MG	
IMBRUVICA ORAL SUSPENSION.	196
IMBRUVICA ORAL TABLET 140	
MG, 280 MG, 420 MG	196
IMJUDO	
IMLYGIC INJECTION SUSPENSIO	
10EXP6 (1 MILLION) PFU/ML	
IMPAVIDO	260
INBRIJA INHALATION CAPSULE,	,
W/INHALATION DEVICE	
indomethacin oral capsule 25 mg, 50 mg	191
indomethacin oral capsule, extended	
release	
INFLECTRA	
infliximab	202

INGREZZA45	50
INGREZZA INITIATION PACK 45	
	0
INLYTA ORAL TABLET 1 MG, 5	
MG	
INQOVI	\$7
INREBIC14	0
itraconazole oral solution21	5
IWILFIN10	
JAKAFI	
javygtor oral tablet, soluble)/
JAYPIRCA ORAL TABLET 100 MG,	
50 MG 32	20
JEMPERLI9	19
<i>jinteli</i>	33
JUXTAPID ORAL CAPSULE 10 MG,	
20 MG, 30 MG, 40 MG, 5 MG, 60 MG 24	12
JYNARQUE ORAL TABLET	.1
JYNARQUE ORAL TABLETS,	
SEQUENTIAL42	
KALYDECO21	6
KANJINTI42	27
KANUMA	50
KERENDIA14	
KESIMPTA PEN	
ketorolac injection solution 15 mg/ml, 30	5
mg/ml, 30 mg/ml (1 ml)18	,2
ketorolac injection syringe 15 mg/ml, 30	
<i>mg/ml</i> 18	52
ketorolac intramuscular solution	52
ketorolac intramuscular syringe	32
ketorolac oral18	
KEVZARA35	
KEYTRUDA	
KIMMTRAK	
KINERET	9
KISQALI FEMARA CO-PACK ORAL	
TABLET 200 MG/DAY(200 MG X 1)-	
2.5 MG, 400 MG/DAY(200 MG X 2)-	
2.5 MG, 600 MG/DAY(200 MG X 3)-	
2.5 MG	8
KISQALI ORAL TABLET 200	-
MG/DAY (200 MG X 1), 400 MG/DAY	
(200 MG X 2), 600 MG/DAY (200 MG	~
X 3)	
KORLYM25	,7

KOSELUGO ORAL CAPSULE 10	
MG, 25 MG	. 368
KRAZATI	9
KYNMOBI SUBLINGUAL FILM 10	
MG, 10-15-20-25-30 MG, 15 MG, 20	
MG, 25 MG, 30 MG	22
lanreotide	.222
lapatinib	223
lenalidomide	226
LENVIMA	
leuprolide (3 month)	.230
leuprolide subcutaneous kit	
lidocaine hcl mucous membrane solution -	
% (40 mg/ml)	
lidocaine topical adhesive patch, medicate	d
5 %	
lidocaine topical ointment	
lidocaine-prilocaine topical cream	240
LONSURF ORAL TABLET 15-6.14	
MG, 20-8.19 MG	
LOQTORZI	.423
LORBRENA ORAL TABLET 100	
MG, 25 MG	.245
LUMAKRAS ORAL TABLET 120	
MG, 320 MG	
LUNSUMIO	
LUPRON DEPOT	
LUPRON DEPOT (3 MONTH)	
LUPRON DEPOT (4 MONTH)	
LUPRON DEPOT (6 MONTH)	232
LUPRON DEPOT-PED	234
LUPRON DEPOT-PED (3 MONTH)	234
LYBALVI	
lyllana	177
LYNPARZA	
LYTGOBI	153
MARGENZA	
MAVENCLAD (10 TABLET PACK)	
MAVENCLAD (4 TABLET PACK)	
MAVENCLAD (5 TABLET PACK)	
MAVENCLAD (6 TABLET PACK)	
MAVENCLAD (7 TABLET PACK)	
MAVENCLAD (8 TABLET PACK)	
MAVENCLAD (9 TABLET PACK)	69
MAVYRET ORAL TABLET	. 161

MAYZENT ORAL TABLET 0.25 MG	,
1 MG, 2 MG	372
MAYZENT STARTER(FOR 1MG	
MAINT)	372
MAYZENT STARTER(FOR 2MG	
	372
megestrol oral suspension 400 mg/10 ml	
(40 mg/ml)	192
megestrol oral tablet	192
MEKINIST ORAL RECON SOLN	425
MEKINIST ORAL TABLET 0.5 MG,	
MG	
MEKTOVI	
MEPSEVII	
methocarbamol oral tablet 500 mg, 750	100
mg	188
mifepristone oral tablet 300 mg	
miglustat	
migustat mimvey	
modafinil oral tablet 100 mg, 200 mg	
morphine concentrate oral solution	
MOUNJARO	
MVASI	
	·· TU
NATPARA	.298
NATPARA NERLYNX	.298 270
NATPARA NERLYNX NEULASTA ONPRO	.298 270 .306
NATPARA NERLYNX NEULASTA ONPRO NINLARO	.298 270 .306 218
NATPARA NERLYNX NEULASTA ONPRO NINLARO <i>nitisinone</i>	.298 270 .306 218 278
NATPARA NERLYNX NEULASTA ONPRO NINLARO nitisinone NIVESTYM	.298 270 .306 218 278 143
NATPARA NERLYNX NEULASTA ONPRO NINLARO <i>nitisinone</i> NIVESTYM NORDITROPIN FLEXPRO	.298 270 .306 218 278 143
NATPARA NERLYNX NEULASTA ONPRO NINLARO NIVESTYM NORDITROPIN FLEXPRO norethindrone ac-eth estradiol oral tablet	.298 270 .306 218 278 143 .381
NATPARA NERLYNX NEULASTA ONPRO NINLARO NINLARO NIVESTYM NORDITROPIN FLEXPRO norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg	.298 270 .306 218 278 143 .381
NATPARA NERLYNX NEULASTA ONPRO NINLARO NINLARO NIVESTYM NORDITROPIN FLEXPRO norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg NOXAFIL ORAL SUSP,DELAYED	.298 270 .306 218 278 143 .381 183
NATPARA NERLYNX NEULASTA ONPRO NINLARO nitisinone NIVESTYM NORDITROPIN FLEXPRO norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg NOXAFIL ORAL SUSP,DELAYED RELEASE FOR RECON	.298 270 .306 218 278 143 .381 183 324
NATPARA NERLYNX NEULASTA ONPRO NINLARO NINLARO NIVESTYM NORDITROPIN FLEXPRO norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg NOXAFIL ORAL SUSP,DELAYED RELEASE FOR RECON NPLATE	.298 270 .306 218 278 143 .381 183 324 352
NATPARA NERLYNX NEULASTA ONPRO NINLARO NINLARO NIVESTYM NORDITROPIN FLEXPRO norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg NOXAFIL ORAL SUSP,DELAYED RELEASE FOR RECON NPLATE NUBEQA	.298 270 .306 218 278 143 .381 183 324 352 85
NATPARA NERLYNX NEULASTA ONPRO NINLARO NINLARO NIVESTYM NORDITROPIN FLEXPRO norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg NOXAFIL ORAL SUSP,DELAYED RELEASE FOR RECON NPLATE NUBEQA NUCALA SUBCUTANEOUS AUTO-	.298 270 .306 218 278 143 .381 183 324 352 85
NATPARA NERLYNX NEULASTA ONPRO NINLARO NINLARO NIVESTYM NORDITROPIN FLEXPRO norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg NOXAFIL ORAL SUSP,DELAYED RELEASE FOR RECON NPLATE NUBEQA NUCALA SUBCUTANEOUS AUTO- INJECTOR	.298 270 .306 218 278 143 .381 183 324 352 85 251
NATPARA NERLYNX NEULASTA ONPRO NINLARO NINLARO NIVESTYM NORDITROPIN FLEXPRO norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg NOXAFIL ORAL SUSP,DELAYED RELEASE FOR RECON NPLATE NUBEQA NUCALA SUBCUTANEOUS AUTO- INJECTOR NUCALA SUBCUTANEOUS RECON	.298 270 .306 218 278 143 .381 183 324 352 85 85
NATPARA NERLYNX NEULASTA ONPRO NINLARO NINLARO NIVESTYM NORDITROPIN FLEXPRO norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg NOXAFIL ORAL SUSP,DELAYED RELEASE FOR RECON NPLATE NUBEQA NUCALA SUBCUTANEOUS AUTO- INJECTOR NUCALA SUBCUTANEOUS RECON SOLN	.298 270 .306 218 278 143 .381 183 324 352 85 85
NATPARA NERLYNX NEULASTA ONPRO NINLARO NINLARO NIVESTYM NORDITROPIN FLEXPRO norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg NOXAFIL ORAL SUSP,DELAYED RELEASE FOR RECON NPLATE NUBEQA NUCALA SUBCUTANEOUS AUTO- INJECTOR NUCALA SUBCUTANEOUS RECON SOLN NUCALA SUBCUTANEOUS	.298 270 .306 218 278 143 .381 183 324 352 85 85
NATPARA NERLYNX NEULASTA ONPRO NINLARO NINLARO NIVESTYM NORDITROPIN FLEXPRO norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg NOXAFIL ORAL SUSP,DELAYED RELEASE FOR RECON NPLATE NUBEQA NUCALA SUBCUTANEOUS AUTO- INJECTOR NUCALA SUBCUTANEOUS RECON SOLN NUCALA SUBCUTANEOUS SYRINGE 100 MG/ML, 40 MG/0.4	.298 270 .306 218 278 143 .381 183 324 352 85 251 N 251
NATPARA NERLYNX NEULASTA ONPRO NINLARO NINLARO NIVESTYM NORDITROPIN FLEXPRO norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg NOXAFIL ORAL SUSP,DELAYED RELEASE FOR RECON NPLATE NUBEQA NUCALA SUBCUTANEOUS AUTO- INJECTOR NUCALA SUBCUTANEOUS RECON SOLN NUCALA SUBCUTANEOUS SYRINGE 100 MG/ML, 40 MG/0.4 ML	.298 270 .306 218 278 143 .381 183 324 352 85 251 N 251
NATPARA NERLYNX NEULASTA ONPRO NINLARO NINLARO NIVESTYM NORDITROPIN FLEXPRO norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg NOXAFIL ORAL SUSP,DELAYED RELEASE FOR RECON NPLATE NUBEQA NUCALA SUBCUTANEOUS AUTO- INJECTOR NUCALA SUBCUTANEOUS RECON SOLN NUCALA SUBCUTANEOUS SYRINGE 100 MG/ML, 40 MG/0.4	.298 270 .306 218 278 143 .381 183 324 352 85 251 N 251 .251 317

NYVEPRIA	301
OCALIVA	. 281
OCREVUS	. 282
ODOMZO	. 385
OFEV	
OGIVRI	.428
OGSIVEO	
OJJAARA	
OLUMIANT	36
ONGENTYS	
ONTRUZANT	.429
ONUREG	34
OPDIVO	.279
OPDUALAG	280
OPSUMIT	
ORENCIA	3
ORENCIA (WITH MALTOSE)	1
ORENCIA CLICKJECT	
ORFADIN ORAL SUSPENSION	. 278
ORGOVYX	
ORILISSA ORAL TABLET 150 MG,	
200 MG	.111
ORKAMBI ORAL GRANULES IN	
PACKET	. 246
ORKAMBI ORAL TABLET	.246
ORSERDU ORAL TABLET 345 MG,	
	110
ORSERDU ORAL TABLET 345 MG, 86 MG	110 23
ORSERDU ORAL TABLET 345 MG, 86 MG OTEZLA	110 23 23
ORSERDU ORAL TABLET 345 MG, 86 MG OTEZLA OTEZLA STARTER	110 23 23 .293
ORSERDU ORAL TABLET 345 MG, 86 MG OTEZLA OTEZLA STARTER oxandrolone	110 23 23 .293 . 247
ORSERDU ORAL TABLET 345 MG, 86 MG OTEZLA OTEZLA STARTER oxandrolone OXLUMO	110 23 23 .293 . 247
ORSERDU ORAL TABLET 345 MG, 86 MG OTEZLA OTEZLA STARTER oxandrolone OXLUMO oxycodone oral concentrate	110 23 23 .293 . 247
ORSERDU ORAL TABLET 345 MG, 86 MG OTEZLA OTEZLA STARTER oxandrolone OXLUMO oxycodone oral concentrate OZEMPIC SUBCUTANEOUS PEN	110 23 23 .293 . 247
ORSERDU ORAL TABLET 345 MG, 86 MG OTEZLA OTEZLA STARTER oxandrolone OXLUMO oxycodone oral concentrate OZEMPIC SUBCUTANEOUS PEN INJECTOR 0.25 MG OR 0.5 MG (2	110 23 23 .293 . 247
ORSERDU ORAL TABLET 345 MG, 86 MG OTEZLA OTEZLA STARTER oxandrolone OXLUMO oxycodone oral concentrate OZEMPIC SUBCUTANEOUS PEN INJECTOR 0.25 MG OR 0.5 MG (2 MG/3 ML), 0.25 MG OR 0.5 MG(2	110 23 23 .293 . 247 .169
ORSERDU ORAL TABLET 345 MG, 86 MG OTEZLA OTEZLA STARTER oxandrolone OXLUMO oxycodone oral concentrate OZEMPIC SUBCUTANEOUS PEN INJECTOR 0.25 MG OR 0.5 MG (2 MG/3 ML), 0.25 MG OR 0.5 MG(2 MG/1.5 ML), 1 MG/DOSE (4 MG/3	110 23 .293 .247 .169
ORSERDU ORAL TABLET 345 MG, 86 MG OTEZLA OTEZLA STARTER oxandrolone OXLUMO oxycodone oral concentrate OZEMPIC SUBCUTANEOUS PEN INJECTOR 0.25 MG OR 0.5 MG (2 MG/3 ML), 0.25 MG OR 0.5 MG(2 MG/1.5 ML), 1 MG/DOSE (4 MG/3 ML), 2 MG/DOSE (8 MG/3 ML)	110 23 .293 .247 .169 .164 .310
ORSERDU ORAL TABLET 345 MG, 86 MG OTEZLA OTEZLA STARTER oxandrolone OXLUMO oxycodone oral concentrate OZEMPIC SUBCUTANEOUS PEN INJECTOR 0.25 MG OR 0.5 MG (2 MG/3 ML), 0.25 MG OR 0.5 MG(2 MG/1.5 ML), 1 MG/DOSE (4 MG/3 ML), 2 MG/DOSE (8 MG/3 ML) PALYNZIQ	110 23 . 293 . 247 . 169 . 164 . 310 . 193
ORSERDU ORAL TABLET 345 MG, 86 MG OTEZLA OTEZLA STARTER oxandrolone OXLUMO oxycodone oral concentrate OZEMPIC SUBCUTANEOUS PEN INJECTOR 0.25 MG OR 0.5 MG (2 MG/3 ML), 0.25 MG OR 0.5 MG(2 MG/1.5 ML), 1 MG/DOSE (4 MG/3 ML), 2 MG/DOSE (8 MG/3 ML) PALYNZIQ paroxetine hcl oral suspension paroxetine hcl oral tablet pazopanib	110 23 .293 .247 .169 .164 .310 .193 .300
ORSERDU ORAL TABLET 345 MG, 86 MG OTEZLA OTEZLA STARTER oxandrolone OXLUMO oxycodone oral concentrate OZEMPIC SUBCUTANEOUS PEN INJECTOR 0.25 MG OR 0.5 MG (2 MG/3 ML), 0.25 MG OR 0.5 MG(2 MG/1.5 ML), 1 MG/DOSE (4 MG/3 ML), 2 MG/DOSE (8 MG/3 ML) PALYNZIQ paroxetine hcl oral suspension paroxetine hcl oral tablet	110 23 .293 .247 .169 .164 .310 .193 .300
ORSERDU ORAL TABLET 345 MG, 86 MG OTEZLA OTEZLA STARTER oxandrolone OXLUMO oxycodone oral concentrate OZEMPIC SUBCUTANEOUS PEN INJECTOR 0.25 MG OR 0.5 MG (2 MG/3 ML), 0.25 MG OR 0.5 MG(2 MG/1.5 ML), 1 MG/DOSE (4 MG/3 ML), 2 MG/DOSE (8 MG/3 ML) PALYNZIQ paroxetine hcl oral suspension paroxetine hcl oral tablet pazopanib PEGASYS PEMAZYRE	110 23 .293 .247 .169 .164 .310 .193 .193 .300 .308 .313
ORSERDU ORAL TABLET 345 MG, 86 MG OTEZLA OTEZLA STARTER oxandrolone OXLUMO oxycodone oral concentrate OZEMPIC SUBCUTANEOUS PEN INJECTOR 0.25 MG OR 0.5 MG (2 MG/3 ML), 0.25 MG OR 0.5 MG (2 MG/1.5 ML), 1 MG/DOSE (4 MG/3 ML), 2 MG/DOSE (8 MG/3 ML) PALYNZIQ paroxetine hcl oral suspension paroxetine hcl oral tablet pazopanib PEGASYS	110 23 .293 .247 .169 .164 .310 .193 .300 .308 .313 .314

PIQRAY ORAL TABLET 200	
MG/DAY (200 MG X 1), 250 MG/DA	Y
(200 MG X1-50 MG X1), 300 MG/DA	Y
(150 MG X 2)	16
pirfenidone oral capsule	318
pirfenidone oral tablet 267 mg, 534 mg,	
801 mg	
PLEGRIDY SUBCUTANEOUS PEN	
INJECTOR 125 MCG/0.5 ML, 63	
MCG/0.5 ML- 94 MCG/0.5 ML	.212
PLEGRIDY SUBCUTANEOUS	
SYRINGE 125 MCG/0.5 ML, 63	
MCG/0.5 ML- 94 MCG/0.5 ML	.212
POMALYST	. 321
posaconazole oral	323
PREMARIN ORAL	174
PREMPHASE	.180
PREMPRO	
PREVYMIS INTRAVENOUS	
SOLUTION 240 MG/12 ML, 480	
MG/24 ML	. 228
PREVYMIS ORAL	
PROMACTA ORAL POWDER IN	
PACKET 12.5 MG, 25 MG	.118
PROMACTA ORAL TABLET 12.5	
MG, 25 MG, 50 MG, 75 MG	. 118
promethazine injection solution	. 185
promethazine oral	. 185
promethazine rectal	185
promethegan	
pyrimethamine	
QINLOCK	. 344
quinine sulfate	328
QULIPTA	30
RADICAVA	106
RAVICTI	
REGRANEX	38
RELEUKO	144
RELISTOR ORAL	. 255
RELISTOR SUBCUTANEOUS	
SOLUTION	254
RELISTOR SUBCUTANEOUS	
SYRINGE 12 MG/0.6 ML, 8 MG/0.4	
ML	
RENFLEXIS	204

RETACRIT INJECTION SOLUTION	
10,000 UNIT/ML, 2,000 UNIT/ML,	
20,000 UNIT/2 ML, 20,000 UNIT/ML,	
3,000 UNIT/ML, 4,000 UNIT/ML,	
40,000 UNIT/ML	128
RETEVMO ORAL CAPSULE 40 MG,	
80 MG	
REVCOVI	
REZLIDHIA	
REZUROCK	
RIABNI	
RINVOQ	
RITUXAN HYCELA	
RIVFLOZA	
ROLVEDON	108
ROZLYTREK ORAL CAPSULE 100	100
MG, 200 MG.	123
ROZLYTREK ORAL PELLETS IN	
PACKET	
RUBRACA	
RUXIENCE	
RYBELSUS	
RYBREVANT	
RYDAPT	
sajazir	198
sapropterin oral tablet, soluble	357
SCEMBLIX ORAL TABLET 20 MG,	
40 MG	
scopolamine base	187
SEROSTIM SUBCUTANEOUS	
RECON SOLN 4 MG, 5 MG, 6 MG	383
SIGNIFOR	299
sildenafil (pulm.hypertension)	
intravenous	369
sildenafil (pulm.hypertension) oral tablet	
SIRTURO	
SKYRIZI	
sodium oxybate	
sodium phenylbutyrate oral tablet	
SOMATULINE DEPOT	270
SUBCUTANEOUS SYRINGE 120	
MG/0.5 ML, 60 MG/0.2 ML, 90 MG/0.1	3
ML	
SOMAVERT	
somaver i	
sor ajento	200

SPRAVATO	133
SPRYCEL ORAL TABLET 100 MG,	
140 MG, 20 MG, 50 MG, 70 MG, 80	
MG	
STELARA	
STIMUFEND	305
STIVARGA	
STRENSIQ	26
sunitinib malate	
SUNOSI	
SUPPRELIN LA	194
SYMDEKO	
SYMLINPEN 120	326
SYMLINPEN 60	326
SYMPAZAN	70
SYNAGIS	296
SYNAREL	264
SYNRIBO	287
TABRECTA	65
tadalafil (pulm. hypertension)	. 390
tadalafil oral tablet 2.5 mg, 5 mg	
TAFINLAR ORAL CAPSULE	79
TAFINLAR ORAL TABLET FOR	
SUSPENSION	80
TAGRISSO	292
TAKHZYRO SUBCUTANEOUS	
SOLUTION	221
TAKHZYRO SUBCUTANEOUS	
SYRINGE 150 MG/ML, 300 MG/2 M	L
(150 MG/ML)	221
TALTZ AUTOINJECTOR	219
TALTZ SYRINGE	. 219
TALVEY	394
TALZENNA	392
TASCENSO ODT	149
TASIGNA ORAL CAPSULE 150 MC	Э,
200 MG, 50 MG	271
tasimelteon	395
TAVALISSE	150
TAZVERIK	
TECENTRIQ	29
TECVAYLI	399
tencon	172
TEPEZZA	403
ТЕРМЕТКО	402

teriflunomide	
testosterone cypionate	407
testosterone enanthate	408
testosterone transdermal gel in metered	_
<i>dose pump 12.5 mgl 1.25 gram (1%),</i>	
20.25 mg/1.25 gram (1.62 %)	406
testosterone transdermal gel in packet 1	%
(25 mg/2.5gram), 1 % (50 mg/5 gram)	406
testosterone transdermal solution in	
metered pump wlapp	406
tetrabenazine	409
THALOMID	411
TIBSOVO	217
TIVDAK	413
TRACLEER ORAL TABLET FOR	
SUSPENSION	53
TRAZIMERA	
TRELSTAR INTRAMUSCULAR	
SUSPENSION FOR	
RECONSTITUTION	439
TREMFYA	168
treprostinil sodium	
tretinoin	
trientine oral capsule 250 mg	437
TRIKAFTA ORAL GRANULES IN	
PACKET, SEQUENTIAL	114
TRIKAFTA ORAL TABLETS,	
SEQUENTIAL	114
TRIPTODUR	
TRULICITY	163
TRUQAP	63
TRUSELTIQ	
TRUXIMA	
TUKYSA ORAL TABLET 150 MG,	50
MG	
TURALIO	
TYSABRI	267
TYVASO	
UBRELVY	
UDENYCA	
UDENYCA AUTOINJECTOR	
UDENYCA ONBODY	
UPTRAVI INTRAVENOUS	
	'

UPTRAVI ORAL TABLET 1,000
MCG, 1,200 MCG, 1,400 MCG, 1,600
MCG, 200 MCG, 400 MCG, 600 MCG,
800 MCG
UPTRAVI ORAL TABLETS, DOSE
PACK
VALCHLOR
VANFLYTA
VEGZELMA47
VELCADE
VENCLEXTA ORAL TABLET 10
MG, 100 MG, 50 MG453
VENCLEXTA STARTING PACK 453
VERQUVO
VERZENIO
vigabatrin
vigadrone
vigpoder
VIBIZIM
VINIZINI IIIO VITRAKVI ORAL CAPSULE 100
MG, 25 MG
VITRAKVI ORAL SOLUTION 224
VIZIMPRO81
VONJO294
voriconazole oral suspension for
reconstitution
VOSEVI
VOWST
VUMERITY
WELIREG
XADAGO
XALKORI ORAL CAPSULE
XALKORI ORAL PELLET 150 MG,
20 MG, 50 MG77
XELJANZ419
XELJANZ XR 419
XERMELO
XGEVA93
XIFAXAN ORAL TABLET 200 MG,
550 MG
XOLAIR SUBCUTANEOUS AUTO-
INJECTOR 300 MG/2 ML
XOLAIR SUBCUTANEOUS RECON
SOLN
5011,

XOLAIR SUBCUTANEOUS	
SYRINGE	288
XOSPATA	
XPOVIO ORAL TABLET 100	
MG/WEEK (50 MG X 2), 40	
MG/WEEK (40 MG X 1), 40MG	
TWICE WEEK (40 MG X 2), 60	
MG/WEEK (60 MG X 1), 60MG	
TWICE WEEK (120 MG/WEEK), 80	
MG/WEEK (40 MG X 2), 80MG	
TWICE WEEK (160 MG/WEEK) 3	366
XTANDI ORAL CAPSULE	125
XTANDI ORAL TABLET 40 MG, 80	
MG	125
XYOSTED	108
yargesa2	259
YERVOY	214
YONSA	7
ZARXIO	45
zebutal1	
ZEJULA ORAL CAPSULE	274
ZEJULA ORAL TABLET	274
ZELBORAF	152
ZIEXTENZO	304
ZIRABEV	49
ZOLADEX	67
ZTALMY	155
ZTLIDO	239
ZURZUVAE ORAL CAPSULE 20	
MG, 25 MG, 30 MG	
ZYDELIG	99
ZYKADIA	67
ZYNLONTA	244
ZYNYZ	336